

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
**WO 00/78953 A2**

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
5/10, C07K 14/47, 14/705, 16/18, 16/28, C12Q 1/68,  
A61K 38/17, G01N 33/50, A01K 67/027

(21) International Application Number: PCT/US00/16668

(22) International Filing Date: 16 June 2000 (16.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/139,923 17 June 1999 (17.06.1999) US  
60/148,177 10 August 1999 (10.08.1999) US  
60/149,357 18 August 1999 (18.08.1999) US  
60/162,287 28 October 1999 (28.10.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 60/139,923 (CIP)  
Filed on 17 June 1999 (17.06.1999)  
US 60/148,177 (CIP)  
Filed on 10 August 1999 (10.08.1999)  
US 60/149,357 (CIP)  
Filed on 18 August 1999 (18.08.1999)  
US 60/162,287 (CIP)  
Filed on 28 October 1999 (28.10.1999)

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YANG, Junming [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). HILLMAN, Jennifer, L.

[US/US]; 230 Monroe Drive, #12, Mountain View, CA 94040 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). BURFORD, Neil [GB/US]; 1308 4th Avenue, San Francisco, CA 94122 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rocksprings Drive, Hayward, CA 94545 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way, #1, Menlo Park, CA 94025 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 00/78953 A2

(54) Title: HUMAN TRANSPORT PROTEINS

(57) Abstract: The invention provides human transport proteins (TPPT) and polynucleotides which identify and encode TPPT. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of TPPT.

## HUMAN TRANSPORT PROTEINS

### TECHNICAL FIELD

5           This invention relates to nucleic acid and amino acid sequences of human transport proteins and to the use of these sequences in the diagnosis, treatment, and prevention of transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer.

### 10           BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded and subdivided into functionally distinct organelles by hydrophobic lipid bilayer membranes. These membranes act as a barrier to most molecules, and maintain the essential differences between the cytosol, the extracellular environment, and the contents of each intracellular organelle. Transport of essential nutrients, certain metal ions, metabolic waste  
15 products, cell signaling molecules, macromolecules, and proteins across lipid membranes and between organelles must be mediated by a variety of transport molecules. Transport between the cytoplasm and the extracellular environment, and between the cytoplasm and luminal spaces of cellular organelles requires specific transport proteins. Each transport protein carries a particular class of molecule, such as ions, sugars, or amino acids, and often is specific to a certain molecular  
20 species of the class.

Cells and organelles require transport molecules to import and export essential nutrients and metal ions including  $K^+$ ,  $NH_4^+$ ,  $P_i$ ,  $SO_4^{2-}$ , sugars, and vitamins, as well as various metabolic waste products. Transport proteins also play roles in antibiotic resistance, toxin secretion, ion balance, synaptic neurotransmission, kidney function, intestinal absorption, tumor growth, and other diverse  
25 cell functions (Griffith, J. and C. Sansom (1998) The Transporter Facts Book, Academic Press, San Diego CA, pp. 3-29). Transport can occur by a passive concentration-dependent mechanism, or can be linked to an energy source such as ATP hydrolysis or an ion gradient. Proteins that function in transport include carrier proteins, which bind to a specific solute and undergo a conformational change that transfers the bound solute across the membrane, and channel proteins, which form  
30 hydrophilic pores that allow specific solutes to diffuse through the membrane down an electrochemical solute gradient.

Transport proteins are often multi-pass transmembrane proteins, which either actively transport molecules across the membrane or passively allow them to cross. Active transport involves directional pumping of a solute across the membrane, usually against an electrochemical gradient.  
35 Active transport is tightly coupled to a source of metabolic energy, such as ATP hydrolysis or an electrochemically favorable ion gradient. Passive transport involves the movement of a solute down

its electrochemical gradient. Transport proteins can be further classified as either carrier proteins or channel proteins. Carrier proteins, which can function in active or passive transport, bind to a specific solute to be transported and undergo a conformational change which transfers the bound solute across the membrane. Channel proteins, which only function in passive transport, form hydrophilic pores across the membrane. When the pores open, specific solutes, such as inorganic ions, pass through the membrane and down the electrochemical gradient of the solute. Examples include facilitative transporters, the secondary active symporters and antiporters driven by ion gradients, and active ATP binding cassette transporters involved in multiple-drug resistance and targeting of antigenic peptides to MHC Class I molecules. Transported substrates range from nutrients and ions to a broad variety of drugs, peptides and proteins.

Information on the action of ARL-6 (ADP-ribosylation like factor), an endoplasmic reticulum transmembrane protein, can be found in Greenfield, J.J. and S. High (1999; J. Cell Sci. 112:1477-1486). Information on reduced folate carrier transporter proteins can be found in Dixon, K.H. et al. (1994; J. Biol. Chem. 269:17-20) and Moscow, J.A. et al. (1995; Cancer Res. 55:5983-5987).

Carrier proteins which transport a single solute from one side of the membrane to the other are called uniporters. In contrast, coupled transporters link the transfer of one solute with simultaneous or sequential transfer of a second solute, either in the same direction (symport) or in the opposite direction (antiport). For example, intestinal and kidney epithelia contain a variety of symporter systems driven by the sodium gradient that exists across the plasma membrane. Sodium moves into the cell down its electrochemical gradient and brings the solute into the cell with it. The sodium gradient that provides the driving force for solute uptake is maintained by the ubiquitous  $\text{Na}^+/\text{K}^+$  ATPase. Sodium-coupled transporters include the mammalian glucose transporter (SGLT1), iodide transporter (NIS), and multivitamin transporter (SMVT). All three transporters have twelve putative transmembrane segments, extracellular glycosylation sites, and cytoplasmically-oriented N- and C-termini.

Mitochondrial carrier proteins are transmembrane-spanning proteins which transport ions and charged metabolites between the cytosol and the mitochondrial matrix. Examples include the ADP, ATP carrier protein; the 2-oxoglutarate/malate carrier; the phosphate carrier protein; the brown fat uncoupling protein which transports protons from the cytosol into the matrix; the pyruvate carrier; the dicarboxylate carrier which transports malate, succinate, fumarate, and phosphate; the tricarboxylate carrier which transports citrate and malate; and the Grave's disease carrier protein, a protein recognized by IgG in patients with active Grave's disease, an autoimmune disorder resulting in hyperthyroidism (Stryer, L. (1995) Biochemistry, W.H. Freeman and Company, New York NY, p. 551; PROSITE PDOC00189 Mitochondrial energy transfer proteins signature; Online Mendelian Inheritance in Man (OMIM) \*275000 Graves Disease).

This class of transporters also includes the mitochondrial uncoupling proteins, which create

proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. The result is energy dissipation in the form of heat. Mitochondrial uncoupling proteins have been implicated as modulators of thermoregulation and metabolic rate, and have been proposed as potential targets for drugs against metabolic diseases such as obesity (Ricquier, D. et al. 5 (1999) *J. Int. Med.* 245:637-642).

A number of metal ions such as iron, zinc, copper, cobalt, manganese, molybdenum, selenium, nickel, and chromium are important as cofactors for a number of enzymes. For example, zinc is required for the function of enzymes such as the extracellular matrix metalloproteinases, and zinc ions stabilize several motifs commonly found in transcription factors, including zinc fingers, zinc 10 clusters, and LIM domains. Zinc and other metal ions must be provided in the diet, and are absorbed by transporters in the gastrointestinal tract. Plasma proteins transport the metal ions to the liver and other target organs, where specific transporters move the ions into cells and cellular organelles as needed. Imbalances in metal ion metabolism have been associated with a number of disease states (Danks, D.M. (1986) *J. Med. Genet.* 23:99-106).

15 The largest and most diverse family of transport proteins known are the ATP-binding cassette (ABC) transporters. As a family, ABC transporters can transport substances that differ markedly in chemical structure and size, ranging from small molecules such as ions, sugars, amino acids, peptides, and phospholipids, to lipopeptides, large proteins, and complex hydrophobic drugs. ABC proteins consist of four modules: two nucleotide-binding domains (NBD), which hydrolyze ATP to supply the 20 energy required for transport, and two membrane-spanning domains (MSD), each containing six putative transmembrane segments. These four modules may be encoded by a single gene, as is the case for the cystic fibrosis transmembrane regulator (CFTR), or by separate genes. When encoded by separate genes, each gene product contains a single NBD and MSD. These "half-molecules" form homo- and heterodimers, such as Tap1 and Tap2, the endoplasmic reticulum-based major 25 histocompatibility (MHC) peptide transport system. Several genetic diseases are attributed to defects in ABC transporters, such as the following diseases and their corresponding proteins: cystic fibrosis (CFTR, an ion channel), adrenoleukodystrophy (adrenoleukodystrophy protein, ALDP), Zellweger syndrome (peroxisomal membrane protein-70, PMP70), and hyperinsulinemic hypoglycemia (sulfonylurea receptor, SUR). Overexpression of the multidrug resistance (MDR) protein, another 30 ABC transporter, in human cancer cells makes the cells resistant to a variety of cytotoxic drugs used in chemotherapy (Taglicht, D. and S. Michaelis (1998) *Methods Enzymol.* 292:131-163).

The nuclear pore complex (NPC) is a large multiprotein complex spanning the nuclear envelope which mediates the transport of proteins and RNA molecules between the nucleus and the cytoplasm, thus contributing to the regulation of gene expression. The NPC allows passive diffusion 35 of ions, small molecules, and macromolecules under about 60kD, while larger macromolecules are transported by facilitated, energy-dependent pathways. Nuclear localization signals (NLS), consisting



of short stretches of amino acids enriched in basic residues, are found on proteins that are targeted to the nucleus, such as the glucocorticoid receptor. The NLS is recognized by the NLS receptor, importin, which then interacts with the monomeric GTP-binding protein Ran. This NLS protein/receptor/Ran complex navigates the nuclear pore with the help of the homodimeric protein nuclear transport factor 2 (NTF2) (Nakielnny, S. and G. Dreyfuss (1997) Curr. Opin. Cell Biol. 9:420-429; Gorlich, D. (1997) Curr. Opin. Cell Biol. 9:412-419). Four O-linked glycoproteins, p62, p58, p54, and p45, exist as a stable "p62 complex" that forms a ring localized on both nucleoplasmic and cytoplasmic surfaces of the NPC. The p62, p58, and p54 proteins all interact directly with the cytosolic transport factors p97 and NTF2, suggesting that the p62 complex is an important ligand binding site near the central gated channel of the NPC (Hu, T. et al. (1996) J. Cell Biol. 134:589-601).

Transport can also occur through intercellular bridges which connect the cytoplasms of sister cells, for example in the male and female germline of species ranging from fruit flies to humans. These bridges allow passage of cytoplasmic materials between cells during development. Intercellular bridges have also been found to connect somatic cells. The nurse cells and oocyte of a *Drosophila* egg chamber, which are derived from a single precursor cell through four rounds of mitosis, are connected to each other through intercellular bridges called ring canals. The cells do not completely separate after mitosis; the mitotic cleavage furrows are transformed into ring canals by the addition of an actin cytoskeleton lining the tunnels between the cells. The *Drosophila* kelch protein functions in organizing actin in the ring canal. Mutations in kelch cause female sterility in *Drosophila*. Kelch contains four protein domains: the NTR domain at the N-terminus, the BTB or POZ domain, the IVR or intervening region; and the kelch repeat domain, which contains six 50-amino acid kelch repeats. The BTB or POZ domain, a 120-amino acid motif that is also found in several zinc-finger containing transcription factors, may be important in dimerization of kelch. Kelch repeats are found in other proteins as well and may be important for actin binding (Robinson, D.N. and L. Cooley (1997) J. Cell Biol. 138:799-810; Cooley, L. (1998) Cell 93:913-915).

### Ion Channels

The electrical potential of a cell is generated and maintained by controlling the movement of ions across the plasma membrane. The movement of ions requires ion channels, which form an ion-selective pore within the membrane. Ion channels share common structural and mechanistic themes. The channel consists of four or five subunits or protein monomers that are arranged like a barrel in the plasma membrane. Each subunit typically consists of six potential transmembrane segments (S1, S2, S3, S4, S5, and S6). The center of the barrel forms a pore lined by  $\alpha$ -helices or  $\beta$ -strands. The side chains of the amino acid residues comprising the  $\alpha$ -helices or  $\beta$ -strands establish the charge (cation or anion) selectivity of the channel. The degree of selectivity, or what specific ions are allowed to pass through the channel, depends on the diameter of the narrowest part of the pore. There

are two basic types of ion channels, ion transporters and gated ion channels. Ion transporters utilize the energy obtained from ATP hydrolysis to actively transport an ion against the ion's concentration gradient. Gated ion channels allow passive flow of an ion down the ion's electrochemical gradient under restricted conditions. Together, these types of ion channels generate, maintain, and utilize an electrochemical gradient that is used in 1) electrical impulse conduction down the axon of a nerve cell, 2) transport of molecules into cells against concentration gradients, 3) initiation of muscle contraction, and 4) endocrine cell secretion.

Transmembrane ATPases are divided into three families. The phosphorylated (P) class ion transporters, including  $\text{Na}^+\text{-K}^+$  ATPase,  $\text{Ca}^{2+}$  ATPase,  $\text{H}^+$  ATPase, and  $\text{Cu}^{++}$  ATPase, are activated by a phosphorylation event. P-class ion transporters are responsible for maintaining resting potential distributions such that cytosolic concentrations of  $\text{Na}^+$  and  $\text{Ca}^{2+}$  are low and cytosolic concentration of  $\text{K}^+$  is high. The vacuolar (V) class of ion transporters include  $\text{H}^+$  pumps on intracellular organelles, such as lysosomes and Golgi. V-class ion transporters are responsible for generating the low pH within the lumen of these organelles that is required for function. The coupling factor (F) class consists of  $\text{H}^+$  pumps in the mitochondria. F-class ion transporters utilize a proton gradient to generate ATP from ADP and inorganic phosphate ( $\text{P}_i$ ).

$\text{Cu}^{++}$  ATPases export copper from cells (PROSITE PDOC00139 E1-E2 ATPases phosphorylation site). Mutations in one  $\text{Cu}^{++}$  ATPase cause Wilson disease, in which toxic amounts of copper accumulate in a number of organs, particularly the liver and brain (Tanzi, R.E. et al. (1993) Nat. Genet. 5:344-350). Mutations in another  $\text{Cu}^{++}$  ATPase cause Menkes disease and occipital horn syndrome. Menkes disease mutations block export of copper from the gastrointestinal tract, leading to skeletal abnormalities, severe mental retardation, neurologic degeneration, and mortality in early childhood (Harrison, M.D. and C.T. Dameron (1999) J. Biochem. Mol. Toxicol. 13:93-106). Occipital horn syndrome mutations cause connective tissue defects (Harrison, supra; Levinson, B. et al. (1996) Hum. Mol. Genet. 5:1737-1742).

The coupling factor (F) class of ion transporters consists of  $\text{H}^+$  pumps in mitochondria, chloroplasts, and bacteria. For example, the  $\text{F}_0\text{F}_1$  ATPase utilizes a proton gradient across the inner mitochondrial membrane to generate ATP from ADP and inorganic phosphate ( $\text{P}_i$ ). The  $\text{F}_0\text{F}_1$  ATPase is composed of the  $\text{F}_0$  complex, which is the transmembrane channel through which protons flow, and the  $\text{F}_1$  complex, where ATP synthesis activity resides.  $\text{F}_0$  has three subunits, A (also known as protein 6), B, and C (Lodish, H. et al. (1995) Molecular Cell Biology, Scientific American Books, New York NY, pp. 752-756; PROSITE PDOC00420 ATP synthase a subunit signature).

Voltage-gated  $\text{Ca}^{2+}$  channels are involved in presynaptic neurotransmitter release, and heart and skeletal muscle contraction. The voltage-gated  $\text{Ca}^{2+}$  channels from skeletal muscle (L-type) and brain (N-type) have been purified and, though their functions differ dramatically, they have similar subunit compositions. The channels are composed of three subunits. The  $\alpha_1$  subunit forms the

membrane pore and voltage sensor, while the  $\alpha_2\delta$  and  $\beta$  subunits modulate the voltage-dependence, gating properties, and the current amplitude of the channel. These subunits are encoded by at least six  $\alpha_1$ , one  $\alpha_2\delta$ , and four  $\beta$  genes. A fourth subunit,  $\gamma$ , has been identified in skeletal muscle (Walker, D. et al. (1998) J. Biol. Chem. 273:2361-2367; and Jay, S.D. et al. (1990) Science 248:490-492). The human  $\beta 4$  subunit is homologous to the mouse epilepsy gene lethargic, and is a candidate for involvement in neurological disorders including ataxia and absence epilepsy (Escayg, A. et al. (1998) Genomics 50:14-22).

Ligand-gated channels open their pores when an extracellular or intracellular mediator binds to the channel. Neurotransmitter-gated channels are channels that open when a neurotransmitter binds to their extracellular domain. These channels exist in the postsynaptic membrane of nerve or muscle cells. There are two types of neurotransmitter-gated channels. Sodium channels open in response to excitatory neurotransmitters, such as acetylcholine, glutamate, and serotonin. This opening causes an influx of  $\text{Na}^+$  and produces the initial localized depolarization that activates the voltage-gated channels and starts the action potential. Chloride channels open in response to inhibitory neurotransmitters, such as  $\gamma$ -aminobutyric acid (GABA) and glycine, leading to hyperpolarization of the membrane and the subsequent generation of an action potential.

Ion channels are expressed in a number of tissues where they are implicated in a variety of processes. CNG channels, while abundantly expressed in photoreceptor and olfactory sensory cells, are also found in kidney, lung, pineal, retinal ganglion cells, testis, aorta, and brain. Calcium-activated  $\text{K}^+$  channels may be responsible for the vasodilatory effects of bradykinin in the kidney and for shunting excess  $\text{K}^+$  from brain capillary endothelial cells into the blood. They are also implicated in repolarizing granulocytes after agonist-stimulated depolarization (Ishi, T.M. et al. (1997) Proc. Natl. Acad. Sci. USA 94:11651-11656). Another transmembrane protein, the leukotriene B<sub>4</sub> receptor (BLT) appears to be involved in inflammation responses and host cell defense against infection. BLT also functions as an HIV coreceptor (Izumi, T. et al. (1997) Nature 387:620-624; Martin, V. et al. (1999) J. Biol. Chem. 274:8597-8603).

Ion channels have been the target for many drug therapies. Neurotransmitter-gated channels have been targeted in therapies for treatment of insomnia, anxiety, depression, and schizophrenia. Voltage-gated channels have been targeted in therapies for arrhythmia, ischemic stroke, head trauma, and neurodegenerative disease (Taylor, C.P. and L.S. Narasimhan (1997) Adv. Pharmacol. 39:47-98).

$\text{K}^+$  channels are located in all cell types, and may be regulated by voltage, ATP concentration, or second messengers such as  $\text{Ca}^{++}$  and cAMP. In non-excitable tissue,  $\text{K}^+$  channels are involved in protein synthesis, control of endocrine secretions, and the maintenance of osmotic equilibrium across membranes. In neurons and other excitable cells, in addition to regulating action potentials and repolarizing membranes,  $\text{K}^+$  channels are responsible for setting resting membrane potential. The cytosol contains non-diffusible anions and, to balance this net negative charge, the cell

contains a  $\text{Na}^+$ - $\text{K}^+$  pump and ion channels that provide the redistribution of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cl}^-$ . The pump actively transports  $\text{Na}^+$  out of the cell and  $\text{K}^+$  into the cell in a 3:2 ratio. Ion channels in the plasma membrane allow  $\text{K}^+$  and  $\text{Cl}^-$  to flow by passive diffusion. Because of the high negative charge within the cytosol,  $\text{Cl}^-$  flows out of the cell. The flow of  $\text{K}^+$  is balanced by an electromotive force  
5 pulling  $\text{K}^+$  into the cell, and a  $\text{K}^+$  concentration gradient pushing  $\text{K}^+$  out of the cell. Thus, the resting membrane potential is primarily regulated by  $\text{K}^+$  flow (Salkoff, L. and T. Jegla (1995) *Neuron* 15:489-492). Information on NY-REN-45, a  $\text{K}^+$  channel integral membrane protein, can be found in Scanlan, M.J. et al. (1998; *Int. J. Cancer* 76:652-658). The emopamil-binding protein (EBP) shares structural features with both pro- and eukaryotic drug transport proteins (Hanner, M. et al. (1995) *J.*  
10 *Biol. Chem.* 270:7551-7557). The  $\text{Na}^+$  channel, transmembrane protein myelin protein zero (MPZ) may be responsible for some sporadic cases of Dejerine-Scottas disease (hereditary motor and sensory neuropathy type III) (Hayasaka, K. et al. (1993) *Nat. Genet.* 5:266-268).

$\text{K}^+$  pore-forming subunits generally have six transmembrane-spanning domains with a short region between the fifth and sixth transmembrane regions that senses membrane potential; and the  
15 amino and carboxy termini are located intracellularly. In mammalian heart, the duration of ventricular action potential is controlled by a  $\text{K}^+$  current. Thus, the  $\text{K}^+$  channel is central to the control of heart rate and rhythm.  $\text{K}^+$  channel dysfunctions are associated with a number of renal diseases including hypertension, hypokalemia, and the associated Bartter's syndrome and Getelman's syndrome, as well as neurological disorders including epilepsy.  $\text{K}^+$  channels have been implicated in  
20 Alzheimer's disease by observations that a significant component of senile plaques, beta amyloid or A beta, also blocks voltage-gated potassium channels in hippocampal neurons (Antes, L.M. et al. (1998) *Seminars Nephrol.* 18:31-45; Stoffel, M. and L.Y. Jan (1998) *Nat. Genet.* 18:6-8; Madeja, M. et al. (1997) *Eur. J. Neurosci.* 9:390-395; Good, T.A. et al. (1996) *Biophys. J.* 70:296-304).

Gated ion channels control ion flow by regulating the opening and closing of pores. These  
25 channels are categorized according to the manner of regulating the gating function. Mechanically-gated channels open pores in response to mechanical stress, voltage-gated channels open pores in response to changes in membrane potential, and ligand-gated channels open pores in the presence of a specific ion, nucleotide, or neurotransmitter.

Voltage-gated  $\text{Na}^+$  channels are responsible for electrical excitability of neurons, skeletal  
30 muscle, heart, and neuroendocrine tissues. For example, the sequential opening and closing of voltage-gated  $\text{Na}^+$  channels results in the propagation of action potentials down neuronal axons.  $\text{Na}^+$  channels isolated from rat brain tissue are heterotrimeric complexes composed of a 260 kDa pore forming  $\alpha$  subunit that associates with two smaller auxiliary subunits,  $\beta 1$  and  $\beta 2$ . The  $\beta 2$  subunit is an integral membrane glycoprotein that contains an extracellular Ig domain, and its association with  $\alpha$   
35 and  $\beta 1$  subunits correlates with increased function of the channel, a change in the channel's gating properties, as well as an increase in whole cell capacitance (Isom, L.L. et al. (1995) *Cell* 83:433-442).

### Integral Membrane Proteins

The majority of known integral membrane proteins are transmembrane proteins (TM) which are characterized by an extracellular, a transmembrane, and an intracellular domain. TM domains are typically comprised of 15 to 25 hydrophobic amino acids which are predicted to adopt an  $\alpha$ -helical conformation. TM proteins are classified as bitopic (Types I and II) and polytopic (Types III and IV) (Singer, S.J. (1990) *Annu. Rev. Cell Biol.* 6:247-96). Bitopic proteins span the membrane once while polytopic proteins contain multiple membrane-spanning segments. TM proteins that act as cell-surface receptor proteins involved in signal transduction include growth and differentiation factor receptors, and receptor-interacting proteins such as *Drosophila* pecanex and frizzled proteins, LIV-1 protein, NF2 protein, and GNS1/SUR4 eukaryotic integral membrane proteins. TM proteins also act as transporters of ions or metabolites, such as gap junction channels (connexins) and ion channels, and as cell anchoring proteins, such as lectins, integrins, and fibronectins. TM proteins act as vesicle organelle-forming molecules, such as calveolins, or as cell recognition molecules, such as cluster of differentiation (CD) antigens, glycoproteins, and mucins. Information on connexin can be found in Kanter, H.L. et al. (1994; *J. Mol. Cell. Cardiol.* 26:861-868).

Many membrane proteins (MPs) contain amino acid sequence motifs that target these proteins to specific subcellular sites. Examples of these motifs include PDZ domains, KDEL, RGD, NGR, and GSL sequence motifs, von Willebrand factor A (vWFA) domains, and EGF-like domains. RGD, NGR, and GSL motif-containing peptides have been used as drug delivery agents in cancer treatments which target tumor vasculature (Arap, W. et al. (1998) *Science*, 279:377-380.) Furthermore, MPs may also contain amino acid sequence motifs, such as the carbohydrate recognition domain (CRD), also known as the C-type lectin domain, that mediate interactions with extracellular or intracellular molecules.

G-protein coupled receptors (GPCR) comprise a superfamily of integral membrane proteins which transduce extracellular signals. GPCRs include receptors for biogenic amines, lipid mediators of inflammation, peptide hormones, and sensory signal mediators. The structure of these highly-conserved receptors consists of seven hydrophobic transmembrane regions, an extracellular N-terminus, and a cytoplasmic C-terminus. Three extracellular loops alternate with three intracellular loops to link the seven transmembrane regions. The most conserved parts of these proteins are the transmembrane regions and the first two cytoplasmic loops. Cysteine disulfide bridges connect the second and third extracellular loops. A conserved, acidic-Arg-aromatic residue triplet present in the second cytoplasmic loop may interact with G proteins. A GPCR consensus pattern is characteristic of most proteins belonging to this superfamily (ExPASy PROSITE document PS00237; and Watson, S. and S. Arkinstall (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp 2-6). Mutations and changes in transcriptional activation of GPCR-encoding genes have been

associated with neurological disorders such as schizophrenia, Parkinson's disease, Alzheimer's disease, drug addiction, and feeding disorders.

Cytochromes are electron-transferring proteins that contain a heme prosthetic group, a porphyrin ring containing a tightly bound iron atom. Cytochromes act as oxidoreductases in such  
 5 diverse cellular processes as respiration, photosynthesis, fatty acid metabolism, and neurotransmitter biosynthesis. The heme iron atom serves as the actual electron carrier by changing from the ferric to the ferrous oxidation state when accepting an electron. Cytochromes accept electrons from one substrate such as NADH or ascorbate and donate them to other electron carriers such as other cytochromes, ubiquinone, or semidehydroascorbic acid (Lodish, H. et al. (1995) Molecular Cell  
 10 Biology, Scientific American Books, New York NY, pp. 759-770, 786-797; Sperling, P. et al. (1995) Eur. J. Biochem. 232:798-805; and Online Mendelian Inheritance in Man (OMIM) \*600019 Cytochrome b561, CYB561).

Cytochrome b5 is an electron donor in membrane-linked redox enzyme systems involved in lipid and drug metabolism. Cytochrome b5 has been found in Golgi, plasma, outer mitochondrial,  
 15 endoplasmic reticulum (ER), and microbody membranes. Conserved amino acids in cytochrome b5 include eight invariant amino acids at W34, H51, P52, G53, G54, G63, F70, and H74, and fifteen conserved amino acids at L24, I35, S36, V41, Y42, N43, T45, W47, A48, L58, D65, T67, L85, T87, and G88 (numbering based on the sunflower cytochrome b5/delta-6 desaturase fusion protein; GI 1040729, Sperling, supra). The invariant residues H51PGG are involved in heme-binding.  
 20 Cytochrome b5-like domains have also been found linked to other enzymes. For example, cytochrome b5-like domains are part of delta-9 fatty acid desaturases in yeast and Histoplasma capsulatum, nitrate reductase, sulfite reductase, flavocytochrome b2, Arabidopsis thaliana acyl lipid desaturase, and Borago officinalis (borage) and Helianthus annuus (sunflower) delta-6 desaturases (Sperling, supra; Sayanova, O. et al (1997) Proc. Natl. Acad. Sci. USA 94:4211-4216; and Mitchell,  
 25 A.G. and C.E. Martin (1997) J. Biol. Chem. 272:28281-28288).

Signal peptides are found on proteins that are targeted to the endoplasmic reticulum (ER). Signal peptides consist of stretches of amino acids enriched in hydrophobic residues. Signal peptides are usually found at the extreme N-terminus of the protein and are recognized by a cytosolic signal-recognition peptide (SRP). The SRP binds to the signal peptide and to an SRP receptor, an integral  
 30 membrane protein in the ER. Once bound to the SRP receptor, the newly formed protein containing the signal peptide is translocated across the ER membrane. Proteins containing signal peptides may end up inserted into the lipid bilayer, or they may end up in the lumen of an organelle or secreted from the cell.

### 35 Disease Correlation

The etiology of numerous human diseases and disorders can be attributed to defects in the

transport of molecules across membranes. Defects in the trafficking of membrane-bound transporters and ion channels are associated with several disorders, e.g. cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, von Gierke disease, and certain forms of diabetes mellitus. Single-gene defect diseases resulting in an inability to transport small molecules across membranes include, e.g., cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease (van't Hoff, W.G. (1996) *Exp. Nephrol.* 4:253-262; Talente, G.M. et al. (1994) *Ann. Intern. Med.* 120:218-226; and Chillon, M. et al. (1995) *New Engl. J. Med.* 332:1475-1480).

Cystinuria is an inherited disease that results from the inability to transport cystine, the disulfide-linked dimer of cysteine, from the urine into the blood. Accumulation of cystine in the urine leads to the formation of cystine stones in the kidneys.

Transthyretin (TTR), present in human plasma, binds to and transports the thyroid hormone thyroxine. Mutations in TTR result in the conversion of TTR to amyloid, an insoluble fibrillar structure. The resulting amyloid plaques have been shown to be the causative agent in the development of familial amyloid polyneuropathy and senile systemic amyloidosis (Miroy, G.J. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:15051-15056).

Stomatin, a 31-kDa erythrocyte integral membrane protein has been linked to the hereditary anemia stomatocytosis. This anemia is characterized by red blood cells that lack stomatin and leak Na<sup>+</sup> and K<sup>+</sup>. Thus, stomatin is presumed to play a role in the regulation of ion transport. Red blood cell ion transport defects are also linked to other disorders such as hypertension (Stewart, G.W. (1997) *Int. J. Biochem. Cell Biol.* 29:271-274).

The discovery of new human transport proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer.

### SUMMARY OF THE INVENTION

The invention features purified polypeptides, human transport proteins, referred to collectively as "TPPT" and individually as "TPPT-1," "TPPT-2," "TPPT-3," "TPPT-4," "TPPT-5," "TPPT-6," "TPPT-7," "TPPT-8," "TPPT-9," "TPPT-10," "TPPT-11," "TPPT-12," "TPPT-13," "TPPT-14," "TPPT-15," "TPPT-16," "TPPT-17," "TPPT-18," "TPPT-19," "TPPT-20," "TPPT-21," "TPPT-22," "TPPT-23," "TPPT-24," "TPPT-25," "TPPT-26," "TPPT-27," "TPPT-28," "TPPT-29," "TPPT-30," "TPPT-31," "TPPT-32," "TPPT-33," "TPPT-34," "TPPT-35," "TPPT-36," "TPPT-37," "TPPT-38," "TPPT-39," "TPPT-40," "TPPT-41," "TPPT-42," and "TPPT-43." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-

43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-43.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-43. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:44-86.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a



polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence  
5 selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70%  
10 sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a  
15 sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary  
20 to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said  
25 hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70%  
30 sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting  
35 the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and a pharmaceutically acceptable excipient. In one embodiment, the pharmaceutical composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional TPPT, comprising administering to a patient in

need of such treatment the pharmaceutical composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:44-86, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

30

#### BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding TPPT.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of TPPT.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

5 Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding TPPT were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

10

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing  
15 particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a  
20 reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be  
25 used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

### 30 DEFINITIONS

"TPPT" refers to the amino acid sequences of substantially purified TPPT obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of  
35 TPPT. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TPPT either by directly interacting with

TPPT or by acting on components of the biological pathway in which TPPT participates.

An "allelic variant" is an alternative form of the gene encoding TPPT. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or  
5 many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding TPPT include those sequences with deletions,  
10 insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as TPPT or a polypeptide with at least one functional characteristic of TPPT. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding TPPT, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding  
15 TPPT. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent TPPT. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of TPPT is retained. For example,  
20 negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

25 The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

30 "Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of TPPT. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small  
35 molecules, or any other compound or composition which modulates the activity of TPPT either by directly interacting with TPPT or by acting on components of the biological pathway in which TPPT

participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind TPPT polypeptides can be prepared using intact polypeptides or using fragments  
5 containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

10 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen  
15 used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified  
20 sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or  
25 translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic TPPT, or of any oligopeptide thereof,  
30 to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

35 A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or

amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding TPPT or fragments of TPPT may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be  
 5 deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from  
 10 one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least  
 15 interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
20	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
25	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
30	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
35	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

40 Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of  
5 hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

10 A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of TPPT or the polynucleotide encoding TPPT which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a  
15 fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected  
20 from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:44-86 comprises a region of unique polynucleotide sequence that  
25 specifically identifies SEQ ID NO:44-86, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:44-86 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:44-86 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:44-86 and the region of SEQ ID NO:44-86 to which the fragment corresponds are routinely  
30 determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-43 is encoded by a fragment of SEQ ID NO:44-86. A fragment of SEQ ID NO:1-43 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-43. For example, a fragment of SEQ ID NO:1-43 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-43. The precise length of  
35 a fragment of SEQ ID NO:1-43 and the region of SEQ ID NO:1-43 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended



purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

5 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps  
10 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of  
15 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent  
20 similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at  
25 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The  
30 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

35 *Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

5 *Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous  
10 nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes  
15 in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some  
20 alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e  
25 sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

30 Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

35 *Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C<sub>0</sub>t or R<sub>0</sub>t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of TPPT which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of TPPT which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of TPPT. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of TPPT.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide,

polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an TPPT may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of TPPT.

"Probe" refers to nucleic acid sequences encoding TPPT, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs

can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is

expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,  
5 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

10 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic  
15 acids encoding TPPT, or fragments thereof, or TPPT itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or  
20 synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

25 The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides  
30 by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

35 A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

10 A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide



polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having  
 5 at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

## 10 THE INVENTION

The invention is based on the discovery of new human transport proteins (TPPT), the polynucleotides encoding TPPT, and the use of these compositions for the diagnosis, treatment, or prevention of transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer.

15 Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding TPPT. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each TPPT were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA  
 20 libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each TPPT and are useful as fragments in hybridization technologies.

25 The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical  
 30 methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding TPPT. The first column of Table 3 lists the nucleotide  
 35 SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:44-86

and to distinguish between SEQ ID NO:44-86 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express TPPT as a fraction of total tissues expressing TPPT. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing TPPT as a fraction of total tissues expressing TPPT. Column 5 lists the vectors used to subclone each cDNA library.

Of particular interest is the expression of SEQ ID NO:50 exclusively in cardiovascular tissue, the expression of SEQ ID NO:56 in nervous and gastrointestinal tissues, the expression of SEQ ID NO:57 in gastrointestinal tissues, and the expression of SEQ ID NO:66 in nervous system tissues. Of particular note is the tissue-specific expression of SEQ ID NO:75. Over 71% of the cDNA libraries expressing SEQ ID NO:75 are derived from lung tissue.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding TPPT were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:44 maps to chromosome 7 within the interval from 38.80 to 42.10 centiMorgans. SEQ ID NO:48 maps to chromosome X within the interval from 107.90 to 122.80 centiMorgans. SEQ ID NO:60 maps to chromosome 2 within the interval from 157.0 to 167.0 centiMorgans. SEQ ID NO:65 maps to chromosome 2 within the interval from 17.4 to 40.7 centiMorgans and to chromosome 5 within the interval from 61.1 to 69.6 centiMorgans. The interval on chromosome 5 from 61.1 to 69.6 centiMorgans also contains genes associated with Cockayne syndrome. SEQ ID NO:69 maps to chromosome 3 within the interval from 157.40 to 162.00 centiMorgans. SEQ ID NO:70 maps to chromosome 3 within the interval from 176.40 to 179.80 centiMorgans. SEQ ID NO:71 maps to chromosome 18 within the interval from the p-terminus to 52.30 centiMorgans. SEQ ID NO:73 maps to chromosome 17 within the interval from 75.70 to 84.20 centiMorgans, and to chromosome 2 within the interval from 204.70 to 209.30 centiMorgans. SEQ ID NO:76 maps to chromosome 20 within the interval from 79.00 to 94.40 centiMorgans. SEQ ID NO:80 maps to chromosome 18 within the interval from 1.60 to 6.20 centiMorgans, and to chromosome 11 within the interval from 117.90 to 126.00 centiMorgans. SEQ ID NO:83 maps to chromosome 17 within the interval from 67.60 to 69.30 centiMorgans, and from 83.8 centiMorgans to the q-terminus, and to chromosome 7 within the interval from 105.20 to 114.50 centiMorgans.

The invention also encompasses TPPT variants. A preferred TPPT variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the TPPT amino acid sequence, and which contains at least one functional or structural characteristic of TPPT.

The invention also encompasses polynucleotides which encode TPPT. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected

from the group consisting of SEQ ID NO:44-86, which encodes TPPT. The polynucleotide sequences of SEQ ID NO:44-86, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

5           The invention also encompasses a variant of a polynucleotide sequence encoding TPPT. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding TPPT. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:44-  
10   86 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:44-86. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of TPPT.

          It will be appreciated by those skilled in the art that as a result of the degeneracy of the  
15   genetic code, a multitude of polynucleotide sequences encoding TPPT, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the  
20   polynucleotide sequence of naturally occurring TPPT, and all such variations are to be considered as being specifically disclosed.

          Although nucleotide sequences which encode TPPT and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring TPPT under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding TPPT or  
25   its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding TPPT and its derivatives without altering the encoded amino acid sequences  
30   include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

          The invention also encompasses production of DNA sequences which encode TPPT and TPPT derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems  
35   using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding TPPT or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:44-86 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 5 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE 10 Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI 15 CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 20 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding TPPT may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, 25 restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids* 30 *Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences 35 are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo

Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in  
5 length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T)  
10 library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-  
15 specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments  
20 which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode TPPT may be cloned in recombinant DNA molecules that direct expression of TPPT, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally  
25 equivalent amino acid sequence may be produced and used to express TPPT.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter TPPT-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic  
30 oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number  
35 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or

improve the biological properties of TPPT, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired  
5 properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of  
10 homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding TPPT may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, TPPT itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved  
15 using the ABI 431A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of TPPT, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid  
25 chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active TPPT, the nucleotide sequences encoding TPPT or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains  
30 the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding TPPT. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding TPPT. Such signals  
35 include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding TPPT and its initiation codon and upstream regulatory sequences are inserted into

the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both

5 natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding TPPT and appropriate transcriptional and translational control

10 elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences

15 encoding TPPT. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or

20 animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984)

25 *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences

30 to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al., (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

35 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding TPPT. For example, routine cloning,

subcloning, and propagation of polynucleotide sequences encoding TPPT can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPT1 plasmid (Life Technologies). Ligation of sequences encoding TPPT into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of TPPT are needed, e.g. for the production of antibodies, vectors which direct high level expression of TPPT may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of TPPT. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of TPPT. Transcription of sequences encoding TPPT may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding TPPT may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses TPPT in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet.



15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of TPPT in cell lines is preferred. For example, sequences encoding TPPT can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous  
5 expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue  
10 culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk<sup>-</sup>* and *ap<sup>r</sup>-* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or  
15 herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which  
20 alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system.  
25 (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding TPPT is inserted within a marker gene sequence, transformed cells containing sequences encoding TPPT can be identified by the absence of marker gene function. Alternatively, a  
30 marker gene can be placed in tandem with a sequence encoding TPPT under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding TPPT and that express TPPT may be identified by a variety of procedures known to those of skill in the art. These  
35 procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or

chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of TPPT using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and  
5 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on TPPT is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and  
10 Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding TPPT  
15 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding TPPT, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety  
20 of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding TPPT may be cultured under  
25 conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode TPPT may be designed to contain signal sequences which direct secretion of TPPT through a prokaryotic or eukaryotic cell membrane.

30 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity.  
35 Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the

American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding TPPT may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric TPPT protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of TPPT activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the TPPT encoding sequence and the heterologous protein sequence, so that TPPT may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled TPPT may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

TPPT of the present invention or fragments thereof may be used to screen for compounds that specifically bind to TPPT. At least one and up to a plurality of test compounds may be screened for specific binding to TPPT. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of TPPT, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) *Current Protocols in Immunology* 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which TPPT binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express TPPT, either as a secreted

protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing TPPT or cell membrane fractions which contain TPPT are then contacted with a test compound and binding, stimulation, or inhibition of activity of either TPPT or the compound is analyzed.

- 5       An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with TPPT, either in solution or affixed to a solid support, and detecting the binding of TPPT to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a
- 10   labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

TPPT of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of TPPT. Such compounds may include agonists, antagonists, or partial or

15   inverse agonists. In one embodiment, an assay is performed under conditions permissive for TPPT activity, wherein TPPT is combined with at least one test compound, and the activity of TPPT in the presence of a test compound is compared with the activity of TPPT in the absence of the test compound. A change in the activity of TPPT in the presence of the test compound is indicative of a compound that modulates the activity of TPPT. Alternatively, a test compound is combined with an

20   in vitro or cell-free system comprising TPPT under conditions suitable for TPPT activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of TPPT may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding TPPT or their mammalian homologs may

25   be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of

30   interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids

35   Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred

to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding TPPT may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding TPPT can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding TPPT is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress TPPT, e.g., by secreting TPPT in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of TPPT and human transport proteins. In addition, the expression of TPPT is closely associated with neurological, cardiovascular, reproductive, gastrointestinal, and hematopoietic/immune tissues, and inflammation, cell proliferation, and cancer. Therefore, TPPT appears to play a role in transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated with increased TPPT expression or activity, it is desirable to decrease the expression or activity of TPPT. In the treatment of disorders associated with decreased TPPT expression or activity, it is desirable to increase the expression or activity of TPPT.

Therefore, in one embodiment, TPPT or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer; cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline

myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis; neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia; and other

5 disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn

10 syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism,

15 hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, and pseudovitamin D-deficiency rickets; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's

20 disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and

25 Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders,

30 peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial

35 frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins,

thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve,

5 mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary congestion and edema, pulmonary embolism, pulmonary

10 hemorrhage, pulmonary infarction, pulmonary hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease, emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis,

15 pulmonary eosinophilia bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung transplantation; a

20 reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm

25 physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis,

30 autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia,

35 irritable bowel syndrome, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis,

pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, 5 fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, 10 myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing TPPT or a fragment or derivative 15 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified TPPT in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not 20 limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of TPPT may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not limited to, those listed above.

In a further embodiment, an antagonist of TPPT may be administered to a subject to treat or 25 prevent a disorder associated with increased expression or activity of TPPT. Examples of such disorders include, but are not limited to, those transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer described above. In one aspect, an antibody which specifically binds TPPT may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to 30 cells or tissues which express TPPT.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding TPPT may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TPPT including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary 35 sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made



by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

5       An antagonist of TPPT may be produced using methods which are generally known in the art. In particular, purified TPPT may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind TPPT. Antibodies to TPPT may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and  
10 fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with TPPT or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to  
15 increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to TPPT  
20 have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of TPPT amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

25       Monoclonal antibodies to TPPT may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and  
30 Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda,  
35 S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce TPPT-specific single

chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte  
5 population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for TPPT may also be generated. For example, such fragments include, but are not limited to,  $F(ab)_2$  fragments produced by pepsin  
10 digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the  $F(ab)_2$  fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired  
15 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between TPPT and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering TPPT epitopes is generally used, but a competitive binding assay may  
20 also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for TPPT. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of TPPT-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions.  
25 The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple TPPT epitopes, represents the average affinity, or avidity, of the antibodies for TPPT. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular TPPT epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the  
30 TPPT-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of TPPT, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons,  
35 New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to

determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of TPPT-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines  
5 for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding TPPT, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA,  
10 RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding TPPT. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding TPPT. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense  
15 sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral  
20 vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.*  
25 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding TPPT may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined  
30 immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal,  
35 R.G. (1995) *Science* 270:404-410; Verma, I.M. and Somia, N. (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated

cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) Nature 335:395-396; Poeschla, E. et al. (1996) Proc. Natl. Acad. Sci. USA. 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in TPPT expression or regulation causes disease, the expression of TPPT from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in TPPT are treated by constructing mammalian expression vectors encoding TPPT and introducing these vectors by mechanical means into TPPT-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) Annu. Rev. Biochem. 62:191-217; Ivics, Z. (1997) Cell 91:501-510; Boulay, J-L. and H. Récipon (1998) Curr. Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of TPPT include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). TPPT may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) Proc. Natl. Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding TPPT from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) Virology 52:456-467), or by electroporation (Neumann, E. et al. (1982) EMBO J. 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to TPPT expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding TPPT under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding TPPT to cells which have one or more genetic abnormalities with respect to the expression of TPPT. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544; and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding TPPT to target cells which have one or more genetic abnormalities with respect to the expression of TPPT. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing TPPT to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has

been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant

5 HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned

10 herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to

15 deliver polynucleotides encoding TPPT to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotech.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA,

20 resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for TPPT into the alphavirus genome in place of the capsid-coding region results in the production of a large number of TPPT-coding RNAs and the synthesis of high levels of TPPT in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent

25 infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of TPPT into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating

30 infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it

35 causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have

been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

5           Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding TPPT.

10           Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of  
15 candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

20           Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding TPPT. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

25           RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine,  
30 and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

          An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding TPPT. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not  
35 limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular

chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased TPPT expression or activity, a compound which specifically inhibits expression of the polynucleotide  
5 encoding TPPT may be therapeutically useful, and in the treatment of disorders associated with decreased TPPT expression or activity, a compound which specifically promotes expression of the polynucleotide encoding TPPT may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method  
10 commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a  
15 polynucleotide encoding TPPT is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding TPPT are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the  
20 polynucleotide encoding TPPT. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide  
25 can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide  
30 nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells  
35 taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved



using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and  
5 monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest  
10 edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such pharmaceutical compositions may consist of TPPT, antibodies to TPPT, and mimetics, agonists, antagonists, or inhibitors of TPPT.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,  
15 intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Pharmaceutical compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol  
20 delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration  
25 enhancers.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of pharmaceutical compositions may be prepared for direct intracellular  
30 delivery of macromolecules comprising TPPT or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, TPPT or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system  
35 (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell

culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

- 5 A therapeutically effective dose refers to that amount of active ingredient, for example TPPT or fragments thereof, antibodies of TPPT, and agonists, antagonists or inhibitors of TPPT, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose
- 10 lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with
- 15 little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

- The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the
- 20 severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

- 25 Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells,
- 30 conditions, locations, etc.

### DIAGNOSTICS

- In another embodiment, antibodies which specifically bind TPPT may be used for the diagnosis of disorders characterized by expression of TPPT, or in assays to monitor patients being treated with TPPT or agonists, antagonists, or inhibitors of TPPT. Antibodies useful for diagnostic
- 35 purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for TPPT include methods which utilize the antibody and a label to detect TPPT in human body fluids

or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

5 A variety of protocols for measuring TPPT, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of TPPT expression. Normal or standard values for TPPT expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to TPPT under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of TPPT expressed in subject, 10 control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding TPPT may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and 15 quantify gene expression in biopsied tissues in which expression of TPPT may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of TPPT, and to monitor regulation of TPPT levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding TPPT or closely related molecules may be used to 20 identify nucleic acid sequences which encode TPPT. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding TPPT, allelic variants, or related sequences.

25 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the TPPT encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:44-86 or from genomic sequences including promoters, enhancers, and introns of the TPPT gene.

Means for producing specific hybridization probes for DNAs encoding TPPT include the 30 cloning of polynucleotide sequences encoding TPPT or TPPT derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels, 35 such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding TPPT may be used for the diagnosis of disorders

- associated with expression of TPPT. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis,
- 5 normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer; cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial
  - 10 myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis; neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia; and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease,
  - 15 cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis,
  - 20 congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases,
  - 25 mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, and pseudovitamin D-deficiency rickets; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple
  - 30 sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis,
  - 35 encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial

nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia,

5 catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon

10 angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus

15 erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary congestion and edema, pulmonary embolism, pulmonary hemorrhage, pulmonary infarction, pulmonary hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease,

20 emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary eosinophilia bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary

25 hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung transplantation; a reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a

30 disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and

35 gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome,

allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding TPPT may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered TPPT expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding TPPT may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding TPPT may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding TPPT in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of TPPT, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding TPPT, under conditions suitable for hybridization or amplification.

5 Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

10 Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

15 With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development  
20 or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding TPPT may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding TPPT, or a fragment of a polynucleotide complementary to the polynucleotide encoding  
25 TPPT, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding TPPT may be used to detect single nucleotide polymorphisms (SNPs). SNPs are  
30 substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding TPPT are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal  
35 tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are

detectable using gel electrophoresis in non-denaturing gels. In fSCCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual  
5 overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

10 Methods which may also be used to quantify the expression of TPPT include radiolabeling or biotinylation nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of  
15 interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large  
20 numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and  
25 monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

30 In another embodiment, antibodies specific for TPPT, or TPPT or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci.*  
35 USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-



2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding TPPT may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) *Proc. Natl. Acad. Sci. USA* 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding TPPT on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, TPPT, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug

screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between TPPT and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with TPPT, or fragments thereof, and washed. Bound TPPT is then detected by methods well known in the art. Purified TPPT can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding TPPT specifically compete with a test compound for binding TPPT. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with TPPT.

In additional embodiments, the nucleotide sequences which encode TPPT may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, in particular U.S. Ser. No. 60/139,923, U.S. Ser. No. 60/148,177, U.S. Ser. No. 60/149,357, and U.S. Ser. No. 60/162,287, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated

using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

- 5 In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic
- 10 oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,
- 15 PBLUEScript plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

- 20 Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96
- 25 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

- Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in
- 30 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

## III. Sequencing and Analysis

- Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.
- 35 Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ

Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic  
5 separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some  
10 of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools,  
15 programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO  
20 software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

25 The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire  
30 annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying  
35 against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such

as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID  
5 NO:44-86. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs  
10 from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer  
15 search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

20 The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair  
25 (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced  
30 either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding TPPT occurred. Analysis involved the categorization of cDNA libraries by  
35 organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous,

reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in

5 Table 3.

#### V. Chromosomal Mapping of TPPT Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:44-49 and SEQ ID NO:54-86 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these  
10 databases that matched SEQ ID NO:44-49 and SEQ ID NO:54-86 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a  
15 mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

The genetic map locations of SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:60, SEQ ID NO:65, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:76, SEQ ID NO:80, and SEQ ID NO:83 are described in The Invention as ranges, or intervals, of human  
20 chromosomes. More than one map location is reported for SEQ ID NO:65, SEQ ID NO:73, SEQ ID NO:80, and SEQ ID NO:83, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:65, SEQ ID NO:73, SEQ ID NO:80, and SEQ ID NO:83 were assembled into their respective clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of  
25 measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Diseases associated with the public and Incyte sequences located within the  
30 indicated intervals are also reported in the Invention section where applicable. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

#### VI. Extension of TPPT Encoding Polynucleotides

35 The full length nucleic acid sequences of SEQ ID NO:44-86 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this

fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target  
5 sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR  
10 was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C,  
15 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN  
20 quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by  
25 electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For  
30 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on  
35 antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was  
5 quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

10 In like manner, the polynucleotide sequences of SEQ ID NO:44-86 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:44-86 are employed to screen cDNAs,  
15 genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase  
20 (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

25 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and  
30 compared.

#### VIII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra),  
mechanical microspotting technologies, and derivatives thereof. The substrate in each of the  
35 aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers.



Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g.,

- 5 Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31.)

- Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The
- 10 array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of
- 15 complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### Tissue or Cell Sample Preparation

- Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and
- 20 poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with
- 25 GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85 °C to the stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc.
- 30 (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### Microarray Preparation

- 35 Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification

uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 µg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

- 5        Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C  
10    oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 µl of the array element DNA, at an average concentration of 100 ng/µl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

- 15        Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in 0.2% SDS and distilled water as before.

20    Hybridization

- Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65 °C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly  
25    larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60 °C. The arrays are washed for 10 min at 45 °C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45 °C in a second wash buffer (0.1X SSC), and dried.

Detection

- 30        Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-  
35    scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The  
5 emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on  
10 the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two  
15 fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a  
20 linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated  
25 to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### **IX. Complementary Polynucleotides**

Sequences complementary to the TPPT-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring TPPT. Although use of oligonucleotides  
30 comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of TPPT. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is  
35 designed to prevent ribosomal binding to the TPPT-encoding transcript.

#### **X. Expression of TPPT**

Expression and purification of TPPT is achieved using bacterial or virus-based expression systems. For expression of TPPT in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express TPPT upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of TPPT in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding TPPT by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, TPPT is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from TPPT at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified TPPT obtained by these methods can be used directly in the assays shown in Examples XI and XV.

#### 30 XI. Demonstration of TPPT Activity

TPPT transport activity is assayed by measuring uptake of labeled substrates into Xenopus laevis oocytes. Oocytes at stages V and VI are injected with TPPT mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 1mM Na<sub>2</sub>HPO<sub>4</sub>, 5 mM Hepes, 3.8 mM NaOH, 50µg/ml gentamycin, pH 7.8) to allow expression of TPPT. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids,

sugars, drugs, ions, and neurotransmitters) is initiated by adding labeled substrate (e.g. radiolabeled with  $^3\text{H}$ , fluorescently labeled with rhodamine, etc.) to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in  $\text{Na}^+$ -free medium, measuring the incorporated label, and comparing with controls. TPPT activity is proportional to the level of

5 internalized labeled substrate.

## XII. Functional Assays

TPPT function is assessed by expressing the sequences encoding TPPT at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice

10 include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10  $\mu\text{g}$  of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu\text{g}$  of an additional plasmid containing sequences encoding a

15 marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of

20 fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with

25 specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of TPPT on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding TPPT and either CD64 or CD64-GFP. CD64 and CD64-

30 GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding TPPT and other genes of interest can be analyzed by northern analysis

35 or microarray techniques.

### XIII. Production of TPPT Specific Antibodies

TPPT substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

5 Alternatively, the TPPT amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

10 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-TPPT activity by, for example, binding the peptide or TPPT to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### XIV. Purification of Naturally Occurring TPPT Using Specific Antibodies

Naturally occurring or recombinant TPPT is substantially purified by immunoaffinity chromatography using antibodies specific for TPPT. An immunoaffinity column is constructed by covalently coupling anti-TPPT antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing TPPT are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of TPPT (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/TPPT binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and TPPT is collected.

### XV. Identification of Molecules Which Interact with TPPT

TPPT, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled TPPT, washed, and any wells with labeled TPPT complex are assayed. Data obtained using different concentrations of TPPT are used to calculate values for the number, affinity, and association of TPPT with the candidate molecules.

35 Alternatively, molecules interacting with TPPT are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, *Nature* 340:245-246), or using commercially

available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

TPPT may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent  
5 No. 6,057,101).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be  
10 understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	44	264114	HNT2AGT01	028972R6 (SPLNFET01), 028972T6 (SPLNFET01), 264114H1 (HNT2AGT01), 452387R6 (TYLNNOT02), 736580R1 (TONSNOT01), 747955R6 (BRAITUT01), 936731R1 (CERVNOT01), 3206282H1 (PENCNOT03), 3344943H1 (SPLNNOT09), 3742964H1 (THYNNOT08), 4028320H1 (BRAINOT23), 4726757H1 (GBLADIT01), 5473562H1 (MCLRUNT01)
2	45	1455669	COLNFET02	1455669H1 (COLNFET02), 2877376F6 (THYRNOT10), 3536452F6 (KIDNNOT25)
3	46	2084989	PANCNOT04	1281527H1 (COLNNOT16), 1412985H1 (BRAINOT12), 2084989H1 (PANCNOT04), 2084989R6 (PANCNOT04), 2084989T6 (PANCNOT04), 2470481F6 (THP1NOT03), 2539015F7 (BONRTUT01), 3109754F6 (BRSTTUT15), 3694831H1 (PANCNOT19), 3700647H1 (SININOT05)
4	47	2501034	ADRETUT05	111466F1 (PITUNOT01), 111466R1 (PITUNOT01), 414042R6 (BRSTNOT01), 687891H1 (UTRSNOT02), 2501034H1 (ADRETUT05)
5	48	2745212	LUNGTUT11	000802H1 (U937NOT01), 008963H1 (HMC1NOT01), 009314H1 (HMC1NOT01), 135428F1 (BMARNOT02), 723168X19 (SYNOOAT01), 1000842R1 (BRSTNOT03), 1370189H1 (BSTMNON02), 1374329H1 (BSTMNON02), 2745212H1 (LUNGTUT11), 4920466H1 (TESTNOT11), SAIA02182F1
6	49	4833111	BRAVTXT03	864776T1 (BRAITUT03), 1911267F6 (CONNTUT01), 4833111H1 (BRAVTXT03), SARA02608F1, SARA02002F1
7	50	876677	LUNGAST01	876677H1 (LUNGAST01), 876677R6 (LUNGAST01), SCDA08642V1
8	51	2326143	OVARNOT02	867305R1 (BRAITUT03), 963058R2 (BRSTTUT03), 1715155F6 (UCMCNOT02), 1727927T6 (PROSNOT14), 2326143H1 (OVARNOT02), 2326143R6 (OVARNOT02), 3360563H1 (PROSTUT16)
9	52	2786302	BRSTNOT13	2786302H1 (BRSTNOT13), 2958321X303D1 (ADRENOT09), 2958321X305D1 (ADRENOT09), 2958321X308D1 (ADRENOT09)
10	53	3735780	SMCCNOS01	551126H1 (BEPINOT01), 2808373H1 (BLADTUT08), 3735780F6 (SMCCNOS01), 3735780H1 (SMCCNOS01), 3735780T6 (SMCCNOS01), 4760604T6 (BRANNOT01)
11	54	039026	HUVENOB01	039026H1 (HUVENOB01), 159164F1 (ADENINB01), 159164R1 (ADENINB01)
12	55	260607	HNT2RAT01	063159R6 (PLACNOB01), 260607R6 (HNT2RAT01), 1272850T1 (TESTTUT02), 1273069H1 (TESTTUT02), 2867453F6 (KIDNNOT20), 3082466H1 (BRAIUNT01), 4796739H1 (LIVRTUT09), 4799318F6 (MYEPUNT01), G1424405



Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
13	56	1429651	SINTBST01	1429651F1 (SINTBST01), 1429651H1 (SINTBST01), 1501096F6 (SINTBST01), 1989621T6 (CORPNOT02), SXLA01343V1, SXLA01183V1, SXLA01559V1, SXLA00812V1
14	57	2069971	ISLTNOT01	2069606F6 (ISLTNOT01), 2069971H1 (ISLTNOT01), 2374634F6 (ISLTNOT01), 2383754F6 (ISLTNOT01), 4171186T6 (SINTNOT21), SXLA01128V1, SXLA01348V1, SXLA01219V1, SXLA00260V1, SXLA00074V1
15	58	2329339	COLNNOT11	658662H1 (BRAINNOT03), 1544110R1 (PROSTUT04), 1657742F6 (URETTUT01), 1750523F6 (STOMTUT02), 2329339H1 (COLNNOT11), 2329339R6 (COLNNOT11), 3858671H1 (LNODNOT03), g1494061, g1891451
16	59	2540219	BONRTUT01	2540219H1 (BONRTUT01), 2540219T6 (BONRTUT01), 2554869F6 (THYMNOT03), g869197
17	60	2722462	LUNGUT10	883601R1 (PANCNOT05), 1525902F6 (UCMCL5T01), 1525902X18C1 (UCMCL5T01), 1525902X311D1 (UCMCL5T01), 1527325T6 (UCMCL5T01), 1554770X311D1 (BLADTUT04), 2417265H1 (HNT3AZT01), 2444786F6 (THPINOT03), 2722462H1 (LUNGUT10), 4293114H1 (BRABDIR01), 5070268T6 (PANCNOT23), SANA01850F1, SAJA01078R1, SANA02081F1, SAJA01813F1
18	61	2739264	OVARNOT09	000573H1 (U937NOT01), 494409F1 (HNT2NOT01), 494409R1 (HNT2NOT01), 2506506F6 (CONUTUT01), 2681059H1 (SINIUCT01), 2744648F6 (BRSTTUT14), 2805590F6 (BLADTUT08), 3770643H1 (BRSTNOT25), 4204278H1 (BRAITUT29), SAEA02093F1
19	62	2758310	THP1AZS08	487309R7 (HNT2AGT01), 1361439F1 (LUNGNOT12), 2758310H1 (THP1AZS08), SCFA05584V1, SCFA05940V1, SCFA05166V1, SCFA05135V1
20	63	2762348	BRSTNOT12	632097R6 (KIDNNOT05), 632097T6 (KIDNNOT05), 2762348H1 (BRSTNOT12), SCCA02837V1, SCCA05356V1, SCCA01377V1, SCCA05963V1, SCCA05364V1, SCCA02307V1, SCCA04327V1, SCCA02009V1
21	64	3715961	PENCNOT09	961523H1 (BRSTTUT03), 1863723F6 (PROSNOT19), 2265329H1 (UTRSNOT02), 2360619R6 (LUNGFEOT05), 2360619T6 (LUNGFEOT05), 2821718H1 (ADRETUT06), 3715961H1 (PENCNOT09), 5016160H1 (BRAXNOT03), 5499583H1 (BRABDIR01)
22	65	5108194	PROSTUS19	1322651X35 (BLADNOT04), 1322651X36 (BLADNOT04), 3494841H1 (ADRETUT07), 4958978F6 (TLYMNOT05), 5108194H1 (PROSTUS19), g1379009, g1527417
23	66	5503122	BRABDIR01	5503122F6 (BRABDIR01), 5503122H1 (BRABDIR01), 5503122R6 (BRABDIR01)

Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
24	67	5517972	LIVRDIR01	805957R1 (BSTMTOT01), 953622R1 (SCORNON01), 1501080F1 (SINTBST01), 1547381R6 (PROSTUT04), 2081843T6 (UTRSNOT08), 2811524F6 (OVARNOT10), 3212921H1 (BLADNOT08), 3250443H1 (SEMVN03), 3269479H1 (BRAINOT20), 3699555F6 (SININOT05), 3700568H1 (SININOT05), 4944050H1 (BRAIFEN05), 5517972H1 (LIVRDIR01)
25	68	5593114	COLCDIT03	2859465F6 (SININOT03), 2859465T6 (SININOT03), 3555656F6 (LUNGNOT31), 3555656T6 (LUNGNOT31), 4345952H1 (TLYMTXT01), 5593114H1 (COLCDIT03), 5874544H1 (COLTDIT04)
26	69	044775	TBLYNOT01	044775H1 (TBLYNOT01), 044775X3 (TBLYNOT01), 455640R1 (KERANOT01), 950702R1 (PANCNOT05), 2418550H1 (HNT3AZT01), 2798917H1 (NPOLNOT01), 2844696H1 (DRGLNOT01), 91718929
27	70	116588	KIDNNOT01	699714R6 (SYNORAT03), 831423R1 (PROSTUT04), 978875R1 (BRSTNOT02), 1350569F1 (LATRTUT02), 1447681R1 (PLACNOT02), 3177382F6 (UTRSTUT04), 3688796H1 (HEAANOT01), 3929008H1 (KIDNNOT19), 92106455, 92163092
28	71	875369	LUNGAST01	571573F1 (OVARNON01), 571573R1 (OVARNON01), 875369H1 (LUNGAST01), 875369R1 (LUNGAST01), 3569021H1 (HEAPNOT01)
29	72	1325518	LPARNOT02	1325518H1 (LPARNOT02), 1325518T6 (LPARNOT02), 1825553F6 (LSUBNOT03), SBAA02035F1
30	73	2060987	OVARNOT03	1378947T1 (LUNGNOT10), 1453290F1 (PENITUT01), 1459818R1 (COLNFET02), 1967477H1 (BRSTNOT04), 2060987H1 (OVARNOT03), 2455371F6 (ENDANOT01), 2499967F7 (ADRETUT05), 3093056T6 (BRSTNOT19), 3213366H1 (BLADNOT08), 4934158H1 (BRSTTUT20), SBYA01942U1
31	74	2172064	ENDCNOT03	2172064CT1 (ENDCNOT03), 2172064H1 (ENDCNOT03), SBLA01269F1
32	75	2219267	LUNGNOT18	2219267F6 (LUNGNOT18), 2219267H1 (LUNGNOT18), 3117478T6 (LUNGNOT13), 3126288T6 (LUNGNOT12), 3558495H1 (LUNGNOT31)
33	76	2308629	NGANNOT01	469862F1 (MMLR1DT01), 469862R1 (MMLR1DT01), 1594203X11C1 (BRAINOT14), 2191933H1 (THYRTUT03)
34	77	2660038	LUNGUT09	1326594F1 (LPARNOT02), 2256143H1 (OVARFUT01), 2278689R6 (PROSNON01), 2528425H1 (GBLANOT02), 2660038H1 (LUNGUT09), 2660038T6 (LUNGUT09), 3449964H1 (UTRSNON03), 5099879H1 (PROSTUS20), 91886680, 97833959

Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
35	78	2670745	ESOGTUT02	259200X12 (HNT2RAT01), 1266477F1 (BRAINT09), 2383364F6 (ISLTNOT01), 2670745H1 (ESOGTUT02), 3181526H1 (TLYJNOT01)
36	79	2676443	KIDNNOT19	607375R6 (BRSTTUT01), 1728626X15C1 (PROSNOT14), 1751773F6 (LIVRTUT01), 1751994T6 (LIVRTUT01), 1796032X14C1 (PROSTUT05), 2010172H1 (TESTNOT03), 2676443H1 (KIDNNOT19)
37	80	3295764	TLYJINT01	063264H1 (PLACNOB01), 434468T6 (THYRNOT01), 487721H1 (HNT2AGT01), 907796R2 (COLNNOT09), 1212556R7 (BRSTTUT01), 1251889H1 (LUNGFET03), 1653370F6 (PROSTUT08), 1653370X309D1 (PROSTUT08), 2192762F6 (THYRTUT03), 2226786F6 (SEMVTUT01), 3295764H1 (TLYJINT01), 3384471H1 (ESOGNOT04), SASA01137F1
38	81	3438320	PENCNOT06	3438320H1 (PENCNOT06), 3501438H1 (PROSTUT13), 3745542H1 (THYMNUT08), 3751060H1 (UTRSNOT18), 4979750F6 (HELATXT04), SADA00043F1, SADA00087F1
39	82	3986488	UTRSTUT05	1634141F6 (COLNNOT19), 1692115X12C1 (PROSTUT10), 1731310F6 (BRSTTUT08), 2046232H1 (THPIT7T01), 3557951H1 (LUNGNOT31), 4726788H1 (GBLADIT01)
40	83	4378816	LUNGNOT37	1318962H1 (BLADNOT04), 1520864F1 (BLADTUT04), 1684381F6 (PROSNOT15), 2055747R6 (BEPINOT01), 4378816H1 (LUNGNOT37)
41	84	4797137	LIVRTUT09	4797137F6 (LIVRTUT09), 4797137H1 (LIVRTUT09), 4797137T6 (LIVRTUT09)
42	85	5470806	MCLRUNT01	5470806H1 (MCLRUNT01), 5470806T6 (MCLRUNT01)
43	86	5473242	MCLRUNT01	5473242F6 (MCLRUNT01), 5473242T6 (MCLRUNT01)

Table 2

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
1	623	S521 S2 T3 S16 S99 S138 S144 T193 T264 T404 S448 S589 S151 T229 T337 T457 S562 S568	N97 N333	BTB domain: C44-F56 POZ domain: N10-Q211 Kelch repeat signature: E379-G392, T398-V412, L438-M452, T498-A512 Ring canal protein repeat: E122-P254	Ring canal protein [Drosophila melanogaster] g577276	MOTIFS BLIMPS-PFAM BLIMPS-PRINTS BLAST-GenBank BLAST-PRODROM BLAST-DOMO
2	99	T17	N15	Signal peptide: M1-G36 Transmembrane region: S25-W45 MRP(2) MRP(1) repeat: C30-V74	Multi-drug resistance- associated protein (MRP)-like protein- 1 (MLP-1) [Rattus norvegicus] g3242458	MOTIFS BLAST-GenBank BLAST-PRODROM SPScan HMMER
3	374	T334 T33 S137 T146 S291 S311 T346	N103 N127 N135 N138	Signal peptide: M1-N52	Tricarboxylate carrier [Rattus sp.] g545998	MOTIFS BLAST-GenBank SPScan
4	271	S234 T126 T169 Y141		Signal peptide: M1-C30 Transmembrane region: L233-F252	Weak similarity with honeybee ATP synthase A chain [Caenorhabditis elegans] g3878801	MOTIFS BLAST-GenBank SPScan HMMER
5	323	S99 S125 S192 T277 S307 S309 T110 Y212		Leucine zipper: L284-L305	Cu <sup>2+</sup> -transporting ATPase homolog [Arabidopsis thaliana] g2464854	BLAST-GenBank MOTIFS
6	274	S96 T198 S215 T29 S121 S164 S170		Mitochondrial energy transfer proteins: G5-L266 Signal peptide: M1-G17	Pet8p [Saccharomyces cerevisiae] g495307	BLAST-GenBank HMMER-PFAM MOTIFS ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO SPScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
7	291	S6 T113 T173 T147 S230 T258	N226 N261	Signal peptide: M1-T42 Transmembrane domain: W29-I54 Band 7 protein family: C50-V62, K90-E224 Membrane stomatin: E14-N283	Stomatin [Homo sapiens] g1161562	MOTIFS BLAST-GenBank SPScan HMMER BLIMPS-BLOCKS BLAST-DOMO BLAST-PRODOM
8	381	S2 S35 T57 S92 T104 S191 S302 S334 S335 S336 T43 T250 T255 T304 S311 S370 Y65	N218 N253 N259		K <sup>+</sup> channel modulatory factor DEBT-91 [Mus musculus] g4838557	MOTIFS BLAST-GenBank
9	190	T160 S17 T71 S77 T78 S111 S134 S142	N87	ABC transporter family: R79-K177 ATP/GTP-binding site motif A (P-loop): G102-S109	ABC2 transporter [Mus musculus] g495259	MOTIFS BLAST-GenBank BLAST-DOMO
10	297	S17 S114 T136 S16	N287	Mitochondrial carrier protein signature: E117-I297 Graves Disease carrier protein: P137-T157, L259-S279	Similar to human ADP/ATP carrier protein [C. elegans] g3879938	MOTIFS BLAST-GenBank HMMER-PFAM BLIMPS-PRINTS
11	89	T37 T47 T60 S64			Mitochondrial import protein Tim9p [Saccharomyces cerevisiae] g3747026	BLAST-GenBank MOTIFS
12	115	T108 T84		Signal peptide: M1-G24 Transmembrane domain: G35-F57 Sodium neurotransmitter symporter signature: R7-S61		MOTIFS SPScan HMMER ProfileScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
13	675	T54 T50 S99 T127 S413 T558 S645 T654 T47 S242 T602 T611 Y501	N243 N247 N301 N601	Transmembrane domains: I29-V48, L103-I121, L177-G196, I210-M229, L417-W435, F481-Y501, Y521-W541 Sodium symporter family domain: Y58-G487 Sodium:solute symporter signature: Y35-G89, M111-R140, L173-G227, P460-G469	Sodium-glucose cotransporter [Oryctolagus cuniculus] g473969	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO
14	320	T84 S304 T11 S75 S80 S164 Y20	N162 N234	Transmembrane domains: I92-L112, I201-K219 Zinc transporter signature: A28-V142, D199-E303 Cation transporter domain: S48-L74	Zinc transporter Znt-2 [Rattus norvegicus] g1256378	BLAST-GenBank MOTIFS HMMER BLIMPS-PRODOM BLAST-PRODOM BLAST-DOMO
15	462	S111 S145 S183 S233 T26 T185 S202 T243	N24 N279	Kelch repeat motifs: C299-N349; F350-R399 Y400-G446 BTB domain: F50-L117 POZ domain: Y27-E215	Ring canal protein [Drosophila melanogaster] g577276	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO
16	98	T22 Y37		Signal peptide: M1-S17 Mitochondrial carrier proteins domain: C4-I89 Mitochondrial carrier proteins signature sequence: V6-G19, G19-A33, G63-E83	Carrier protein (cl) [Caenorhabditis elegans] g472902	BLAST-GenBank MOTIFS SPScan HMMER-PFAM ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
17	748	S55 S196 T254 S307 S327 T491 T534 T550 T571 S635 S648 S677 T696 S283 S291 T314 S629 S701 Y556	N531 N543 N548 N627	Signal peptide: M1-A61 Transmembrane domains: L39-L56, I167-F186, C229-F252, G438-L455, M492-F509, L598-I618 Ion transport proteins signature: F85-V251, L369-I618	Voltage-gated calcium channel [Rattus norvegicus] g4586963	BLAST-GenBank MOTIFS SPScan HMMER HMMER-PFAM BLIMPS-PRINTS
18	507	T200 S183 T232 T284 T349 T150 T252 S253 S319 S383 Y454 S460 S104 T178 S320 S321 T498 T531 Y365	N220 N250 N364 N496	Signal peptide: M1-G26	Nucleoporin p54 [Rattus norvegicus] g1537070	BLAST-GenBank MOTIFS SPScan
19	592	S460 S104 T178 S320 S321 T498 T531 Y365		ABC precursor signature: N153-Q162, F210-A229, G234-I254, V312-G332, T366-V378	ABC transporter [Methanobacterium thermo.] g2622773	BLAST-GenBank MOTIFS BLIMPS-PRODOM BLAST-PRODOM
20	841	T98 S120 S203 T214 T276 S388 T438 T700 T838 T167 T179 S280 T370 S435 S531 S539 S666 S693 S830	N368 N490 N624	Transmembrane domains: Y451-D469, M544-F562, F577-F597, G775-M797 Vacuolar ion transport subunit signature: M10-F831	Vacuolar H <sup>+</sup> /ATPase subunit [Rattus norvegicus] g206430	BLAST-GenBank MOTIFS HMMER BLIMPS-PRODOM BLAST-PRODOM BLAST-DOMO
21	253	S50 T139 T152 T177 S202 T143 Y55		Mitochondrial carrier proteins domain: Y31-S248 Mitochondrial energy transfer proteins signature sequence: I62-Q86, I110-G122	Mitochondrial uncoupling protein UCP-4 [Homo sapiens] g4324701	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
22	229	S69 S26 S109 T162 S178 S25 S64 S65 T210 S219		Signal peptide: M1-A47 Mitochondrial carrier proteins domain: Q32-G220 Mitochondrial carrier proteins signature sequence: S36-T49, T49-V63, G92-E112, T144-T162, Y187-F205	Grave's disease carrier protein [Bos taurus] g387	BLAST-GenBank MOTIFS SPScan HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
23	170	S26 S31 S149 S164 T22 T157	N66 N145	Dihydroxyipyridine-sensitive L-type calcium channel signature: Y2-A47, I49-V77, A83-N100, R106-E131 SH3 domain: V59-R122	Voltage-dependent calcium channel beta-4 subunit [Homo sapiens] g2058727	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO
24	655	T194 S195 S232 T362 S655 S4 S88 T135 T153 S187 T214 S322 T345 S353 S443 T609 S261 S381 S384	N338 N418 N557 N596	Transmembrane domains: I396-K417, Y494-S522, T538-V556 ABC transporters domain: F73-G262 ABC transporter family signature sequence: I78-L89, V186-D217	Breast cancer resistance protein (multidrug transporter) [Homo sapiens] g4038352	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO
25	184	T51 S29 T100 S138 S151 Y78	N27		Cation transport protein [E. coli] g495778	BLAST-GenBank MOTIFS
26	154	S54 S42 S62 T78 Y104		Mitochondrial energy transfer proteins signatures: P89-L97, M1-E41, M73-L152 Mitochondrial carrier protein domain: G2-L152	Similar to carrier protein C2 [C. elegans] g3879669	MOTIFS HMMER-PFAM BLAST-PRODOM BLAST-DOMO BLAST-GenBank



Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
27	438	S170 T5 T51 T265 T300 S425	N50 N423	Transmembrane domains: C91-L111, L237-I257, I305-M332, M332-L354, L391-V408, I186-A204	Multidrug efflux transporter [Bacillus subtilis] g2635104	MOTIFS HMMER BLAST-GenBank
28	237	S10 S47 T72 S28 S96 S148 T173 T222 S6 S21 T32 T61 T192	N35	Nucleic acid-binding protein E5.1 domain: S6-K128	ARL-6 interacting protein-4 [Mus musculus] g4927204	MOTIFS BLAST-DBOM BLAST-GenBank
29	219	T66 S194 T200		Signal peptide: M1-R19 or M1-K15 Caseins alpha/beta signature: M1-N39	Surface antigen [Trypanosoma cruzi] g161956	MOTIFS HMMER SPScan ProfileScan BLAST-GenBank
30	707	S31 T6 T55 T263 T328 T546 T580 T594 S662 S673 T32 S50 S231 T244 T306 T385 S439 S476 S533 S553 S624	N343 N570 N638 N703	Potassium channel signature: A62-T81 Potassium channel integral membrane protein domain: S13-D117	NY-REN-45 antigen (similar to potassium channel protein) [Homo sapiens] g5360115	MOTIFS BLIMPS-PRINTS BLAST-DBOM BLAST-GenBank
31	279	T18 T245 T206	N181	Signal cleavage: M1-G45 Connexin domains: M1-V99, V20-Y44 Connexin signatures: L33-V86, L152-F205, F51-P73, S76-L96, L133-Y159, C169-T189, I190-L218 Gap junction protein connexin transmembrane regions: F5-Y97, L133- K223, M1-S130	Gap junction protein (similar to connexin) [Homo sapiens] g3006230	MOTIFS SPScan HMMER BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PFAM ProfileScan BLAST-PRODOR BLAST-DBOM BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
32	154	S114		Signal peptide: M1-A35 or M1-A14 Transmembrane domain: F83-L102	mBOCT (potent organic cation transporter) [Mus musculus] g4589468	MOTIFS HMME SPScan BLAST-GenBank
33	289	T83 T205 S269 T279	N60	Mitochondrial energy transfer proteins signatures: M1-G147, P17, P115, N185-K280, A101-Q181, Y184-I278 Mitochondrial carrier protein domains: M1-E176, N185-K280 Mitochondrial transmembrane transport protein regions: P17-R182, P180-I278	Mitochondrial solute carrier [Onchocerca volvulus] g1518458	MOTIFS HMME-PFAM BLAST-PRODOM ProfileScan BLAST-GenBank
34	300	S189 S195 S204 T257		Mitochondrial energy transfer proteins signatures: P19-W27, D2-I53, L209-L295 Mitochondrial carrier protein domain: D2-Y295 Transport protein domain: P122-Y295	YKL522=mitochondria 1 ADP/ATP carrier protein homolog [Saccharomyces cerevisiae] g254449	MOTIFS HMME-PFAM ProfileScan BLAST-PRODOM BLAST-DOMO BLAST-GenBank
35	382	S34 S207 T221 S312 T40 S53 T112 T117 T277 S337	N96 N372	Kelch motifs: H191-G249, E250-D301	Similarity to Human host cell factor C1 [Homo sapiens] g3875291	MOTIFS HMME BLAST-PFAM BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
36	287	T36 T118 S180 S230 T84 S168 T244		Mitochondrial energy transfer proteins signatures: P26-L34, P219-L227, L97-G193, W10-V89, D197-F281, P96-Y194 Mitochondrial carrier protein domain: A5-F281 Mitochondrial brown fat region: Y82-Q94, V151-S168, Y194-C212	Mitochondrial dicarboxylate carrier [Rattus norvegicus] g3646426	MOTIFS HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO BLAST-GenBank
37	497	T65 T135 S147 T360 S8 T22 S45 S291	N63 N314 N414	Transmembrane domains: M114-T137, M364-M380, Y390-A413, A421-D444, F456-V478 Folate transporter domains: W30-H218, I253-K484	Reduced folate carrier [Homo sapiens] g1041934	MOTIFS HMMER BLAST-PRODOM BLAST-DOMO SPScan BLAST-GenBank
38	228	T21 S124 T145 S158 T190 T95 S132 S137 T177		Heme-binding domain in cytochrome b5: Y19-G98 Cytochrome b5 family domain: H28-P75	cytochrome b5 containing fusion protein [Helianthus annuus] g1040729 P=1.2e-07	MOTIFS HMMER-PFAM BLAST-GenBank ProfileScan
39	273	T63 S158 T48	N214	Transmembrane domains: L85-N105, F180-Y200 Intermembrane space domain: L30-L251	Sqv-7-like protein (similar to nucleotide-sugar transporters) [Homo sapiens] g4008517	MOTIFS HMMER BLAST-DOMO BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
40	206	S187 S201	N158	Signal peptide: M1-G29 or M1-A27 Emopamil binding protein: G37-S187, L15-K203 Transmembrane domain: Y164-L183	C-8,7 sterol isomerase, aSI1 [Arabidopsis thaliana] g2772934	MOTIFS HMMER ProfileScan BLAST-DOMO BLAST-GenBank
41	235	S192 S200 S56 T95 T146 S199 T207 S229 T53 T61 T69 T119 T148 Y70	N123	Transmembrane domain: F15-I34, M155-V174 Channel myelin protein: L18-M181 Sodium channel beta-2 subunit precursor: F15-E210 Immunoglobulin domain: I34-V136	Myelin protein zero (MPZ) [Homo sapiens] g2160399	MOTIFS HMMER BLIMPS-PRINTS BLAST-PRODUM BLAST-DOMO BLAST-GenBank
42	147	T79 T116 S3 S66 Y89 Y98	N118	Signal peptide: M1-G23 or M1-A20 Transthyretin signature: S28-S132 Transthyretin domain: G21-Q146	Transthyretin precursor [Sus scrofa] g1009702	MOTIFS HMMER ProfileScan BLAST-PRODUM BLAST-DOMO BLAST-GenBank BLIMPS-BLOCKS BLIMPS-PRINTS
43	147	T5 S88 T39		Globin domain: V2-H147 Heme oxygen transport protein domain: L32-H147	III beta-3 globin [Rattus norvegicus] g395943	MOTIFS HMMER-PFAM BLAST-PRODUM BLAST-DOMO BLIMPS-BLOCKS BLIMPS-PRINTS

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
44	1567-1611 2107-2151	Gastrointestinal (0.203) Hematopoietic/Immune (0.188) Nervous (0.156)	Cell Proliferation and Cancer (0.547) Inflammation (0.422)	PBLUESCRIPT
45	1-92 351-434	Endocrine (0.333) Developmental (0.167) Gastrointestinal (0.167) Musculoskeletal (0.167) Reproductive (0.167)	Cell Proliferation and Cancer (0.833) Inflammation (0.167)	pINCY
46	920-964 1352-1396	Reproductive (0.304) Gastrointestinal (0.174) Cardiovascular (0.130) Hematopoietic/Immune (0.130) Nervous (0.130)	Cell Proliferation and Cancer (0.478) Inflammation (0.391)	PSPORT1
47	1-80 768-848	Nervous (0.273) Reproductive (0.273) Gastrointestinal (0.127) Hematopoietic/Immune (0.127)	Cell Proliferation and Cancer (0.564) Inflammation (0.400)	pINCY
48	111-194 687-758	Reproductive (0.221) Nervous (0.185) Gastrointestinal (0.124)	Cell Proliferation and Cancer (0.552) Inflammation (0.343)	pINCY
49	1-97	Nervous (0.234) Hematopoietic/Immune (0.191) Gastrointestinal (0.149)	Cell Proliferation and Cancer (0.617) Inflammation (0.340)	pINCY
50	218-262	Cardiovascular (1.000)	Cancer (0.333) Inflammation/Trauma (0.333) Cell Proliferation (0.333)	PSPORT1
51	811-855	Hematopoietic/Immune (0.180) Gastrointestinal (0.146) Reproductive (0.281)	Cancer (0.393) Inflammation/Trauma (0.515) Cell Proliferation (0.146)	PSPORT1
52	595-639	Gastrointestinal (0.286) Reproductive (0.714)	Cancer (0.429) Inflammation/Trauma (0.429)	pINCY
53	96-140	Cardiovascular (0.167) Hematopoietic/Immune (0.167) Nervous (0.250) Reproductive (0.167)	Cancer (0.250) Inflammation/Trauma (0.167) Cell Proliferation (0.167)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
54	507-551	Reproductive (0.323) Gastrointestinal (0.154) Nervous (0.123)	Cancer (0.446) Inflammation/Trauma (0.308) Cell Proliferation (0.185)	PBLUESCRIPT
55	455-499	Urologic (0.333) Nervous (0.222) Reproductive (0.222)	Cancer (0.667) Cell Proliferation (0.333)	PBLUESCRIPT
56	1835-1879	Nervous (0.625) Gastrointestinal (0.375)	Inflammation/Trauma (0.375) Cancer (0.250) Neurological (0.250)	pINCY
57	811-855	Gastrointestinal (1.000)	Inflammation/Trauma (0.667)	pINCY
58	390-434	Reproductive (0.320) Nervous (0.240) Urologic (0.120)	Cancer (0.520) Inflammation/Trauma (0.240) Cell Proliferation (0.160)	PSPORT1
59	413-457	Gastrointestinal (0.333) Musculoskeletal (0.333) Nervous (0.333)	Cancer (0.333) Neurological (0.333)	pINCY
60	2021-2084	Nervous (0.197) Gastrointestinal (0.184) Reproductive (0.184)	Cancer (0.461) Inflammation/Trauma (0.316) Cell Proliferation (0.118)	pINCY
61	65-109	Nervous (0.226) Reproductive (0.208) Cardiovascular (0.113) Gastrointestinal (0.113)	Cancer (0.528) Inflammation/Trauma (0.301) Cell Proliferation (0.208)	pINCY
62	379-423 1867-1911	Reproductive (0.282) Gastrointestinal (0.205) Nervous (0.154)	Cancer (0.538) Inflammation/Trauma (0.282) Cell Proliferation (0.103)	PSPORT1
63	362-406 1193-1237	Urologic (0.500) Reproductive (0.333) Cardiovascular (0.167)	Cancer (0.667) Inflammation/Trauma (0.333)	pINCY
64	394-438	Nervous (0.294) Reproductive (0.265) Cardiovascular (0.118)	Cancer (0.382) Inflammation/Trauma (0.235) Cell Proliferation (0.118)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
65	768-812	Reproductive (0.300) Endocrine (0.200) Gastrointestinal (0.200) Hematopoietic/Immune (0.200)	Inflammation/Trauma (0.500) Cancer (0.400)	pINCY
66	77-121	Nervous (1.000)	Neurological (1.000)	pINCY
67	1999-2043	Reproductive (0.324) Nervous (0.265) Gastrointestinal (0.235)	Cancer (0.500) Inflammation/Trauma (0.294) Cell Proliferation (0.118)	pINCY
68	561-605	Hematopoietic/Immune (0.455) Gastrointestinal (0.182) Nervous (0.182)	Inflammation/Trauma (0.546) Cell Proliferation (0.182)	pINCY
69	679-729	Nervous (0.292) Gastrointestinal (0.208) Hematopoietic/Immune (0.125)	Cancer (0.250) Cell Proliferation (0.375) Inflammation/Trauma (0.416)	PBLUESCRIPT
70	95-366 1078-1185	Reproductive (0.206) Hematopoietic/Immune (0.186) Cardiovascular (0.127)	Cancer (0.373) Inflammation/Trauma (0.382) Cell Proliferation (0.176)	PBLUESCRIPT
71	33-152	Reproductive (0.275) Nervous (0.163) Gastrointestinal (0.137)	Cancer (0.438) Inflammation/Trauma (0.314) Cell Proliferation (0.176)	PSPORT1
72	81-779	Gastrointestinal (1.000)	Cancer (1.000)	pINCY
73	719-817 1202-1414	Reproductive (0.311) Hematopoietic/Immune (0.203) Gastrointestinal (0.122)	Cancer (0.459) Inflammation/Trauma (0.379) Cell Proliferation (0.203)	PSPORT1
74	1-848	Nervous (0.750) Dermatologic (0.250)	Cancer (0.250) Cell Proliferation (0.250) Inflammation/Trauma (0.500)	pINCY
75	1-478	Cardiovascular (0.714) Developmental (0.143) Hematopoietic/Immune (0.143)	Cancer (0.571) Cell Proliferation (0.286) Inflammation (0.143)	pINCY
76	1-134	Reproductive (0.253) Nervous (0.241) Gastrointestinal (0.127) Hematopoietic (0.127)	Cancer (0.494) Inflammation (0.215) Cell Proliferation (0.127)	PSPORT1

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
77	510-719 960-1100	Reproductive (0.467) Cardiovascular (0.133) Gastrointestinal (0.133)	Cancer (0.467) Inflammation/Trauma (0.467)	pINCY
78	180-293	Reproductive (0.230) Nervous (0.225) Gastrointestinal (0.124)	Cancer (0.478) Inflammation/Trauma (0.292) Cell Proliferation (0.191)	pINCY
79	192-653 795-935	Reproductive (0.417) Gastrointestinal (0.292) Urologic (0.125)	Cancer (0.750) Cell Proliferation (0.125) Inflammation/Trauma (0.167)	pINCY
80	139-1044	Reproductive (0.245) Nervous (0.143) Developmental (0.122)	Cancer (0.490) Inflammation/Trauma (0.286) Cell Proliferation (0.224)	pINCY
81	233-916	Reproductive (0.667) Cardiovascular (0.167) Nervous (0.167)	Cancer (0.500) Cell Proliferation (0.333) Inflammation (0.167)	pINCY
82	1-153 760-816	Gastrointestinal (0.282) Hematopoietic/Immune (0.205) Reproductive (0.205)	Inflammation/Trauma (0.461) Cancer (0.308) Cell Proliferation (0.205)	pINCY
83	57-299	Nervous (0.179) Reproductive (0.179) Gastrointestinal (0.128)	Cancer (0.564) Cell Proliferation (0.256) Inflammation/Trauma (0.180)	pINCY
84	1-707	Gastrointestinal (0.500) Hematopoietic/Immune (0.500)	Cancer (0.500) Inflammation (0.500)	pINCY
85	451-594	Hematopoietic/Immune (1.000)	Cell Proliferation (1.000)	pINCY
86	8-124 161-187 407-472	Developmental (1.000)	Cell Proliferation (1.000)	pINCY



Table 4

SEQ ID NO:	Library	Library Comment
44	HNT2AGT01	Library was constructed at Stratagene (STR937233), using RNA isolated from the hNT2 cell line derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor. Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
45	COLNFET02	Library was constructed using RNA isolated from the colon tissue of a Caucasian female fetus, who died at 20 weeks' gestation.
46	PANCNOT04	Library was constructed using RNA isolated from the pancreatic tissue of a 5-year-old Caucasian male, who died in a motor vehicle accident. Serology was positive for cytomegalovirus (CMV).
47	ADRETUT05	Library was constructed RNA isolated from adrenal tumor tissue removed from a 52-year-old Caucasian female during a unilateral adrenalectomy. Pathology indicated a pheochromocytoma.
48	LUNGTUT11	Library was constructed using RNA isolated from lung tumor tissue removed from the right lower lobe of a 57-year-old Caucasian male during a segmental lung resection. Pathology indicated an infiltrating grade 4 squamous cell carcinoma. Multiple intrapulmonary peribronchial lymph nodes showed metastatic squamous cell carcinoma. Patient history included a benign brain neoplasm and tobacco abuse. Family history included spinal cord cancer, type II diabetes, cerebrovascular disease, and malignant prostate neoplasm.
49	BRAVXT03	Library was constructed using RNA isolated from treated astrocytes removed from the brain of a female fetus who died after 22 weeks' gestation. The cells were treated with tumor necrosis factor (TNF) alpha and interleukin 1 (IL-1), 10ng/ml each for 24 hours.
50	LUNGAST01	Library was constructed using RNA isolated from the lung tissue of a 17-year-old Caucasian male, who died from head trauma. Patient history included asthma.
51	OVARNOT02	Library was constructed using RNA isolated from ovarian tissue removed from a 59-year-old Caucasian female who died of a myocardial infarction. Patient history included cardiomyopathy, coronary artery disease, previous myocardial infarctions, hypercholesterolemia, hypotension, and arthritis.
52	BRSTNOT13	Library was constructed using RNA isolated from breast tissue removed from the left medial lateral breast of a 36-year-old Caucasian female during bilateral simple mastectomy and total breast reconstruction. Pathology indicated benign breast tissue. Patient history included a breast neoplasm, depressive disorder, hyperlipidemia, chronic stomach ulcer, and an ectopic pregnancy. Family history included myocardial infarction, cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, skin cancer, breast cancer, depressive disorder, esophageal cancer, bone cancer, Hodgkin's lymphoma, bladder cancer, and a heart condition.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
53	SMCCNOS01	Library was constructed using 7.56 x 10 <sup>6</sup> clones from a coronary artery smooth muscle cell library and was subjected to two rounds of subtraction hybridization for 48 hours with 6.12 x 10 <sup>6</sup> clones from a control coronary artery smooth muscle cell library. The starting library for subtraction was constructed using RNA isolated from coronary artery smooth muscle cells removed from a 3-year-old Caucasian male. The cells were treated with TNF alpha & IL-1 beta 10ng/ml each for 20 hours. The hybridization probe for subtraction was derived from a similarly constructed library from RNA isolated from untreated coronary artery smooth muscle cells from the same donor.
54	HUVENOB01	Library was constructed using RNA isolated from HUV-EC-C (ATCC CRL 1730) cells.
55	HNT2RAT01	Library was constructed at Stratagene (STR937231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
56	SINTBST01	Library was constructed using RNA isolated from ileum tissue obtained from an 18-year-old Caucasian female during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
57	ISLTNOT01	Library was constructed using RNA isolated from a pooled collection of pancreatic islet cells.
58	COLNNOT11	Library was constructed using RNA isolated from colon tissue removed from a 60-year-old Caucasian male during a left hemicolectomy.
59	BONRTUT01	Library was constructed using RNA isolated from rib tumor tissue removed from a 16-year-old Caucasian male during a rib osteotomy and a wedge resection of the lung. Pathology indicated metastatic grade 3 (of 4) osteosarcoma, forming a mass involving the chest wall.
60	LUNGTVT10	Library was constructed using RNA isolated from lung tumor tissue removed from the left upper lobe of a 65-year-old Caucasian female during a segmental lung resection. Pathology indicated metastatic grade 2 myxoid liposarcoma and metastatic grade 4 liposarcoma. Patient history included soft tissue cancer, breast cancer, and secondary lung cancer.
61	OVARNOT09	Library was constructed using RNA isolated from ovarian tissue removed from a 28-year-old Caucasian female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. Pathology indicated multiple follicular cysts ranging in size from 0.4 to 1.5 cm in the right and left ovaries, chronic cervicitis and squamous metaplasia of the cervix, and endometrium in weakly proliferative phase. Family history included benign hypertension, hyperlipidemia, and atherosclerotic coronary artery disease.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
62	THP1AZS08	Library was constructed using 5.76 million clones from a 5-aza-2'-deoxycytidine (AZ) treated THP-1 promonocyte cell line library. Starting RNA was made from THP-1 promonocyte cells treated for three days with 0.8 micromolar AZ. 5.76 million clones from the AZ-treated THP-1 cell library were then subjected to two rounds of subtractive hybridization with 5 million clones from the untreated THP-1 cell library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (1991) Nucleic Acids Res. 19:1954, and Bonaldo et al. (1996) Genome Research 6:791. THP-1 (ATCC TIB 202) is a human promonocyte cell line derived from peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
63	BRSTNOT12	Library was constructed using RNA isolated from diseased breast tissue removed from a 32-year-old Caucasian female during a bilateral reduction mammoplasty. Pathology indicated nonproliferative fibrocystic disease. Family history included benign hypertension and atherosclerotic coronary artery disease.
64	PENCNOT09	Library was constructed using RNA isolated from penis right corpora cavernosa tissue.
65	PROSTUS19	Library was constructed using 2.36 million clones from a prostate tumor library which was subjected to two rounds of subtraction hybridization with 2.36 million clones from a normal prostate library. The starting library for subtraction was constructed using RNA isolated from prostate tumor tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenocarcinoma (Gleason grade 3+3) involving the prostate peripherally with invasion of the capsule. Adenofibromatous hyperplasia was present. The patient presented with elevated prostate-specific antigen (PSA). Patient history included diverticulitis of the colon, asbestosis, and thrombophlebitis. Family history included benign hypertension, multiple myeloma, hyperlipidemia, and rheumatoid arthritis. The hybridization probe for subtraction was derived from a similarly constructed library, except that NotI-anchored oligo(dT) primer was used. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (1991) Nucleic Acids Res. 19:1954 and Bonaldo, et al. (1996) Genome Research 6:791.
66	BRABDIR01	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. Patient history included Huntington's disease, emphysema, and tobacco abuse.
67	LIVRDIR01	Library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis. Serology was positive for anti-mitochondrial antibody.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
68	COLCDIT03	Library was constructed using RNA isolated from diseased colon polyp tissue removed from the cecum of a 67-year-old female. Pathology indicated a benign cecum polyp. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma that arose in tubulovillous adenoma forming a fungating mass in the cecum.
69	TBLYN0T01	Library was constructed at Stratagene (STR937214) using RNA isolated from a hybrid of T-B lymphoblasts from an untreated leukemic cell line.
70	KIDN0T01	Library was constructed using RNA isolated from the kidney tissue of a 64-year-old Caucasian female, who died from an intracranial bleed. Patient history included rheumatoid arthritis and tobacco use.
71	LUNGAST01	Library was constructed using RNA isolated from the lung tissue of a 17-year-old Caucasian male, who died from head trauma. Patient history included asthma.
72	LPARN0T02	Library was constructed using RNA isolated from tissue obtained from the left parotid (salivary) gland of a 70-year-old male with parotid cancer.
73	OVAR0T03	Library was constructed using RNA isolated from ovarian tissue removed from a 43-year-old Caucasian female during a bilateral salpingo-oophorectomy. Pathology for the associated tumor tissue indicated grade 2 mucinous cyst adenocarcinoma. The patient presented with stress incontinence. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, cerebrovascular disease, breast cancer, and uterine cancer.
74	ENDC0T03	Library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
75	LUNG0T18	Library was constructed using RNA isolated from left upper lobe lung tissue removed from a 66-year-old Caucasian female. Pathology for the associated tumor tissue indicated a grade 2 adenocarcinoma. Patient history included cerebrovascular disease, atherosclerotic coronary artery disease, and pulmonary insufficiency. Family history included a myocardial infarction and atherosclerotic coronary artery disease.
76	NGANN0T01	Library was constructed using RNA isolated from tumorous neuroganglion tissue removed from a 9-year-old Caucasian male during a soft tissue excision of the chest wall. Pathology indicated a ganglioneuroma. Family history included asthma.
77	LUNG0T09	Library was constructed using RNA isolated from lung tumor tissue removed from a 68-year-old Caucasian male during segmental lung resection. Pathology indicated invasive grade 3 squamous cell carcinoma and a metastatic tumor. Patient history included type II diabetes, thyroid disorder, depressive disorder, hyperlipidemia, esophageal ulcer, and tobacco use.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
78	ESOGTUT02	Library was constructed using RNA isolated from esophageal tumor tissue obtained from a 61-year-old Caucasian male during a partial esophagectomy, proximal gastrectomy, pyloromyotomy, and regional lymph node excision. Pathology indicated an invasive grade 3 adenocarcinoma in the esophagus. Family history included atherosclerotic coronary artery disease, type II diabetes, chronic liver disease, primary cardiomyopathy, benign hypertension, and cerebrovascular disease.
79	KIDNNOT19	Library was constructed using RNA isolated from kidney tissue removed a 65-year-old Caucasian male during an exploratory laparotomy and nephroureterectomy. Pathology for the associated tumor tissue indicated a grade 1 renal cell carcinoma within the upper pole of the left kidney. Patient history included malignant melanoma of the abdominal skin, benign neoplasm of colon, cerebrovascular disease, and umbilical hernia. Family history included myocardial infarction, atherosclerotic coronary artery disease, cerebrovascular disease, and prostate cancer.
80	TLYJINT01	Library was constructed using RNA isolated from a Jurkat cell line derived from the T cells of a male. Patient history included acute T-cell leukemia. This is an uninduced Jurkat cell line library from the same donor.
81	PENCNOT06	Library was constructed using RNA isolated from penis corpora cavernosa tissue removed from a 3-year-old Black male. Pathology for the associated tumor tissue indicated invasive grade 4 urothelial carcinoma forming a soft tissue scrotal mass that invaded the cavernous body of the penis and encased both testicles. Right inguinal lymph node showed metastatic grade 4 urothelial carcinoma, with extranodal invasion.
82	UTRSTUT05	Library was constructed using RNA isolated from uterine tumor tissue removed from a 41-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology indicated uterine leiomyoma. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Patient history included a ventral hernia and a benign ovarian neoplasm.
83	LUNGNOT37	Library was constructed using polyA RNA isolated from lung tissue removed from a 15-year-old Caucasian female who died from a closed head injury. Serology was positive for cytomegalovirus.
84	LIVRTUT09	Library was constructed using RNA isolated from an untreated C3A hepatocyte cell line which is a derivative of Hep G2, a cell line derived from a hepatoblastoma removed from a 15-year-old Caucasian male.
85	MCLRUNT01	Library was constructed using RNA isolated from untreated peripheral blood mononuclear cell tissue obtained from buffy coat, removed from a 60-year-old male.
86	MCLRUNT01	Library was constructed using RNA isolated from untreated peripheral blood mononuclear cell tissue obtained from buffy coat, removed from a 60-year-old male.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score $\geq$ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28,
  - 10 SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43,
- b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID
  - 15 NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35,
    - 20 SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43,
- c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID
  - 25 NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, and
- 30 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID
  - 35 NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID



NO:41, SEQ ID NO:42, and SEQ ID NO:43.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

15

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86.

25

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

30

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

35

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

5 11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
- 10 NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
- 15 b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
- 20 NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
- c) a polynucleotide sequence complementary to a),
- 25 d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

30

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe
- 35 specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

5

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

10

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

15

17. A pharmaceutical composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43.

20

18. A method for treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 16.

25

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

30

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A pharmaceutical composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

35

21. A method for treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 20.

- 5           22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
  - b) detecting antagonist activity in the sample.

- 10           23. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional TPPT, comprising administering to a patient in need of such treatment a pharmaceutical composition  
15 of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:
- a) combining the polypeptide of claim 1 with at least one test compound under suitable  
20 conditions, and
  - b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of  
25 claim 1, said method comprising:
- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
  - b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound,  
and
  - 30 c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

- 35           27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method

comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

5           28. An isolated polynucleotide comprising a polynucleotide sequence of SEQ ID NO:83.

29. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 28.

10           30. A cell transformed with a recombinant polynucleotide of claim 29.

31. A transgenic organism comprising a recombinant polynucleotide of claim 29.

32. A method for producing a polypeptide comprising an amino acid sequence of SEQ ID  
15 NO:40, the method comprising:

- a) culturing the cell of claim 30 under conditions suitable for expression of the polypeptide,  
and
- b) recovering the polypeptide so expressed.

20           33. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 28, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

## SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.  
 LAL, Preeti  
 YANG, Junming  
 YUE, Henry  
 HILLMAN, Jennifer L.  
 TANG, Y. Tom  
 BANDMAN, Olga  
 BURFORD, Neil  
 BAUGHN, Mariah R.  
 AZIMZAI, Yalda  
 LU, Dyung Aina M.  
 AU-YOUNG, Janice  
 PATTERSON, Chandra

<120> HUMAN TRANSPORT PROTEINS

<130> PF-0709 PCT

<140> To Be Assigned  
 <141> Herewith

<150> 60/139,923; 60/148,177; 60/149,357; 60/162,287  
 <151> 1999-06-17; 1999-08-10; 1999-08-18; 1999-10-28

<160> 86  
 <170> PERL Program

<210> 1  
 <211> 623  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 264114CD1

<400> 1  
 Met Ser Thr Gln Asp Glu Arg Gln Ile Asn Thr Glu Tyr Ala Val  
 1 5 10 15  
 Ser Leu Leu Glu Gln Leu Lys Leu Phe Tyr Glu Gln Gln Leu Phe  
 20 25 30  
 Thr Asp Ile Val Leu Ile Val Glu Gly Thr Glu Phe Pro Cys His  
 35 40 45  
 Lys Met Val Leu Ala Thr Cys Ser Ser Tyr Phe Arg Ala Met Phe  
 50 55 60  
 Met Ser Gly Leu Ser Glu Ser Lys Gln Thr His Val His Leu Arg  
 65 70 75  
 Asn Val Asp Ala Ala Thr Leu Gln Ile Ile Ile Thr Tyr Ala Tyr  
 80 85 90  
 Thr Gly Asn Leu Ala Met Asn Asp Ser Thr Val Glu Gln Leu Tyr  
 95 100 105  
 Glu Thr Ala Cys Phe Leu Gln Val Glu Asp Val Leu Gln Arg Cys  
 110 115 120  
 Arg Glu Tyr Leu Ile Lys Lys Ile Asn Ala Glu Asn Cys Val Arg  
 125 130 135  
 Leu Leu Ser Phe Ala Asp Leu Phe Ser Cys Glu Glu Leu Lys Gln  
 140 145 150  
 Ser Ala Lys Arg Met Val Glu His Lys Phe Thr Ala Val Tyr His  
 155 160 165  
 Gln Asp Ala Phe Met Gln Leu Ser His Asp Leu Leu Ile Asp Ile  
 170 175 180  
 Leu Ser Ser Asp Asn Leu Asn Val Glu Lys Glu Glu Thr Val Arg  
 185 190 195  
 Glu Ala Ala Met Leu Trp Leu Glu Tyr Asn Thr Glu Ser Arg Ser

Gln Tyr Leu Ser	200	Ser Val Leu Ser Gln	205	Arg Ile Asp Ala	210
	215		220		225
Ser Glu Val Thr	230	Gln Arg Ala Trp Phe	235	Gln Gly Leu Pro Pro	240
Asp Lys Ser Val	245	Val Val Gln Gly Leu	250	Lys Ser Met Pro	255
Phe Phe Lys Pro	260	Arg Leu Gly Met Thr	265	Lys Glu Glu Met Met	270
Phe Ile Glu Ala	275	Ser Ser Glu Asn Pro	280	Ser Ser Leu Tyr Ser	285
Val Cys Tyr Ser	290	Pro Gln Ala Glu Lys	295	Val Tyr Lys Leu Cys	300
Pro Pro Ala Asp	305	Leu His Lys Val Gly	310	Val Val Thr Pro	315
Asn Asp Ile Tyr	320	Ile Ala Gly Gly Gln	325	Val Pro Leu Lys Asn	330
Lys Thr Asn His	335	Ser Lys Thr Ser Lys	340	Gln Thr Ala Phe	345
Thr Val Asn Cys	350	Phe Tyr Trp Phe Asp	355	Gln Gln Asn Thr	360
Phe Pro Lys Thr	365	Pro Met Leu Phe Val	370	Ile Lys Pro Ser	375
Val Cys Cys Glu	380	Gly Tyr Ile Tyr Ala	385	Ile Gly Gly Asp Ser	390
Gly Gly Glu Leu	395	Asn Arg Arg Thr Val	400	Arg Tyr Asp Thr	405
Lys Asp Glu Trp	410	Thr Met Val Ser Pro	415	Pro Cys Ala Trp	420
Trp Ser Ala Ala	425	Val Val Val His Asp	430	Cys Ile Tyr Val Met	435
Leu Asn Leu Met	440	Tyr Cys Tyr Phe Pro	445	Arg Ser Asp Ser Trp	450
Glu Met Ala Met	455	Arg Gln Thr Ser Arg	460	Ser Phe Ala Ser Ala	465
Ala Phe Gly Asp	470	Lys Ile Phe Tyr Ile	475	Gly Gly Leu His Ile	480
Thr Asn Ser Gly	485	Ile Arg Leu Pro Ser	490	Gly Thr Val Asp Gly	495
Ser Val Thr Val	500	Glu Ile Tyr Asp Val	505	Asn Lys Asn Glu Trp	510
Met Ala Ala Asn	515	Ile Pro Ala Lys Arg	520	Tyr Ser Asp Pro Cys	525
Arg Ala Val Val	530	Ile Ser Asn Ser Leu	535	Cys Val Phe Met Arg	540
Thr His Leu Asn	545	Glu Arg Ala Lys Tyr	550	Val Thr Tyr Gln Tyr	555
Leu Glu Leu Asp	560	Arg Trp Ser Leu Arg	565	Gln His Ile Ser Glu	570
Val Leu Trp Asp	575	Leu Gly Arg Asp Phe	580	Arg Cys Thr Val Gly	585
Leu Tyr Pro Ser	590	Cys Leu Glu Glu Ser	595	Pro Trp Lys Pro Pro	600
Tyr Leu Phe Ser	605	Thr Asp Gly Thr Glu	610	Glu Phe Glu Leu Asp	615
Glu Met Val Ala	620	Leu Pro Pro Val			

&lt;210&gt; 2

&lt;211&gt; 99

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1455669CD1

&lt;400&gt; 2

Met Ala Ala Pro Ala Glu Pro Cys Ala Gly Gln Gly Val Trp Asn

1	5	10	15
Gln Thr Glu Pro Glu	Pro Ala Ala Thr Ser	Leu Leu Ser Leu Cys	
	20	25	30
Phe Leu Arg Thr Ala	Gly Val Trp Val Pro	Pro Met Tyr Leu Trp	
	35	40	45
Val Leu Gly Pro Ile	Tyr Leu Leu Phe Ile	His His His Gly Arg	
	50	55	60
Gly Tyr Leu Arg Met	Ser Pro Leu Phe Lys	Ala Lys Met Val Ala	
	65	70	75
Ala Ile Pro Gly Ser	Leu Glu Pro Gly Asn	Val Arg Gly Arg Gln	
	80	85	90
Gly Thr Gly Trp Asn	Leu Val Lys Ser		
	95		

&lt;210&gt; 3

&lt;211&gt; 374

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2084989CD1

&lt;400&gt; 3

Met Glu Ser Lys Met	Gly Glu Leu Pro Leu	Asp Ile Asn Ile Gln
1	5	10
Glu Pro Arg Trp Asp	Gln Ser Thr Phe Leu	Gly Arg Ala Arg His
	20	25
Phe Phe Thr Val Thr	Asp Pro Arg Asn Leu	Leu Leu Ser Gly Ala
	35	40
Gln Leu Glu Ala Ser	Arg Asn Ile Val Gln	Asn Tyr Arg Ala Gly
	50	55
Val Val Thr Pro Gly	Ile Thr Glu Asp Gln	Leu Trp Arg Ala Lys
	65	70
Tyr Val Tyr Asp Ser	Ala Phe His Pro Asp	Thr Gly Glu Lys Val
	80	85
Val Leu Ile Gly Arg	Met Ser Ala Gln Val	Pro Met Asn Met Thr
	95	100
Ile Thr Gly Cys Met	Leu Thr Phe Tyr Arg	Lys Thr Pro Thr Val
	110	115
Val Phe Trp Gln Trp	Val Asn Gln Ser Phe	Asn Ala Ile Val Asn
	125	130
Tyr Ser Asn Arg Ser	Gly Asp Thr Pro Ile	Thr Val Arg Gln Leu
	140	145
Gly Thr Ala Tyr Val	Ser Ala Thr Thr Gly	Ala Val Ala Thr Ala
	155	160
Leu Gly Leu Lys Ser	Leu Thr Lys His Leu	Pro Pro Leu Val Gly
	170	175
Arg Phe Val Pro Phe	Ala Ala Val Ala Ala	Asn Cys Ile Asn
	185	190
Ile Pro Leu Met Arg	Gln Arg Glu Leu Gln	Val Gly Ile Pro Val
	200	205
Ala Asp Glu Ala Gly	Gln Arg Leu Gly Tyr	Ser Val Thr Ala Ala
	215	220
Lys Gln Gly Ile Phe	Gln Val Val Ile Ser	Arg Ile Cys Met Ala
	230	235
Ile Pro Ala Met Ala	Ile Pro Pro Leu Ile	Met Asp Thr Leu Glu
	245	250
Lys Lys Asp Phe Leu	Lys Val Gly Asp Cys	Thr Ser Leu Val Leu
	260	265
Glu Trp Ala Met Ala	Gly Arg Ser Asp Gln	Ala Pro Thr Leu Ser
	275	280
Pro Ala Ser Pro Asp	Ser Leu Arg Leu Ala	Ser Pro Ser Pro Asp
	290	295
Pro Cys Thr Ala Ser	Ser Thr Phe Val His	Ser Ala Arg Met Asn
	305	310
Trp Ala Gly Val Lys	Glu Leu Cys Arg Gly	Arg Arg Arg Gly Gln
	320	325
Arg Lys Glu Thr Asn	Phe Ile Ser Val Thr	Pro Val Ala Ser Asp



	335		340		345
Thr Gln Lys Gly	Thr Val Ile Val Met	Leu Asp Leu Met Leu	Ile		
	350		355		360
Leu Leu Pro Pro	Ser Ala Ser Ile Leu	Arg Gly Thr His Gly			
	365		370		

&lt;210&gt; 4

&lt;211&gt; 271

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2501034CD1

&lt;400&gt; 4

Met Gly Asn Gly Gly	Arg Ser Gly Leu Gln	Gln Gly Lys Gly Asn
1	5	10
Val Asp Gly Val Ala	Ala Thr Pro Thr Ala	Ala Ser Ala Ser Cys
	20	25
Gln Tyr Arg Cys Ile	Glu Cys Asn Gln Glu	Ala Lys Glu Leu Tyr
	35	40
Arg Asp Tyr Asn His	Gly Val Leu Lys Ile	Thr Ile Cys Lys Ser
	50	55
Cys Gln Lys Pro Val	Asp Lys Tyr Ile Glu	Tyr Asp Pro Val Ile
	65	70
Ile Leu Ile Asn Ala	Ile Leu Cys Lys Ala	Gln Ala Tyr Arg His
	80	85
Ile Leu Phe Asn Thr	Gln Ile Asn Ile His	Gly Lys Leu Cys Ile
	95	100
Phe Cys Leu Leu Cys	Glu Ala Tyr Leu Arg	Trp Trp Gln Leu Gln
	110	115
Asp Ser Asn Gln Asn	Thr Ala Pro Asp Asp	Leu Ile Arg Tyr Ala
	125	130
Lys Glu Trp Asp Phe	Tyr Arg Met Phe Ala	Ile Ala Ala Leu Glu
	140	145
Gln Thr Ala Tyr Phe	Ile Gly Ile Phe Thr	Phe Leu Trp Val Glu
	155	160
Arg Pro Met Thr Ala	Lys Lys Lys Pro Asn	Phe Ile Leu Leu Leu
	170	175
Lys Ala Leu Leu Leu	Ser Ser Tyr Gly Lys	Leu Leu Leu Ile Pro
	185	190
Ala Val Ile Trp Glu	His Asp Tyr Thr Ser	Val Cys Leu Lys Leu
	200	205
Ile Lys Val Phe Val	Leu Thr Ser Asn Phe	Gln Ala Ile Arg Val
	215	220
Thr Leu Asn Ile Asn	Arg Lys Leu Ser Phe	Leu Ala Val Leu Ser
	230	235
Gly Leu Leu Leu Glu	Ser Ile Met Val Tyr	Phe Phe Gln Ser Met
	245	250
Glu Trp Asp Val Gly	Ser Asp Tyr Ala Ile	Phe Lys Ser Gln Asp
	260	265
Phe		270

&lt;210&gt; 5

&lt;211&gt; 323

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2745212CD1

&lt;400&gt; 5

Met Ala Pro Lys Gln	Asp Pro Lys Pro Lys	Phe Gln Glu Gly Glu
1	5	10
Arg Val Leu Cys Phe	His Gly Pro Leu Leu	Tyr Glu Ala Lys Cys
	20	25
Val Lys Val Ala Ile	Lys Asp Lys Gln Val	Lys Tyr Phe Ile His

	35		40		45
Tyr Ser Gly Trp	Asn Lys Asn Trp Asp	Glu Trp Val Pro Glu Ser			
	50		55		60
Arg Val Leu Lys	Tyr Val Asp Thr Asn Leu	Gln Lys Gln Arg Glu			
	65		70		75
Leu Gln Lys Ala	Asn Gln Glu Gln Tyr Ala	Glu Gly Lys Met Arg			
	80		85		90
Gly Ala Ala Pro	Gly Lys Lys Thr Ser Gly	Leu Gln Gln Lys Asn			
	95		100		105
Val Glu Val Lys	Thr Lys Lys Asn Lys Gln	Lys Thr Pro Gly Asn			
	110		115		120
Gly Asp Gly Gly	Ser Thr Ser Glu Thr Pro	Gln Pro Pro Arg Lys			
	125		130		135
Lys Arg Ala Arg	Val Asp Pro Thr Val Glu	Asn Glu Glu Thr Phe			
	140		145		150
Met Asn Arg Val	Glu Val Lys Val Lys Ile	Pro Glu Glu Leu Lys			
	155		160		165
Pro Trp Leu Val	Asp Asp Trp Asp Leu Ile	Thr Arg Gln Lys Gln			
	170		175		180
Leu Phe Tyr Leu	Pro Ala Lys Lys Asn Val	Asp Ser Ile Leu Glu			
	185		190		195
Asp Tyr Ala Asn	Tyr Lys Lys Ser Arg Gly	Asn Thr Asp Asn Lys			
	200		205		210
Glu Tyr Ala Val	Asn Glu Val Val Ala Gly	Ile Lys Glu Tyr Phe			
	215		220		225
Asn Val Met Leu	Gly Thr Gln Leu Leu Tyr	Lys Phe Glu Arg Pro			
	230		235		240
Gln Tyr Ala Glu	Ile Leu Ala Asp His Pro	Asp Ala Pro Met Ser			
	245		250		255
Gln Val Tyr Gly	Ala Pro His Leu Leu Arg	Leu Phe Val Arg Ile			
	260		265		270
Gly Ala Met Leu	Ala Tyr Thr Pro Leu Asp	Glu Lys Ser Leu Ala			
	275		280		285
Leu Leu Leu Asn	Tyr Leu His Asp Phe Leu	Lys Tyr Leu Ala Lys			
	290		295		300
Asn Ser Ala Thr	Leu Phe Ser Ala Ser Asp	Tyr Glu Val Ala Pro			
	305		310		315
Pro Glu Tyr His	Arg Lys Ala Val				
	320				

&lt;210&gt; 6

&lt;211&gt; 274

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4833111CD1

&lt;400&gt; 6

Met Asp Arg Pro Gly	Phe Val Ala Ala Leu Val Ala Gly Gly Val
1	5 10 15
Ala Gly Val Ser Val	Asp Leu Ile Leu Phe Pro Leu Asp Thr Ile
	20 25 30
Lys Thr Arg Leu Gln	Ser Pro Gln Gly Phe Ser Lys Ala Gly Gly
	35 40 45
Phe His Gly Ile Tyr	Ala Gly Val Pro Ser Ala Ala Ile Gly Ser
	50 55 60
Phe Pro Asn Ala Ala	Ala Phe Phe Ile Thr Tyr Glu Tyr Val Lys
	65 70 75
Trp Phe Leu His Ala	Asp Ser Ser Ser Tyr Leu Thr Pro Met Lys
	80 85 90
His Met Leu Ala Ala	Ser Ala Gly Glu Val Val Ala Cys Leu Ile
	95 100 105
Arg Val Pro Ser Glu	Val Val Lys Gln Arg Ala Gln Val Ser Ala
	110 115 120
Ser Thr Arg Thr Phe	Gln Ile Phe Ser Asn Ile Leu Tyr Glu Glu
	125 130 135
Gly Ile Gln Gly Leu	Tyr Arg Gly Tyr Lys Ser Thr Val Leu Arg

	140		145		150
Glu Ile Pro Phe	Ser Leu Val Gln Phe	Pro Leu Trp Glu Ser	Leu		
	155		160		165
Lys Ala Leu Trp	Ser Trp Arg Gln Asp	His Val Val Asp Ser	Trp		
	170		175		180
Gln Ser Ala Val	Cys Gly Ala Phe Ala	Gly Gly Phe Ala Ala	Ala		
	185		190		195
Val Thr Thr Pro	Leu Asp Val Ala Lys	Thr Arg Ile Thr Leu	Ala		
	200		205		210
Lys Ala Gly Ser	Ser Thr Ala Asp Gly	Asn Val Leu Ser Val	Leu		
	215		220		225
His Gly Val Trp	Arg Ser Gln Gly Leu	Ala Gly Leu Phe Ala	Gly		
	230		235		240
Val Phe Pro Arg	Met Ala Ala Ile Ser	Leu Gly Gly Phe Ile	Phe		
	245		250		255
Leu Gly Ala Tyr	Asp Arg Thr His Ser	Leu Leu Leu Glu Val	Gly		
	260		265		270
Arg Lys Ser Pro					

&lt;210&gt; 7

&lt;211&gt; 291

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 876677CD1

&lt;400&gt; 7

Met Asp Ser Arg	Val Ser Ser Pro	Glu Lys Gln Asp	Lys Glu Asn
1	5	10	15
Phe Val Gly Val	Asn Asn Lys Arg	Leu Gly Val Cys	Gly Trp Ile
	20	25	30
Leu Phe Ser Leu	Ser Phe Leu Leu	Val Ile Ile Thr	Phe Pro Ile
	35	40	45
Ser Ile Trp Met	Cys Leu Lys Ile	Ile Lys Glu Tyr	Glu Arg Ala
	50	55	60
Val Val Phe Arg	Leu Gly Arg Ile	Gln Ala Asp Lys	Ala Lys Gly
	65	70	75
Pro Gly Leu Ile	Leu Val Leu Pro	Cys Ile Asp Val	Phe Val Lys
	80	85	90
Val Asp Leu Arg	Thr Val Thr Cys	Asn Ile Pro Pro	Gln Glu Ile
	95	100	105
Leu Thr Arg Asp	Ser Val Thr Thr	Gln Val Asp Gly	Val Val Tyr
	110	115	120
Tyr Arg Ile Tyr	Ser Ala Val Ser	Ala Val Ala Asn	Val Asn Asp
	125	130	135
Val His Gln Ala	Thr Phe Leu Leu	Ala Gln Thr Thr	Leu Arg Asn
	140	145	150
Val Leu Gly Thr	Gln Thr Leu Ser	Gln Ile Leu Ala	Gly Arg Glu
	155	160	165
Glu Ile Ala His	Ser Ile Gln Thr	Leu Leu Asp Asp	Ala Thr Glu
	170	175	180
Leu Trp Gly Ile	Arg Val Ala Arg	Val Glu Ile Lys	Asp Val Arg
	185	190	195
Ile Pro Val Gln	Leu Gln Arg Ser	Met Ala Ala Glu	Ala Glu Ala
	200	205	210
Thr Arg Glu Ala	Arg Ala Lys Val	Leu Ala Ala Glu	Gly Glu Met
	215	220	225
Asn Ala Ser Lys	Ser Leu Lys Ser	Ala Ser Met Val	Leu Ala Glu
	230	235	240
Ser Pro Ile Ala	Leu Gln Leu Arg	Tyr Leu Gln Thr	Leu Ser Thr
	245	250	255
Val Ala Thr Glu	Lys Asn Ser Thr	Ile Val Phe Pro	Leu Pro Met
	260	265	270
Asn Ile Leu Glu	Gly Ile Gly Gly	Val Ser Tyr Asp	Asn His Lys
	275	280	285
Lys Leu Pro Asn	Lys Ala		

290

<210> 8  
 <211> 381  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2326143CD1

<400> 8  
 Met Ser Arg His Glu Gly Val Ser Cys Asp Ala Cys Leu Lys Gly  
 1 5 10 15  
 Asn Phe Arg Gly Arg Arg Tyr Lys Cys Leu Ile Cys Tyr Asp Tyr  
 20 25 30  
 Asp Leu Cys Ala Ser Cys Tyr Glu Ser Gly Ala Thr Thr Thr Arg  
 35 40 45  
 His Thr Thr Asp His Pro Met Gln Cys Ile Leu Thr Arg Val Asp  
 50 55 60  
 Phe Asp Leu Tyr Tyr Gly Gly Glu Ala Phe Ser Val Glu Gln Pro  
 65 70 75  
 Gln Ser Phe Thr Cys Pro Tyr Cys Gly Lys Met Gly Tyr Thr Glu  
 80 85 90  
 Thr Ser Leu Gln Glu His Val Thr Ser Glu His Ala Glu Thr Ser  
 95 100 105  
 Thr Glu Val Ile Cys Pro Ile Cys Ala Ala Leu Pro Gly Gly Asp  
 110 115 120  
 Pro Asn His Val Thr Asp Asp Phe Ala Ala His Leu Thr Leu Glu  
 125 130 135  
 His Arg Ala Pro Arg Asp Leu Asp Glu Ser Ser Gly Val Arg His  
 140 145 150  
 Val Arg Arg Met Phe His Pro Gly Arg Gly Leu Gly Gly Pro Arg  
 155 160 165  
 Ala Arg Arg Ser Asn Met His Phe Thr Ser Ser Ser Thr Gly Gly  
 170 175 180  
 Leu Ser Ser Ser Gln Ser Ser Tyr Ser Pro Ser Asn Arg Glu Ala  
 185 190 195  
 Met Asp Pro Ile Ala Glu Leu Leu Ser Gln Leu Ser Gly Val Arg  
 200 205 210  
 Arg Ser Ala Gly Gly Gln Leu Asn Ser Ser Gly Pro Ser Ala Ser  
 215 220 225  
 Gln Leu Gln Gln Leu Gln Met Gln Leu Gln Leu Glu Arg Gln His  
 230 235 240  
 Ala Gln Ala Ala Arg Gln Gln Leu Glu Thr Ala Arg Asn Ala Thr  
 245 250 255  
 Arg Arg Thr Asn Thr Ser Ser Val Thr Thr Thr Ile Thr Gln Ser  
 260 265 270  
 Thr Ala Thr Thr Asn Ile Ala Asn Thr Glu Ser Ser Gln Gln Thr  
 275 280 285  
 Leu Gln Asn Ser Gln Phe Leu Leu Thr Arg Leu Asn Asp Pro Lys  
 290 295 300  
 Met Ser Glu Thr Glu Arg Gln Ser Met Glu Ser Glu Arg Ala Asp  
 305 310 315  
 Arg Ser Leu Phe Val Gln Glu Leu Leu Leu Ser Thr Leu Val Arg  
 320 325 330  
 Glu Glu Ser Ser Ser Ser Asp Glu Asp Asp Arg Gly Glu Met Ala  
 335 340 345  
 Asp Phe Gly Ala Met Gly Cys Val Asp Ile Met Pro Leu Asp Val  
 350 355 360  
 Ala Leu Glu Asn Leu Asn Leu Lys Glu Ser Asn Lys Gly Asn Glu  
 365 370 375  
 Pro Pro Pro Pro Pro Leu  
 380

<210> 9  
 <211> 190  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2786302CD1

<400> 9  
 Met Lys Tyr Gly Asn Glu Ile Met Asn Lys Asp Pro Val Phe Arg  
 1 5 10 15  
 Ile Ser Pro Arg Ser Arg Glu Thr His Pro Asn Pro Glu Glu Pro  
 20 25 30  
 Glu Glu Glu Asp Glu Asp Val Gln Ala Glu Arg Val Gln Ala Ala  
 35 40 45  
 Asn Ala Leu Thr Ala Pro Asn Leu Glu Glu Glu Pro Val Ile Thr  
 50 55 60  
 Ala Ser Cys Leu His Lys Glu Tyr Tyr Glu Thr Lys Lys Ser Cys  
 65 70 75  
 Phe Ser Thr Arg Lys Lys Lys Ile Ala Ile Arg Asn Val Ser Phe  
 80 85 90  
 Cys Val Lys Lys Gly Glu Val Leu Gly Leu Leu Gly His Asn Gly  
 95 100 105  
 Ala Gly Lys Ser Thr Ser Ile Lys Met Ile Thr Gly Cys Thr Lys  
 110 115 120  
 Pro Thr Ala Gly Val Val Val Leu Gln Gly Ser Arg Ala Ser Val  
 125 130 135  
 Arg Gln Gln His Asp Asn Ser Leu Lys Phe Leu Gly Tyr Cys Pro  
 140 145 150  
 Gln Glu Asn Ser Leu Trp Pro Lys Leu Thr Met Lys Glu His Leu  
 155 160 165  
 Glu Leu Tyr Ala Ala Val Glu Arg Leu Gly Gln Lys Arg Cys Cys  
 170 175 180  
 Ser Gln Tyr Phe Thr Ile Gly Gly Arg Ser  
 185 190

<210> 10  
 <211> 297  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3735780CD1

<400> 10  
 Met Met Asp Ser Glu Ala His Glu Lys Arg Pro Pro Ile Leu Thr  
 1 5 10 15  
 Ser Ser Lys Gln Asp Ile Ser Pro His Ile Thr Asn Val Gly Glu  
 20 25 30  
 Met Lys His Tyr Leu Cys Gly Cys Cys Ala Ala Phe Asn Asn Val  
 35 40 45  
 Ala Ile Thr Phe Pro Ile Gln Lys Val Leu Phe Arg Gln Gln Leu  
 50 55 60  
 Tyr Gly Ile Lys Thr Arg Asp Ala Ile Leu Gln Leu Arg Arg Asp  
 65 70 75  
 Gly Phe Arg Asn Leu Tyr Arg Gly Ile Leu Pro Pro Leu Met Gln  
 80 85 90  
 Lys Thr Thr Thr Leu Ala Leu Met Phe Gly Leu Tyr Glu Asp Leu  
 95 100 105  
 Ser Cys Leu Leu His Lys His Val Ser Ala Pro Glu Phe Ala Thr  
 110 115 120  
 Ser Gly Val Ala Ala Val Leu Ala Gly Thr Thr Glu Ala Ile Phe  
 125 130 135  
 Thr Pro Leu Glu Arg Val Gln Thr Leu Leu Gln Asp His Lys His  
 140 145 150  
 His Asp Lys Phe Thr Asn Thr Tyr Gln Ala Phe Lys Ala Leu Lys  
 155 160 165  
 Cys His Gly Ile Gly Glu Tyr Tyr Arg Gly Leu Val Pro Ile Leu  
 170 175 180  
 Phe Arg Asn Gly Leu Ser Asn Val Leu Phe Phe Gly Leu Arg Gly  
 185 190 195  
 Pro Ile Lys Glu His Leu Pro Thr Ala Thr Thr His Ser Ala His

200	205	210
Leu Val Asn Asp Phe Ile Cys Gly Gly	Leu Leu Gly Ala Met Leu	
215	220	225
Gly Phe Leu Phe Phe Pro Ile Asn Val	Val Lys Thr Arg Ile Gln	
230	235	240
Ser Gln Ile Gly Gly Glu Phe Gln Ser	Phe Pro Lys Val Phe Gln	
245	250	255
Lys Ile Trp Leu Glu Arg Asp Arg Lys	Leu Ile Asn Leu Phe Arg	
260	265	270
Gly Ala His Leu Asn Tyr His Arg Ser	Leu Ile Ser Trp Gly Ile	
275	280	285
Ile Asn Ala Thr Tyr Glu Phe Leu Leu	Lys Val Ile	
290	295	

<210> 11  
 <211> 89  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 039026CD1

<400> 11	
Met Ala Ala Gln Ile Pro Glu Ser Asp Gln Ile Lys Gln Phe Lys	
1 5 10 15	
Glu Phe Leu Gly Thr Tyr Asn Lys Leu Thr Glu Thr Cys Phe Leu	
20 25 30	
Asp Cys Val Lys Asp Phe Thr Thr Arg Glu Val Lys Pro Glu Glu	
35 40 45	
Thr Thr Cys Ser Glu His Cys Leu Gln Lys Tyr Leu Lys Met Thr	
50 55 60	
Gln Arg Ile Ser Met Arg Phe Gln Glu Tyr His Ile Gln Gln Asn	
65 70 75	
Glu Ala Leu Ala Ala Lys Ala Gly Leu Leu Gly Gln Pro Arg	
80 85	

<210> 12  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 260607CD1

<400> 12	
Met Ala Leu Ile Pro Ser Arg Val Trp Leu Pro Phe Ala Val Trp	
1 5 10 15	
Val Val Asp Ser Ala Pro Val Arg Gly Leu Val Arg Arg Glu Pro	
20 25 30	
Phe Leu Arg Thr Gly Ser Phe Ile Ala Leu Phe Tyr Phe Pro Pro	
35 40 45	
Leu Leu Pro Val Leu Ile Asn Leu Phe Ser Phe Phe Leu Thr Pro	
50 55 60	
Ser Phe Trp Arg Gln Leu Gly Ala Ile Leu Val Tyr Ala Ser Leu	
65 70 75	
Leu Ala Glu Lys Thr Pro Phe Lys Thr Gln Arg Thr Leu Glu Gly	
80 85 90	
Asp Ala Leu Val Gly Ser Val Ser Ile Phe Leu Cys Ala Lys Asp	
95 100 105	
Arg Gln Thr Glu Ala Glu Arg Gly Cys Ser	
110 115	

<210> 13  
 <211> 675  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature

&lt;223&gt; Incyte ID No: 1429651CD1

&lt;400&gt; 13

Met	Glu	Ser	Gly	Thr	Ser	Ser	Pro	Gln	Pro	Pro	Gln	Leu	Asp	Pro
1				5					10					15
Leu	Asp	Ala	Phe	Pro	Gln	Lys	Gly	Leu	Glu	Pro	Gly	Asp	Ile	Ala
				20					25					30
Val	Leu	Val	Leu	Tyr	Phe	Leu	Phe	Val	Leu	Ala	Val	Gly	Leu	Trp
				35					40					45
Ser	Thr	Val	Lys	Thr	Lys	Arg	Asp	Thr	Val	Lys	Gly	Tyr	Phe	Leu
				50					55					60
Ala	Gly	Gly	Asp	Met	Val	Trp	Trp	Pro	Val	Gly	Ala	Ser	Leu	Phe
				65					70					75
Ala	Ser	Asn	Val	Gly	Ser	Gly	His	Phe	Ile	Gly	Leu	Ala	Gly	Ser
				80					85					90
Gly	Ala	Ala	Thr	Gly	Ile	Ser	Val	Ser	Ala	Tyr	Glu	Leu	Asn	Gly
				95					100					105
Leu	Phe	Ser	Val	Leu	Met	Leu	Ala	Trp	Ile	Phe	Leu	Pro	Ile	Tyr
				110					115					120
Ile	Ala	Gly	Gln	Val	Thr	Thr	Met	Pro	Glu	Tyr	Leu	Arg	Lys	Arg
				125					130					135
Phe	Gly	Gly	Ile	Arg	Ile	Pro	Ile	Ile	Leu	Ala	Val	Leu	Tyr	Leu
				140					145					150
Phe	Ile	Tyr	Ile	Phe	Thr	Lys	Ile	Ser	Val	Asp	Met	Tyr	Ala	Gly
				155					160					165
Ala	Ile	Phe	Ile	Gln	Gln	Ser	Leu	His	Leu	Asp	Leu	Tyr	Leu	Ala
				170					175					180
Ile	Val	Gly	Leu	Leu	Ala	Ile	Thr	Ala	Val	Tyr	Thr	Val	Ala	Gly
				185					190					195
Gly	Leu	Ala	Ala	Val	Ile	Tyr	Thr	Asp	Ala	Leu	Gln	Thr	Leu	Ile
				200					205					210
Met	Leu	Ile	Gly	Ala	Leu	Thr	Leu	Met	Gly	Tyr	Ser	Phe	Ala	Ala
				215					220					225
Val	Gly	Gly	Met	Glu	Gly	Leu	Lys	Glu	Lys	Tyr	Phe	Leu	Ala	Leu
				230					235					240
Ala	Ser	Asn	Arg	Ser	Glu	Asn	Ser	Ser	Cys	Gly	Leu	Pro	Arg	Glu
				245					250					255
Asp	Ala	Phe	His	Ile	Phe	Arg	Asp	Pro	Leu	Thr	Ser	Asp	Leu	Pro
				260					265					270
Trp	Pro	Gly	Val	Leu	Phe	Gly	Met	Ser	Ile	Pro	Ser	Leu	Trp	Tyr
				275					280					285
Trp	Cys	Thr	Asp	Gln	Val	Ile	Val	Gln	Arg	Thr	Leu	Ala	Ala	Lys
				290					295					300
Asn	Leu	Ser	His	Ala	Lys	Gly	Gly	Ala	Leu	Met	Ala	Ala	Tyr	Leu
				305					310					315
Lys	Val	Leu	Pro	Leu	Phe	Ile	Met	Val	Phe	Pro	Gly	Met	Val	Ser
				320					325					330
Arg	Ile	Leu	Phe	Pro	Asp	Gln	Val	Ala	Cys	Ala	Asp	Pro	Glu	Ile
				335					340					345
Cys	Gln	Lys	Ile	Cys	Ser	Asn	Pro	Ser	Gly	Cys	Ser	Asp	Ile	Ala
				350					355					360
Tyr	Pro	Lys	Leu	Val	Leu	Glu	Leu	Leu	Pro	Thr	Gly	Leu	Arg	Gly
				365					370					375
Leu	Met	Met	Ala	Val	Met	Val	Ala	Ala	Leu	Met	Ser	Ser	Leu	Thr
				380					385					390
Ser	Ile	Phe	Asn	Ser	Ala	Ser	Thr	Ile	Phe	Thr	Met	Asp	Leu	Trp
				395					400					405
Asn	His	Leu	Arg	Pro	Arg	Ala	Ser	Glu	Lys	Glu	Leu	Met	Ile	Val
				410					415					420
Gly	Arg	Val	Phe	Val	Leu	Leu	Leu	Val	Leu	Val	Ser	Ile	Leu	Trp
				425					430					435
Ile	Pro	Val	Val	Gln	Ala	Ser	Gln	Gly	Gly	Gln	Leu	Phe	Ile	Tyr
				440					445					450
Ile	Gln	Ser	Ile	Ser	Ser	Tyr	Leu	Gln	Pro	Pro	Val	Ala	Val	Val
				455					460					465
Phe	Ile	Met	Gly	Cys	Phe	Trp	Lys	Arg	Thr	Asn	Glu	Lys	Gly	Ala
				470					475					480
Phe	Trp	Gly	Leu	Ile	Ser	Gly	Leu	Leu	Leu	Gly	Leu	Val	Arg	Leu

	485		490		495
Val Leu Asp Phe	Ile Tyr Val Gln Pro	Arg Cys Asp Gln Pro	Asp		
	500		505		510
Glu Arg Pro Val	Leu Val Lys Ser Ile	His Tyr Leu Tyr Phe	Ser		
	515		520		525
Met Ile Leu Ser	Thr Val Thr Leu Ile	Thr Val Ser Thr Val	Ser		
	530		535		540
Trp Phe Thr Glu	Pro Pro Ser Lys Glu	Met Val Ser His Leu	Thr		
	545		550		555
Trp Phe Thr Arg	His Asp Pro Val Val	Gln Lys Glu Gln Ala	Pro		
	560		565		570
Pro Ala Ala Pro	Leu Ser Leu Thr Leu	Ser Gln Asn Gly Met	Pro		
	575		580		585
Glu Ala Ser Ser	Ser Ser Ser Val Gln	Phe Glu Met Val Gln	Glu		
	590		595		600
Asn Thr Ser Lys	Thr His Ser Cys Asp	Met Thr Pro Lys Gln	Ser		
	605		610		615
Lys Val Val Lys	Ala Ile Leu Trp Leu	Cys Gly Ile Gln Glu	Lys		
	620		625		630
Gly Lys Glu Glu	Leu Pro Ala Arg Ala	Glu Ala Ile Ile Val	Ser		
	635		640		645
Leu Glu Glu Asn	Pro Leu Val Lys Thr	Leu Leu Asp Val Asn	Leu		
	650		655		660
Ile Phe Cys Val	Ser Cys Ala Ile Phe	Ile Trp Gly Tyr Phe	Ala		
	665		670		675

&lt;210&gt; 14

&lt;211&gt; 320

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2069971CD1

&lt;400&gt; 14

Met Tyr His Cys His	Ser Gly Ser Lys Pro	Thr Glu Lys Gly Ala
1	5	10
Asn Glu Tyr Ala Tyr	Ala Lys Trp Lys Leu	Cys Ser Ala Ser Ala
	20	25
Ile Cys Phe Ile Phe	Met Ile Ala Glu Val	Val Gly Gly His Ile
	35	40
Ala Gly Ser Leu Ala	Val Val Thr Asp Ala	Ala His Leu Leu Ile
	50	55
Asp Leu Thr Ser Phe	Leu Leu Ser Leu Phe	Ser Leu Trp Leu Ser
	65	70
Ser Lys Pro Pro Ser	Lys Arg Leu Thr Phe	Gly Trp His Arg Ala
	80	85
Glu Ile Leu Gly Ala	Leu Leu Ser Ile Leu	Cys Ile Trp Val Val
	95	100
Thr Gly Val Leu Val	Tyr Leu Ala Cys Glu	Arg Leu Leu Tyr Pro
	110	115
Asp Tyr Gln Ile Gln	Ala Thr Val Met Ile	Ile Val Ser Ser Cys
	125	130
Ala Val Ala Ala Asn	Ile Val Leu Thr Val	Val Val Leu His Gln Arg
	140	145
Cys Leu Gly His Asn	His Lys Glu Val Gln	Ala Asn Ala Ser Val
	155	160
Arg Ala Ala Phe Val	His Ala Leu Gly Asp	Leu Phe Gln Ser Ile
	170	175
Ser Val Leu Ile Ser	Ala Leu Ile Ile Tyr	Phe Lys Pro Glu Tyr
	185	190
Lys Ile Ala Asp Pro	Ile Cys Thr Phe Ile	Phe Ser Ile Leu Val
	200	205
Leu Ala Ser Thr Ile	Thr Ile Leu Lys Asp	Phe Ser Ile Leu Leu
	215	220
Met Glu Gly Val Pro	Lys Ser Leu Asn Tyr	Ser Gly Val Lys Glu
	230	235



```

Leu Ile Leu Ala Val Asp Gly Val Leu Ser Val His Ser Leu His
245 250 255
Ile Trp Ser Leu Thr Met Asn Gln Val Ile Leu Ser Ala His Val
260 265 270
Ala Thr Ala Ala Ser Arg Asp Ser Gln Val Val Arg Arg Glu Ile
275 280 285
Ala Lys Ala Leu Ser Lys Ser Phe Thr Met His Ser Leu Thr Ile
290 295 300
Gln Met Glu Ser Pro Val Asp Gln Asp Pro Asp Cys Leu Phe Cys
305 310 315
Glu Asp Pro Cys Asp
320

```

<210> 15  
 <211> 462  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2329339CD1

```

<400> 15
Met Ala Glu Glu Gln Glu Phe Thr Gln Leu Cys Lys Leu Pro Ala
1 5 10 15
Gln Pro Ser His Pro His Cys Val Asn Asn Thr Tyr Arg Ser Ala
20 25 30
Gln His Ser Gln Ala Leu Leu Arg Gly Leu Leu Ala Leu Arg Asp
35 40 45
Ser Gly Ile Leu Phe Asp Val Val Leu Val Val Glu Gly Arg His
50 55 60
Ile Glu Ala His Arg Ile Leu Leu Ala Ala Ser Cys Asp Tyr Phe
65 70 75
Arg Gly Met Phe Ala Gly Gly Leu Lys Glu Met Glu Gln Glu Glu
80 85 90
Val Leu Ile His Gly Val Ser Tyr Asn Ala Met Cys Gln Ile Leu
95 100 105
His Phe Ile Tyr Thr Ser Glu Leu Glu Leu Ser Leu Ser Asn Val
110 115 120
Gln Glu Thr Leu Val Ala Ala Cys Gln Leu Gln Ile Pro Glu Ile
125 130 135
Ile His Phe Cys Cys Asp Phe Leu Met Ser Trp Val Asp Glu Glu
140 145 150
Asn Ile Leu Asp Val Tyr Arg Leu Ala Glu Leu Phe Asp Leu Ser
155 160 165
Arg Leu Thr Glu Gln Leu Asp Thr Tyr Ile Leu Lys Asn Phe Val
170 175 180
Ala Phe Ser Arg Thr Asp Lys Tyr Arg Gln Leu Pro Leu Glu Lys
185 190 195
Val Tyr Ser Leu Leu Ser Ser Asn Arg Leu Glu Val Ser Cys Glu
200 205 210
Thr Glu Val Tyr Glu Gly Ala Leu Leu Tyr His Tyr Ser Leu Glu
215 220 225
Gln Val Gln Ala Asp Gln Ile Ser Leu His Glu Pro Pro Lys Leu
230 235 240
Leu Glu Thr Val Arg Phe Pro Leu Met Glu Ala Glu Val Leu Gln
245 250 255
Arg Leu His Asp Lys Leu Asp Pro Ser Pro Leu Arg Asp Thr Val
260 265 270
Ala Ser Gly Leu Met Tyr His Arg Asn Glu Ser Leu Gln Pro Ser
275 280 285
Leu Gln Ser Pro Gln Thr Glu Leu Arg Ser Asp Phe Gln Cys Val
290 295 300
Val Gly Phe Gly Gly Ile His Ser Thr Pro Ser Thr Val Leu Ser
305 310 315
Asp Gln Ala Lys Tyr Leu Asn Pro Leu Leu Gly Glu Trp Lys His
320 325 330
Phe Thr Ala Ser Leu Ala Pro Arg Met Ser Asn Gln Gly Ile Ala
335 340 345

```

```

Val Leu Asn Asn Phe Val Tyr Leu Ile Gly Gly Asp Asn Asn Val
350 355 360
Gln Gly Phe Arg Ala Glu Ser Arg Cys Trp Arg Tyr Asp Pro Arg
365 370 375
His Asn Arg Trp Phe Gln Ile Gln Ser Leu Gln Gln Glu His Ala
380 385 390
Asp Leu Ser Val Cys Val Val Gly Arg Tyr Ile Tyr Ala Val Ala
395 400 405
Gly Arg Asp Tyr His Asn Asp Leu Asn Ala Val Glu Arg Tyr Asp
410 415 420
Pro Ala Thr Asn Ser Trp Ala Tyr Val Ala Pro Leu Lys Arg Glu
425 430 435
Val Tyr Ala His Ala Gly Ala Thr Leu Glu Gly Lys Met Tyr Ile
440 445 450
Thr Cys Gly Arg Lys Leu Ile Pro Phe Ser Glu Gly
455 460

```

&lt;210&gt; 16

&lt;211&gt; 98

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2540219CD1

&lt;400&gt; 16

```

Met Arg Ala Cys Ala Val Trp Leu Ala Gly Gly Met Ala Gly Ala
1 5 10 15
Ile Ser Trp Gly Thr Ala Thr Pro Met Asp Val Val Lys Ser Arg
20 25 30
Leu Gln Ala Asp Gly Val Tyr Leu Asn Lys Tyr Lys Gly Val Leu
35 40 45
Asp Cys Ile Ser Gln Ser Tyr Gln Lys Glu Gly Leu Lys Val Phe
50 55 60
Phe Arg Gly Ile Thr Val Asn Ala Val Arg Gly Phe Pro Met Ser
65 70 75
Ala Ala Met Phe Leu Gly Tyr Glu Leu Ser Leu Gln Ala Ile Arg
80 85 90
Gly Asp His Ala Val Thr Ser Pro
95

```

&lt;210&gt; 17

&lt;211&gt; 748

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2722462CD1

&lt;400&gt; 17

```

Met Asn Tyr Gln Glu Ala Ala Ile Tyr Leu Gln Glu Gly Glu Asn
1 5 10 15
Asn Asp Lys Phe Phe Thr His Pro Lys Asp Ala Lys Ala Leu Ala
20 25 30
Ala Tyr Leu Phe Ala His Asn His Leu Phe Tyr Leu Met Glu Leu
35 40 45
Ala Thr Ala Leu Leu Leu Leu Leu Ser Leu Cys Glu Ala Pro
50 55 60
Ala Val Pro Ala Leu Arg Leu Gly Ile Tyr Val His Ala Thr Leu
65 70 75
Glu Leu Phe Ala Leu Met Val Val Val Phe Glu Leu Cys Met Lys
80 85 90
Leu Arg Trp Leu Gly Leu His Thr Phe Ile Arg His Lys Arg Thr
95 100 105
Met Val Lys Thr Ser Val Leu Val Val Gln Phe Val Glu Ala Ile
110 115 120
Val Val Leu Val Arg Gln Met Ser His Val Arg Val Thr Arg Ala
125 130 135

```

Leu Arg Cys Ile	Phe	Leu Val Asp Cys	Arg	Tyr Cys Gly Gly	Val
	140		145		150
Arg Arg Asn Leu	Arg	Gln Ile Phe Gln	Ser	Leu Pro Pro Phe	Met
	155		160		165
Asp Ile Leu Leu	Leu	Leu Leu Phe Phe	Met	Ile Ile Phe Ala	Ile
	170		175		180
Leu Gly Phe Tyr	Leu	Phe Ser Pro Asn	Pro	Ser Asp Pro Tyr	Phe
	185		190		195
Ser Thr Leu Glu	Asn	Ser Ile Val Ser	Leu	Phe Val Leu Leu	Thr
	200		205		210
Thr Ala Asn Phe	Pro	Asp Val Met Met	Pro	Ser Tyr Ser Arg	Asn
	215		220		225
Pro Trp Ser Cys	Val	Phe Phe Ile Val	Tyr	Leu Ser Ile Glu	Leu
	230		235		240
Tyr Phe Ile Met	Asn	Leu Leu Leu Ala	Val	Val Phe Asp Thr	Phe
	245		250		255
Asn Asp Ile Glu	Lys	Arg Lys Phe Lys	Ser	Leu Leu Leu His	Lys
	260		265		270
Arg Thr Ala Ile	Gln	His Ala Tyr Arg	Leu	Leu Ile Ser Gln	Arg
	275		280		285
Arg Pro Ala Gly	Ile	Ser Tyr Arg Gln	Phe	Glu Gly Leu Met	Arg
	290		295		300
Phe Tyr Lys Pro	Arg	Met Ser Ala Arg	Glu	Arg Tyr Leu Thr	Phe
	305		310		315
Lys Ala Leu Asn	Gln	Asn Asn Thr Pro	Leu	Leu Ser Leu Lys	Asp
	320		325		330
Phe Tyr Asp Ile	Tyr	Glu Val Ala Ala	Leu	Lys Trp Lys Ala	Lys
	335		340		345
Lys Asn Arg Glu	His	Trp Phe Asp Glu	Leu	Pro Arg Thr Ala	Leu
	350		355		360
Leu Ile Phe Lys	Gly	Ile Asn Ile Leu	Val	Lys Ser Lys Ala	Phe
	365		370		375
Gln Tyr Phe Met	Tyr	Leu Val Val Ala	Val	Asn Gly Val Trp	Ile
	380		385		390
Leu Val Glu Thr	Phe	Met Leu Lys Gly	Gly	Asn Phe Phe Ser	Lys
	395		400		405
His Val Pro Trp	Ser	Tyr Leu Val Phe	Leu	Thr Ile Tyr Gly	Val
	410		415		420
Glu Leu Phe Leu	Lys	Val Ala Gly Leu	Gly	Pro Val Glu Tyr	Leu
	425		430		435
Ser Ser Gly Trp	Asn	Leu Phe Asp Phe	Ser	Val Thr Val Phe	Ala
	440		445		450
Phe Leu Gly Leu	Leu	Ala Leu Ala Leu	Asn	Met Glu Pro Phe	Tyr
	455		460		465
Phe Ile Val Val	Leu	Arg Pro Leu Gln	Leu	Leu Arg Leu Phe	Lys
	470		475		480
Leu Lys Glu Arg	Tyr	Arg Asn Val Leu	Asp	Thr Met Phe Glu	Leu
	485		490		495
Leu Pro Arg Met	Ala	Ser Leu Gly Leu	Thr	Leu Leu Ile Phe	Tyr
	500		505		510
Tyr Ser Phe Ala	Ile	Val Gly Met Glu	Phe	Phe Cys Gly Ile	Val
	515		520		525
Phe Pro Asn Cys	Cys	Asn Thr Ser Thr	Val	Ala Asp Ala Tyr	Arg
	530		535		540
Trp Arg Asn His	Thr	Val Gly Asn Arg	Thr	Val Val Glu Glu	Gly
	545		550		555
Tyr Tyr Tyr Leu	Asn	Asn Phe Asp Asn	Ile	Leu Asn Ser Phe	Val
	560		565		570
Thr Leu Phe Glu	Leu	Thr Val Val Asn	Asn	Trp Tyr Ile Ile	Met
	575		580		585
Glu Gly Val Thr	Ser	Gln Thr Ser His	Trp	Ser Arg Leu Tyr	Phe
	590		595		600
Met Thr Phe Tyr	Ile	Val Thr Met Val	Val	Met Thr Ile Ile	Val
	605		610		615
Ala Phe Ile Leu	Glu	Ala Phe Val Phe	Arg	Met Asn Tyr Ser	Arg
	620		625		630
Lys Asn Gln Asp	Ser	Glu Val Asp Gly	Gly	Ile Thr Leu Glu	Lys
	635		640		645

Glu	Ile	Ser	Lys	Glu	Glu	Leu	Val	Ala	Val	Leu	Glu	Leu	Tyr	Arg
				650					655					660
Glu	Ala	Arg	Gly	Ala	Ser	Ser	Asp	Val	Thr	Arg	Leu	Leu	Glu	Thr
				665					670					675
Leu	Ser	Gln	Met	Glu	Arg	Tyr	Gln	Gln	His	Ser	Met	Val	Phe	Leu
				680					685					690
Gly	Arg	Arg	Ser	Arg	Thr	Lys	Ser	Asp	Leu	Ser	Leu	Lys	Met	Tyr
				695					700					705
Gln	Glu	Glu	Ile	Gln	Glu	Trp	Tyr	Glu	Glu	His	Ala	Arg	Glu	Gln
				710					715					720
Glu	Gln	Gln	Arg	Gln	Leu	Ser	Ser	Ser	Ala	Ala	Pro	Ala	Ala	Gln
				725					730					735
Gln	Pro	Pro	Gly	Ser	Arg	Gln	Arg	Ser	Gln	Thr	Val	Thr		
				740					745					

&lt;210&gt; 18

&lt;211&gt; 507

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2739264CD1

&lt;400&gt; 18

Met	Ala	Phe	Asn	Phe	Gly	Ala	Pro	Ser	Gly	Thr	Ser	Gly	Thr	Ala
1				5					10					15
Ala	Ala	Thr	Ala	Ala	Pro	Ala	Gly	Gly	Phe	Gly	Gly	Phe	Gly	Thr
				20					25					30
Thr	Ser	Thr	Thr	Ala	Gly	Ser	Ala	Phe	Ser	Phe	Ser	Ala	Pro	Thr
				35					40					45
Asn	Thr	Gly	Thr	Thr	Gly	Leu	Phe	Gly	Gly	Thr	Gln	Asn	Lys	Gly
				50					55					60
Phe	Gly	Phe	Gly	Thr	Gly	Phe	Gly	Thr	Thr	Thr	Gly	Thr	Ser	Thr
				65					70					75
Gly	Leu	Gly	Thr	Gly	Leu	Gly	Thr	Gly	Leu	Gly	Phe	Gly	Gly	Phe
				80					85					90
Asn	Thr	Gln	Gln	Gln	Gln	Gln	Thr	Thr	Leu	Gly	Gly	Leu	Phe	Ser
				95					100					105
Gln	Pro	Thr	Gln	Ala	Pro	Thr	Gln	Ser	Asn	Gln	Leu	Ile	Asn	Thr
				110					115					120
Ala	Ser	Ala	Leu	Ser	Ala	Pro	Thr	Leu	Leu	Gly	Asp	Glu	Arg	Asp
				125					130					135
Ala	Ile	Leu	Ala	Lys	Trp	Asn	Gln	Leu	Gln	Ala	Phe	Trp	Gly	Thr
				140					145					150
Gly	Lys	Gly	Tyr	Phe	Asn	Asn	Asn	Ile	Pro	Pro	Val	Glu	Phe	Thr
				155					160					165
Gln	Glu	Asn	Pro	Phe	Cys	Arg	Phe	Lys	Ala	Val	Gly	Tyr	Ser	Cys
				170					175					180
Met	Pro	Ser	Asn	Lys	Asp	Glu	Asp	Gly	Leu	Val	Val	Leu	Val	Phe
				185					190					195
Asn	Lys	Lys	Glu	Thr	Glu	Ile	Arg	Ser	Gln	Gln	Gln	Gln	Leu	Val
				200					205					210
Glu	Ser	Leu	His	Lys	Val	Leu	Gly	Gly	Asn	Gln	Thr	Leu	Thr	Val
				215					220					225
Asn	Val	Glu	Gly	Thr	Lys	Thr	Leu	Pro	Asp	Asp	Gln	Thr	Glu	Val
				230					235					240
Val	Ile	Tyr	Val	Val	Glu	Arg	Ser	Pro	Asn	Gly	Thr	Ser	Arg	Arg
				245					250					255
Val	Pro	Ala	Thr	Thr	Leu	Tyr	Ala	His	Phe	Glu	Gln	Ala	Asn	Ile
				260					265					270
Lys	Thr	Gln	Leu	Gln	Gln	Leu	Gly	Val	Thr	Leu	Ser	Met	Thr	Arg
				275					280					285
Thr	Glu	Leu	Ser	Pro	Ala	Gln	Ile	Lys	Gln	Leu	Leu	Gln	Asn	Pro
				290					295					300
Pro	Ala	Gly	Val	Asp	Pro	Ile	Ile	Trp	Glu	Gln	Ala	Lys	Val	Asp
				305					310					315
Asn	Pro	Asp	Ser	Glu	Lys	Leu	Ile	Pro	Val	Pro	Met	Val	Gly	Phe
				320					325					330

```

Lys Glu Leu Leu Arg Arg Leu Lys Val Gln Asp Gln Met Thr Lys
335 340 345
Gln His Gln Thr Arg Leu Asp Ile Ile Ser Glu Asp Ile Ser Glu
350 355 360
Leu Gln Lys Asn Gln Thr Thr Ser Val Ala Lys Ile Ala Gln Tyr
365 370 375
Lys Arg Lys Leu Met Asp Leu Ser His Arg Thr Leu Gln Val Leu
380 385 390
Ile Lys Gln Glu Ile Gln Arg Lys Ser Gly Tyr Ala Ile Gln Ala
395 400 405
Asp Glu Glu Gln Leu Arg Val Gln Leu Asp Thr Ile Gln Gly Glu
410 415 420
Leu Asn Ala Pro Thr Gln Phe Lys Gly Arg Leu Asn Glu Leu Met
425 430 435
Ser Gln Ile Arg Met Gln Asn His Phe Gly Ala Val Arg Ser Glu
440 445 450
Glu Arg Tyr Tyr Ile Asp Ala Asp Leu Leu Arg Glu Ile Lys Gln
455 460 465
His Leu Lys Gln Gln Gln Glu Gly Leu Ser His Leu Ile Ser Ile
470 475 480
Ile Lys Asp Asp Leu Glu Asp Ile Lys Leu Val Glu His Gly Leu
485 490 495
Asn Glu Thr Ile His Ile Arg Gly Gly Val Phe Ser
500 505

```

&lt;210&gt; 19

&lt;211&gt; 592

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2758310CD1

&lt;400&gt; 19

```

Met Trp Phe Cys Gly Gln Ser Thr Pro Phe Gly Cys Glu Leu His
1 5 10 15
Asp Thr Cys Val Gln Leu Cys His Phe His Ser Ala Leu Leu His
20 25 30
Arg Arg Gln Lys Pro Trp Pro Ser Pro Ala Val Phe Phe Arg Arg
35 40 45
Asn Val Arg Gly Leu Pro Pro Arg Phe Ser Ser Pro Thr Pro Leu
50 55 60
Trp Arg Lys Val Leu Ser Thr Ala Val Val Gly Ala Pro Leu Leu
65 70 75
Leu Gly Ala Arg Tyr Val Met Ala Glu Ala Arg Glu Lys Arg Arg
80 85 90
Met Arg Leu Val Val Asp Gly Met Gly Arg Phe Gly Arg Ser Leu
95 100 105
Lys Val Gly Leu Gln Ile Ser Leu Asp Tyr Trp Trp Cys Thr Asn
110 115 120
Val Val Leu Arg Gly Trp Lys Ser Pro Gly Tyr Leu Glu Val Met
125 130 135
Ser Ala Cys His Gln Arg Ala Ala Asp Ala Leu Val Ala Gly Ala
140 145 150
Ile Ser Asn Gly Gly Leu Tyr Val Lys Leu Gly Gln Gly Leu Cys
155 160 165
Ser Phe Asn His Leu Leu Pro Pro Glu Tyr Thr Arg Thr Leu Arg
170 175 180
Val Leu Glu Asp Arg Ala Leu Lys Arg Gly Phe Gln Glu Val Asp
185 190 195
Glu Leu Phe Leu Glu Asp Phe Gln Ala Leu Pro His Glu Leu Phe
200 205 210
Gln Glu Phe Asp Tyr Gln Pro Ile Ala Ala Ala Ser Leu Ala Gln
215 220 225
Val His Arg Ala Lys Leu His Asp Gly Thr Ser Val Ala Val Lys
230 235 240
Val Gln Tyr Ile Asp Leu Arg Asp Arg Phe Asp Gly Asp Ile His
245 250 255

```

Thr	Leu	Glu	Leu	Leu	Leu	Arg	Leu	Val	Glu	Val	Met	His	Pro	Ser	
				260					265						270
Phe	Gly	Phe	Ser	Trp	Val	Leu	Gln	Asp	Leu	Lys	Gly	Thr	Leu	Ala	
				275					280						285
Gln	Glu	Leu	Asp	Phe	Glu	Asn	Glu	Gly	Arg	Asn	Ala	Glu	Arg	Cys	
				290					295						300
Ala	Arg	Glu	Leu	Ala	His	Phe	Pro	Tyr	Val	Val	Val	Pro	Arg	Val	
				305					310						315
His	Trp	Asp	Lys	Ser	Ser	Lys	Arg	Val	Leu	Thr	Ala	Asp	Phe	Cys	
				320					325						330
Ala	Gly	Cys	Lys	Val	Asn	Asp	Val	Glu	Ala	Ile	Arg	Ser	Gln	Gly	
				335					340						345
Leu	Ala	Val	His	Asp	Ile	Ala	Glu	Lys	Leu	Ile	Lys	Ala	Phe	Ala	
				350					355						360
Glu	Gln	Ile	Phe	Tyr	Thr	Gly	Phe	Ile	His	Ser	Asp	Pro	His	Pro	
				365					370						375
Gly	Asn	Val	Leu	Val	Arg	Lys	Gly	Pro	Asp	Gly	Lys	Ala	Glu	Leu	
				380					385						390
Val	Leu	Leu	Asp	His	Gly	Leu	Tyr	Gln	Phe	Leu	Glu	Glu	Lys	Asp	
				395					400						405
Arg	Ala	Ala	Leu	Cys	Gln	Leu	Trp	Arg	Ala	Ile	Ile	Leu	Arg	Asp	
				410					415						420
Asp	Ala	Ala	Met	Arg	Ala	His	Ala	Ala	Ala	Leu	Gly	Val	Gln	Asp	
				425					430						435
Tyr	Leu	Leu	Phe	Ala	Glu	Met	Leu	Met	Gln	Arg	Pro	Val	Arg	Leu	
				440					445						450
Gly	Gln	Leu	Trp	Gly	Ser	His	Leu	Leu	Ser	Arg	Glu	Glu	Ala	Ala	
				455					460						465
Tyr	Met	Val	Asp	Met	Ala	Arg	Glu	Arg	Phe	Glu	Ala	Val	Met	Ala	
				470					475						480
Val	Leu	Arg	Glu	Leu	Pro	Arg	Pro	Met	Leu	Leu	Val	Leu	Arg	Asn	
				485					490						495
Ile	Asn	Thr	Val	Arg	Ala	Ile	Asn	Val	Ala	Leu	Gly	Ala	Pro	Val	
				500					505						510
Asp	Arg	Tyr	Phe	Leu	Met	Ala	Lys	Arg	Ala	Val	Arg	Gly	Trp	Ser	
				515					520						525
Arg	Leu	Ala	Gly	Ala	Thr	Tyr	Arg	Gly	Val	Tyr	Gly	Thr	Ser	Leu	
				530					535						540
Leu	Arg	His	Ala	Lys	Val	Val	Trp	Glu	Met	Leu	Lys	Phe	Glu	Val	
				545					550						555
Ala	Leu	Arg	Leu	Glu	Thr	Leu	Ala	Met	Arg	Leu	Thr	Ala	Leu	Leu	
				560					565						570
Ala	Arg	Ala	Leu	Val	His	Leu	Ser	Leu	Val	Pro	Pro	Ala	Glu	Glu	
				575					580						585
Leu	Tyr	Gln	Tyr	Leu	Glu	Thr									590

&lt;210&gt; 20

&lt;211&gt; 841

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2762348CD1

&lt;400&gt; 20

Met	Ala	Ser	Val	Phe	Arg	Ser	Glu	Glu	Met	Cys	Leu	Ser	Gln	Leu	
1				5					10					15	
Phe	Leu	Gln	Val	Glu	Ala	Ala	Tyr	Cys	Cys	Val	Ala	Glu	Leu	Gly	
				20					25					30	
Glu	Leu	Gly	Leu	Val	Gln	Phe	Lys	Asp	Leu	Asn	Met	Asn	Val	Asn	
				35					40					45	
Ser	Phe	Gln	Arg	Lys	Phe	Val	Asn	Glu	Val	Arg	Arg	Cys	Glu	Ser	
				50					55					60	
Leu	Glu	Arg	Ile	Leu	Arg	Phe	Leu	Glu	Asp	Glu	Met	Gln	Asn	Glu	
				65					70					75	
Ile	Val	Val	Gln	Leu	Leu	Glu	Lys	Ser	Pro	Leu	Thr	Pro	Leu	Pro	
				80					85					90	

Arg	Glu	Met	Ile	Thr	Leu	Glu	Thr	Val	Leu	Glu	Lys	Leu	Glu	Gly	
				95					100					105	
Glu	Leu	Gln	Glu	Ala	Asn	Gln	Asn	Gln	Gln	Ala	Leu	Lys	Gln	Ser	
				110					115					120	
Phe	Leu	Glu	Leu	Thr	Glu	Leu	Lys	Tyr	Leu	Leu	Lys	Lys	Thr	Gln	
				125					130					135	
Asp	Phe	Phe	Glu	Thr	Glu	Thr	Asn	Leu	Ala	Asp	Asp	Phe	Phe	Thr	
				140					145					150	
Glu	Asp	Thr	Ser	Gly	Leu	Leu	Glu	Leu	Lys	Ala	Val	Pro	Ala	Tyr	
				155					160					165	
Met	Thr	Gly	Lys	Leu	Gly	Phe	Ile	Ala	Gly	Cys	Asp	Pro	Thr	Gly	
				170					175					180	
Lys	Arg	Met	Ala	Ser	Phe	Glu	Arg	Leu	Leu	Trp	Arg	Val	Cys	Arg	
				185					190					195	
Gly	Asn	Val	Tyr	Leu	Lys	Phe	Ser	Glu	Met	Asp	Ala	Pro	Leu	Glu	
				200					205					210	
Asp	Pro	Val	Thr	Lys	Glu	Glu	Ile	Gln	Lys	His	Ile	Phe	Ile	Ile	
				215					220					225	
Phe	Tyr	Gln	Gly	Glu	Gln	Leu	Arg	Gln	Lys	Ile	Lys	Lys	Ile	Cys	
				230					235					240	
Asp	Gly	Phe	Arg	Ala	Thr	Val	Tyr	Pro	Cys	Pro	Glu	Pro	Ala	Val	
				245					250					255	
Glu	Arg	Arg	Glu	Met	Leu	Glu	Ser	Val	Asn	Val	Arg	Leu	Glu	Asp	
				260					265					270	
Leu	Ile	Thr	Val	Ile	Thr	Gln	Thr	Glu	Ser	His	Arg	Gln	Arg	Leu	
				275					280					285	
Leu	Gln	Glu	Ala	Ala	Ala	Asn	Trp	His	Ser	Trp	Leu	Ile	Lys	Val	
				290					295					300	
Gln	Lys	Met	Lys	Ala	Val	Tyr	His	Ile	Leu	Asn	Met	Cys	Asn	Ile	
				305					310					315	
Asp	Val	Thr	Gln	Gln	Cys	Val	Ile	Ala	Glu	Ile	Trp	Phe	Pro	Val	
				320					325					330	
Ala	Asp	Ala	Thr	Arg	Ile	Lys	Arg	Ala	Leu	Glu	Gln	Gly	Met	Glu	
				335					340					345	
Leu	Ser	Gly	Ser	Ser	Met	Ala	Pro	Ile	Met	Thr	Thr	Val	Gln	Ser	
				350					355					360	
Lys	Thr	Ala	Pro	Pro	Thr	Phe	Asn	Arg	Thr	Asn	Lys	Phe	Thr	Ala	
				365					370					375	
Gly	Phe	Gln	Asn	Ile	Val	Asp	Ala	Tyr	Gly	Val	Gly	Ser	Tyr	Arg	
				380					385					390	
Glu	Ile	Asn	Pro	Ala	Pro	Tyr	Thr	Ile	Ile	Thr	Phe	Pro	Phe	Leu	
				395					400					405	
Phe	Ala	Val	Met	Phe	Gly	Asp	Cys	Gly	His	Gly	Thr	Val	Met	Leu	
				410					415					420	
Leu	Ala	Ala	Leu	Trp	Met	Ile	Leu	Asn	Glu	Arg	Arg	Leu	Leu	Ser	
				425					430					435	
Gln	Lys	Thr	Asp	Asn	Glu	Ile	Trp	Asn	Thr	Phe	Phe	His	Gly	Arg	
				440					445					450	
Tyr	Leu	Ile	Leu	Leu	Met	Gly	Ile	Phe	Ser	Ile	Tyr	Thr	Gly	Leu	
				455					460					465	
Ile	Tyr	Asn	Asp	Cys	Phe	Ser	Lys	Ser	Leu	Asn	Ile	Phe	Gly	Ser	
				470					475					480	
Ser	Trp	Ser	Val	Gln	Pro	Met	Phe	Arg	Asn	Gly	Thr	Trp	Asn	Thr	
				485					490					495	
His	Val	Met	Glu	Glu	Ser	Leu	Tyr	Leu	Gln	Leu	Asp	Pro	Ala	Ile	
				500					505					510	
Pro	Gly	Val	Tyr	Phe	Gly	Asn	Pro	Tyr	Pro	Phe	Gly	Ile	Asp	Pro	
				515					520					525	
Ile	Trp	Asn	Leu	Ala	Ser	Asn	Lys	Leu	Thr	Phe	Leu	Asn	Ser	Tyr	
				530					535					540	
Lys	Met	Lys	Met	Ser	Val	Ile	Leu	Gly	Ile	Val	Gln	Met	Val	Phe	
				545					550					555	
Gly	Val	Ile	Leu	Ser	Leu	Phe	Asn	His	Ile	Tyr	Phe	Arg	Arg	Thr	
				560					565					570	
Leu	Asn	Ile	Ile	Leu	Gln	Phe	Ile	Pro	Glu	Met	Ile	Phe	Ile	Leu	
				575					580					585	
Cys	Leu	Phe	Gly	Tyr	Leu	Val	Phe	Met	Ile	Ile	Phe	Lys	Trp	Cys	
				590					595					600	

Cys	Phe	Asp	Val	His	Val	Ser	Gln	His	Ala	Pro	Ser	Ile	Leu	Ile	
					605				610						615
His	Phe	Ile	Asn	Met	Phe	Leu	Phe	Asn	Tyr	Ser	Asp	Ser	Ser	Asn	
					620				625						630
Ala	Pro	Leu	Tyr	Lys	His	Gln	Gln	Glu	Val	Gln	Ser	Phe	Phe	Val	
					635				640						645
Val	Met	Ala	Leu	Ile	Ser	Val	Pro	Trp	Met	Leu	Leu	Ile	Lys	Pro	
					650				655						660
Phe	Ile	Leu	Arg	Ala	Ser	His	Arg	Lys	Ser	Gln	Leu	Gln	Ala	Ser	
					665				670						675
Arg	Ile	Gln	Glu	Asp	Ala	Thr	Glu	Asn	Ile	Glu	Gly	Asp	Ser	Ser	
					680				685						690
Ser	Pro	Ser	Ser	Arg	Ser	Gly	Gln	Arg	Thr	Ser	Ala	Asp	Thr	His	
					695				700						705
Gly	Ala	Leu	Asp	Asp	His	Gly	Glu	Glu	Phe	Asn	Phe	Gly	Asp	Val	
					710				715						720
Phe	Val	His	Gln	Ala	Ile	His	Thr	Ile	Glu	Tyr	Cys	Leu	Gly	Cys	
					725				730						735
Ile	Ser	Asn	Thr	Ala	Ser	Tyr	Leu	Arg	Leu	Trp	Ala	Leu	Ser	Leu	
					740				745						750
Ala	His	Ala	Gln	Leu	Ser	Glu	Val	Leu	Trp	Thr	Met	Val	Met	Asn	
					755				760						765
Ser	Gly	Leu	Gln	Thr	Arg	Gly	Trp	Gly	Gly	Ile	Val	Gly	Val	Phe	
					770				775						780
Ile	Ile	Phe	Ala	Val	Phe	Ala	Val	Leu	Thr	Val	Ala	Ile	Leu	Leu	
					785				790						795
Ile	Met	Glu	Gly	Leu	Ser	Ala	Phe	Leu	His	Ala	Leu	Arg	Leu	His	
					800				805						810
Trp	Val	Glu	Phe	Gln	Asn	Lys	Phe	Tyr	Val	Gly	Asp	Gly	Tyr	Lys	
					815				820						825
Phe	Ser	Pro	Phe	Ser	Phe	Lys	His	Ile	Leu	Asp	Gly	Thr	Ala	Glu	
					830				835						840

Glu

&lt;210&gt; 21

&lt;211&gt; 253

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3715961CD1

&lt;400&gt; 21

Met	Ser	Glu	Cys	Pro	Leu	Ile	Leu	Tyr	Ile	His	Lys	His	Ile	Asp	
1				5					10					15	
Thr	Tyr	Ser	Gln	Ser	Tyr	Leu	Phe	Asn	Asp	Leu	Phe	Tyr	Pro	Val	
				20					25					30	
Tyr	Ser	Gly	Gly	Arg	Met	Val	Thr	Tyr	Glu	His	Leu	Arg	Glu	Val	
				35					40					45	
Val	Phe	Gly	Lys	Ser	Glu	Asp	Glu	His	Tyr	Pro	Leu	Trp	Lys	Ser	
				50					55					60	
Val	Ile	Gly	Gly	Met	Met	Ala	Gly	Val	Ile	Gly	Gln	Phe	Leu	Ala	
				65					70					75	
Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Gln	Met	Gln	Met	Glu	Gly	Lys	
				80					85					90	
Arg	Lys	Leu	Glu	Gly	Lys	Pro	Leu	Arg	Phe	Arg	Gly	Val	His	His	
				95					100					105	
Ala	Phe	Ala	Lys	Ile	Leu	Ala	Glu	Gly	Gly	Ile	Arg	Gly	Leu	Trp	
				110					115					120	
Ala	Gly	Trp	Val	Pro	Asn	Ile	Gln	Arg	Ala	Ala	Leu	Val	Asn	Met	
				125					130					135	
Gly	Asp	Leu	Thr	Thr	Tyr	Asp	Thr	Val	Lys	His	Tyr	Leu	Val	Leu	
				140					145					150	
Asn	Thr	Pro	Leu	Glu	Asp	Asn	Ile	Met	Thr	His	Gly	Leu	Ser	Ser	
				155					160					165	
Leu	Cys	Ser	Gly	Leu	Val	Ala	Ser	Ile	Leu	Gly	Thr	Pro	Ala	Asp	
				170					175					180	



Val	Ile	Lys	Ser	Arg	Ile	Met	Asn	Gln	Pro	Arg	Asp	Lys	Gln	Gly	
				185					190					195	
Arg	Gly	Leu	Leu	Tyr	Lys	Ser	Ser	Thr	Asp	Cys	Leu	Ile	Gln	Ala	
				200					205					210	
Val	Gln	Gly	Glu	Gly	Phe	Met	Ser	Leu	Tyr	Lys	Gly	Phe	Leu	Pro	
				215					220					225	
Ser	Trp	Leu	Arg	Met	Thr	Pro	Trp	Ser	Met	Val	Phe	Trp	Leu	Thr	
				230					235					240	
Tyr	Glu	Lys	Ile	Arg	Glu	Met	Ser	Gly	Val	Ser	Pro	Phe			
				245					250						

&lt;210&gt; 22

&lt;211&gt; 229

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5108194CD1

&lt;400&gt; 22

Met	Gly	Asn	Gly	Val	Lys	Glu	Gly	Pro	Val	Arg	Leu	His	Glu	Asp	
1				5					10					15	
Ala	Glu	Ala	Val	Leu	Ser	Ser	Ser	Val	Ser	Ser	Lys	Arg	Asp	His	
				20					25					30	
Arg	Gln	Val	Leu	Ser	Ser	Leu	Leu	Ser	Gly	Ala	Leu	Ala	Gly	Ala	
				35					40					45	
Leu	Ala	Lys	Thr	Ala	Val	Ala	Pro	Leu	Asp	Arg	Thr	Lys	Ile	Ile	
				50					55					60	
Phe	Gln	Val	Ser	Ser	Lys	Arg	Phe	Ser	Ala	Lys	Glu	Ala	Phe	Arg	
				65					70					75	
Val	Leu	Tyr	Tyr	Thr	Tyr	Leu	Asn	Glu	Gly	Phe	Leu	Ser	Leu	Trp	
				80					85					90	
Arg	Gly	Asn	Ser	Ala	Thr	Met	Val	Arg	Val	Val	Pro	Tyr	Ala	Ala	
				95					100					105	
Ile	Gln	Phe	Ser	Ala	His	Glu	Glu	Tyr	Lys	Arg	Ile	Leu	Gly	Ser	
				110					115					120	
Tyr	Tyr	Gly	Phe	Arg	Gly	Glu	Ala	Leu	Pro	Pro	Trp	Pro	Arg	Leu	
				125					130					135	
Phe	Ala	Gly	Ala	Leu	Ala	Gly	Thr	Thr	Ala	Ala	Ser	Leu	Thr	Tyr	
				140					145					150	
Pro	Leu	Asp	Leu	Val	Arg	Ala	Arg	Met	Ala	Val	Thr	Pro	Lys	Glu	
				155					160					165	
Met	Tyr	Ser	Asn	Ile	Phe	His	Val	Phe	Ile	Arg	Ile	Ser	Arg	Glu	
				170					175					180	
Glu	Gly	Leu	Lys	Thr	Leu	Tyr	His	Gly	Phe	Met	Pro	Thr	Val	Leu	
				185					190					195	
Gly	Val	Ile	Pro	Tyr	Ala	Gly	Leu	Ser	Phe	Phe	Thr	Tyr	Glu	Thr	
				200					205					210	
Leu	Lys	Ser	Leu	His	Arg	Glu	Tyr	Ser	Gly	Arg	Lys	Leu	Ile	Pro	
				215					220					225	

Phe Ser Glu Gly

&lt;210&gt; 23

&lt;211&gt; 170

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5503122CD1

&lt;400&gt; 23

Met	Tyr	Asp	Asn	Leu	Tyr	Leu	His	Gly	Ile	Glu	Asp	Ser	Glu	Ala	
1				5					10					15	
Gly	Ser	Ala	Asp	Ser	Tyr	Thr	Ser	Arg	Pro	Ser	Asp	Ser	Asp	Val	
				20					25					30	
Ser	Leu	Glu	Glu	Asp	Arg	Glu	Ala	Ile	Arg	Gln	Glu	Arg	Glu	Gln	
				35					40					45	

```
<210> 24
<211> 655
<212> PRT
<213> Homo sapiens
```

<400> 24

21/60

```

Asp Ser Thr Ala Val Ala Leu Asn Arg Glu Glu Asp Phe Lys Ala
305 310 315
Thr Glu Ile Ile Glu Pro Ser Lys Gln Asp Lys Pro Leu Ile Glu
320 325 330
Lys Leu Ala Glu Ile Tyr Val Asn Ser Ser Phe Tyr Lys Glu Thr
335 340 345
Lys Ala Glu Leu His Gln Leu Ser Gly Gly Glu Lys Lys Lys Lys
350 355 360
Ile Thr Val Phe Lys Glu Ile Ser Tyr Thr Thr Ser Phe Cys His
365 370 375
Gln Leu Arg Trp Val Ser Lys Arg Ser Phe Lys Asn Leu Leu Gly
380 385 390
Asn Pro Gln Ala Ser Ile Ala Gln Ile Ile Val Thr Val Val Leu
395 400 405
Gly Leu Val Ile Gly Ala Ile Tyr Phe Gly Leu Lys Asn Asp Ser
410 415 420
Thr Gly Ile Gln Asn Arg Ala Gly Val Leu Phe Phe Leu Thr Thr
425 430 435
Asn Gln Cys Phe Ser Ser Val Ser Ala Val Glu Leu Phe Val Val
440 445 450
Glu Lys Lys Leu Phe Ile His Glu Tyr Ile Ser Gly Tyr Tyr Arg
455 460 465
Val Ser Ser Tyr Phe Leu Gly Lys Leu Leu Ser Asp Leu Leu Pro
470 475 480
Met Arg Met Leu Pro Ser Ile Ile Phe Thr Cys Ile Val Tyr Phe
485 490 495
Met Leu Gly Leu Lys Pro Lys Ala Asp Ala Phe Phe Val Met Met
500 505 510
Phe Thr Leu Met Met Val Ala Tyr Ser Ala Ser Ser Met Ala Leu
515 520 525
Ala Ile Ala Ala Gly Gln Ser Val Val Ser Val Ala Thr Leu Leu
530 535 540
Met Thr Ile Cys Phe Val Phe Met Met Ile Phe Ser Gly Leu Leu
545 550 555
Val Asn Leu Thr Thr Ile Ala Ser Trp Leu Ser Trp Leu Gln Tyr
560 565 570
Phe Ser Ile Pro Arg Tyr Gly Phe Thr Ala Leu Gln His Asn Glu
575 580 585
Phe Leu Gly Gln Asn Phe Cys Pro Gly Leu Asn Ala Thr Gly Asn
590 595 600
Asn Pro Cys Asn Tyr Ala Thr Cys Thr Gly Glu Glu Tyr Leu Val
605 610 615
Lys Gln Gly Ile Asp Leu Ser Pro Trp Gly Leu Trp Lys Asn His
620 625 630
Val Ala Leu Ala Cys Met Ile Val Ile Phe Leu Thr Ile Ala Tyr
635 640 645
Leu Lys Leu Leu Phe Leu Lys Lys Tyr Ser
650 655

```

&lt;210&gt; 25

&lt;211&gt; 184

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5593114CD1

&lt;400&gt; 25

```

Met Trp Val Phe Gly Tyr Gly Ser Leu Ile Trp Lys Val Asp Phe
1 5 10 15
Pro Tyr Gln Asp Lys Leu Val Gly Tyr Ile Thr Asn Tyr Ser Arg
20 25 30
Arg Phe Trp Gln Gly Ser Thr Asp His Arg Gly Val Pro Gly Lys
35 40 45
Pro Gly Arg Val Val Thr Leu Val Glu Asp Pro Ala Gly Cys Val
50 55 60
Trp Gly Val Ala Tyr Arg Leu Pro Val Gly Lys Glu Glu Glu Val
65 70 75

```

Lys Ala Tyr Leu Asp Phe Arg Glu Lys Gly Gly Tyr Arg Thr Thr  
 80 85 90  
 Thr Val Ile Phe Tyr Pro Lys Asp Pro Thr Thr Lys Pro Phe Ser  
 95 100 105  
 Val Leu Leu Tyr Ile Gly Thr Cys Asp Asn Pro Asp Tyr Leu Gly  
 110 115 120  
 Pro Ala Pro Leu Glu Asp Ile Ala Glu Gln Ile Phe Asn Ala Ala  
 125 130 135  
 Gly Pro Ser Gly Arg Asn Thr Glu Tyr Leu Phe Glu Leu Ala Asn  
 140 145 150  
 Ser Ile Arg Asn Leu Val Pro Glu Glu Ala Asp Glu His Leu Phe  
 155 160 165  
 Ala Leu Glu Lys Leu Val Lys Glu Arg Leu Glu Gly Lys Gln Asn  
 170 175 180  
 Leu Asn Cys Ile

<210> 26  
 <211> 154  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 044775CD1

<400> 26  
 Met Gly Ala Phe Glu Cys Val Arg Lys Val Tyr Gln Thr Asp Gly  
 1 5 10 15  
 Leu Lys Gly Phe Tyr Arg Gly Met Ser Ala Ser Tyr Ala Gly Ile  
 20 25 30  
 Ser Glu Thr Val Ile His Phe Val Ile Tyr Glu Ser Ile Lys Gln  
 35 40 45  
 Lys Leu Leu Glu Tyr Lys Thr Ala Ser Thr Met Glu Asn Asp Glu  
 50 55 60  
 Glu Ser Val Lys Glu Ala Ser Asp Phe Val Gly Met Met Leu Ala  
 65 70 75  
 Ala Ala Thr Ser Lys Thr Cys Ala Thr Thr Ile Ala Tyr Pro His  
 80 85 90  
 Glu Val Val Arg Thr Arg Leu Arg Glu Glu Gly Thr Lys Tyr Arg  
 95 100 105  
 Ser Phe Phe Gln Thr Leu Ser Leu Leu Val Gln Glu Glu Gly Tyr  
 110 115 120  
 Gly Ser Leu Tyr Arg Gly Leu Thr Thr His Leu Val Arg Gln Ile  
 125 130 135  
 Pro Asn Thr Ala Ile Met Met Ala Thr Tyr Glu Leu Val Val Tyr  
 140 145 150  
 Leu Leu Asn Gly

<210> 27  
 <211> 438  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 116588CD1

<400> 27  
 Met Leu Leu Val Thr Pro Arg Pro Glu Arg Gly Gly Arg Gly Thr  
 1 5 10 15  
 Glu Leu Gly Glu Phe Cys Gly Thr Pro Leu Leu Phe Ser Ser Tyr  
 20 25 30  
 Phe Cys Tyr Asp Asn Pro Ala Ala Leu Gln Thr Gln Val Lys Arg  
 35 40 45  
 Asp Met Gln Val Asn Thr Thr Lys Phe Met Leu Leu Tyr Ala Trp  
 50 55 60  
 Tyr Ser Trp Pro Asn Val Val Leu Cys Phe Phe Gly Gly Phe Leu  
 65 70 75

```

Ile Asp Arg Val Phe Gly Ile Arg Trp Gly Thr Ile Ile Phe Ser
80 85 90
Cys Phe Val Cys Ile Gly Gln Val Val Phe Ala Leu Gly Gly Ile
95 100 105
Phe Asn Ala Phe Trp Leu Met Glu Phe Gly Arg Phe Val Phe Gly
110 115 120
Ile Gly Gly Glu Ser Leu Ala Val Ala Gln Asn Thr Tyr Ala Val
125 130 135
Ser Trp Phe Lys Gly Lys Glu Leu Asn Leu Val Phe Gly Leu Gln
140 145 150
Leu Ser Met Ala Arg Ile Gly Ser Thr Val Asn Met Asn Leu Met
155 160 165
Gly Trp Leu Tyr Ser Lys Ile Glu Ala Leu Leu Gly Ser Ala Gly
170 175 180
His Thr Thr Leu Gly Ile Thr Leu Met Ile Gly Gly Val Thr Cys
185 190 195
Ile Leu Ser Leu Ile Cys Ala Leu Ala Leu Ala Tyr Leu Asp Gln
200 205 210
Arg Ala Glu Arg Ile Leu His Lys Glu Gln Gly Lys Thr Gly Glu
215 220 225
Val Ile Lys Leu Thr Asp Val Lys Asp Phe Ser Leu Pro Leu Trp
230 235 240
Leu Ile Phe Ile Ile Cys Val Cys Tyr Tyr Val Ala Val Phe Pro
245 250 255
Phe Ile Gly Leu Gly Lys Val Phe Phe Thr Glu Lys Phe Gly Phe
260 265 270
Ser Ser Gln Ala Ala Ser Ala Ile Asn Ser Val Val Tyr Val Ile
275 280 285
Ser Ala Pro Met Ser Pro Val Phe Gly Leu Leu Val Asp Lys Thr
290 295 300
Gly Lys Asn Ile Ile Trp Val Leu Cys Ala Val Ala Ala Thr Leu
305 310 315
Val Ser His Met Met Leu Ala Phe Thr Met Trp Asn Pro Trp Ile
320 325 330
Ala Met Cys Leu Leu Gly Leu Ser Tyr Ser Leu Leu Ala Cys Ala
335 340 345
Leu Trp Pro Met Val Ala Phe Val Val Pro Glu His Gln Leu Gly
350 355 360
Thr Ala Tyr Gly Phe Met Gln Ser Ile Gln Asn Leu Gly Leu Ala
365 370 375
Ile Ile Ser Ile Ile Ala Gly Met Ile Leu Asp Ser Arg Gly Tyr
380 385 390
Leu Phe Leu Glu Val Phe Phe Ile Ala Cys Val Ser Leu Ser Leu
395 400 405
Leu Ser Val Val Leu Leu Tyr Leu Val Asn Arg Ala Gln Gly Gly
410 415 420
Asn Leu Asn Tyr Ser Ala Arg Gln Arg Glu Glu Ile Lys Phe Ser
425 430 435
His Thr Glu

```

&lt;210&gt; 28

&lt;211&gt; 237

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 875369CD1

&lt;400&gt; 28

```

Met Ala His Val Gly Ser Arg Lys Arg Ser Arg Ser Arg Ser Arg
1 5 10 15
Ser Arg Gly Arg Gly Ser Glu Lys Arg Lys Lys Lys Ser Arg Lys
20 25 30
Asp Thr Ser Arg Asn Cys Ser Ala Ser Thr Ser Gln Gly Arg Lys
35 40 45
Ala Ser Thr Ala Pro Gly Ala Glu Ala Ser Pro Ser Pro Cys Ile
50 55 60

```

```

Thr Glu Arg Ser Lys Gln Lys Ala Arg Arg Arg Thr Arg Ser Ser
65 70 75
Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
80 85 90
Ser Ser Ser Ser Ser Ser Ser Ser Asp Gly Arg Lys Lys Arg Gly
95 100 105
Lys Tyr Lys Asp Lys Arg Arg Lys Lys Lys Lys Lys Arg Lys Lys
110 115 120
Leu Lys Lys Lys Gly Lys Glu Lys Ala Glu Ala Gln Gln Val Glu
125 130 135
Ala Leu Pro Gly Pro Ser Leu Asp Gln Trp His Arg Ser Ala Gly
140 145 150
Glu Glu Glu Asp Gly Pro Val Leu Thr Asp Glu Gln Lys Ser Arg
155 160 165
Ile Gln Ala Met Lys Pro Met Thr Lys Glu Glu Trp Asp Ala Arg
170 175 180
Gln Ser Ile Ile Arg Lys Val Val Asp Pro Glu Thr Gly Arg Thr
185 190 195
Arg Leu Ile Lys Gly Asp Gly Glu Val Leu Glu Glu Ile Val Thr
200 205 210
Lys Glu Arg His Arg Glu Ile Asn Lys Gln Ala Thr Arg Gly Asp
215 220 225
Cys Leu Ala Phe Gln Met Arg Ala Gly Leu Leu Pro
230 235

```

&lt;210&gt; 29

&lt;211&gt; 219

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1325518CD1

&lt;400&gt; 29

```

Met Lys Leu Leu Leu Trp Ala Cys Ile Val Cys Val Ala Phe Ala
1 5 10 15
Arg Lys Arg Arg Phe Pro Phe Ile Gly Glu Asp Asp Asn Asp Asp
20 25 30
Gly His Pro Leu His Pro Ser Leu Asn Ile Pro Tyr Gly Ile Arg
35 40 45
Asn Leu Pro Pro Pro Leu Tyr Tyr Arg Pro Val Asn Thr Val Pro
50 55 60
Ser Tyr Pro Gly Asn Thr Tyr Thr Asp Thr Gly Leu Pro Ser Tyr
65 70 75
Pro Trp Ile Leu Thr Ser Pro Gly Phe Pro Tyr Val Tyr His Ile
80 85 90
Arg Gly Phe Pro Leu Ala Thr Gln Leu Asn Val Pro Pro Leu Pro
95 100 105
Pro Arg Gly Phe Pro Phe Val Pro Pro Ser Arg Phe Phe Ser Ala
110 115 120
Ala Ala Ala Pro Ala Ala Pro Pro Ile Ala Ala Glu Pro Ala Ala
125 130 135
Ala Ala Pro Leu Thr Ala Thr Pro Val Ala Ala Glu Pro Ala Ala
140 145 150
Gly Ala Pro Val Ala Ala Glu Pro Ala Ala Glu Ala Pro Val Gly
155 160 165
Ala Glu Pro Ala Ala Glu Ala Pro Val Ala Ala Glu Pro Ala Ala
170 175 180
Glu Ala Pro Val Gly Val Glu Pro Ala Ala Glu Glu Pro Ser Pro
185 190 195
Ala Glu Pro Ala Thr Ala Lys Pro Ala Ala Pro Glu Pro His Pro
200 205 210
Ser Pro Ser Leu Glu Gln Ala Asn Gln
215

```

&lt;210&gt; 30

&lt;211&gt; 707

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

<400> 30

26/60

Ala Leu Glu Ser	470	Ala Asp Gly His Gly	475	Gly Cys Ser Ala Gly	480
	485		490		495
Asp Ile Gly Pro	500	Tyr Gly Glu Arg Asp	505	Asp Gln Gln Val Phe	510
Gln Lys Val Val	515	Pro Ser Ala Ser Gln	520	Leu Phe Val Arg Leu	525
Ser Thr Gly Gln	530	Arg Val Cys Ser Val	535	Arg Ser Val Asp Gly	540
Pro Thr Thr Ala	545	Phe Thr Val Leu Glu	550	Cys Glu Gly Ser Arg	555
Leu Gly Ser Arg	560	Pro Arg Arg Tyr Leu	565	Leu Thr Gly Gln Ala	570
Gly Ser Leu Ala	575	Met Trp Asp Leu Thr	580	Thr Ala Met Asp Gly	585
Gly Gln Ala Pro	590	Ala Gly Gly Leu Thr	595	Glu Gln Glu Leu Met	600
Gln Leu Glu His	605	Cys Glu Leu Ala Pro	610	Pro Ala Pro Ser Ala	615
Ser Trp Gly Cys	620	Leu Pro Ser Pro Ser	625	Pro Arg Ile Ser Leu	630
Ser Leu His Ser	635	Ala Ser Ser Asn Thr	640	Ser Leu Ser Gly His	645
Gly Ser Pro Ser	650	Pro Pro Gln Ala Glu	655	Ala Arg Arg Arg Gly	660
Gly Ser Phe Val	665	Glu Arg Cys Gln Glu	670	Leu Val Arg Ser Gly	675
Asp Leu Arg Arg	680	Pro Pro Thr Pro Ala	685	Pro Trp Pro Ser Ser	690
Leu Gly Thr Pro	695	Leu Thr Pro Pro Lys	700	Met Lys Leu Asn Glu	705
Ser Phe					

&lt;210&gt; 31

&lt;211&gt; 279

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2172064CD1

&lt;400&gt; 31

Met Cys Gly Arg Phe	1	Leu Arg Arg Leu Leu	10	Ala Glu Glu Ser Arg	15
Arg Ser Thr Pro Val	20	Gly Arg Leu Leu Leu	25	Pro Val Leu Leu Gly	30
Phe Arg Leu Val Leu	35	Leu Ala Ala Ser Gly	40	Pro Gly Val Tyr Gly	45
Asp Glu Gln Ser Glu	50	Phe Val Cys His Thr	55	Gln Gln Pro Gly Cys	60
Lys Ala Ala Cys Phe	65	Asp Ala Phe His Pro	70	Leu Ser Pro Leu Arg	75
Ser Trp Val Phe Gln	80	Val Ile Leu Val Ala	85	Val Pro Ser Ala Leu	90
Tyr Met Gly Phe Thr	95	Leu Tyr His Val Ile	100	Trp His Trp Glu Leu	105
Ser Gly Lys Gly Lys	110	Glu Glu Glu Thr Leu	115	Ile Gln Gly Arg Glu	120
Gly Asn Thr Asp Val	125	Pro Gly Ala Gly Ser	130	Leu Arg Leu Leu Trp	135
Ala Tyr Val Ala Gln	140	Leu Gly Ala Arg Leu	145	Val Leu Glu Gly Ala	150
Ala Leu Gly Leu Gln	155	Tyr His Leu Tyr Gly	160	Phe Gln Met Pro Ser	165
Ser Phe Ala Cys Arg	170	Arg Glu Pro Cys Leu	175	Gly Ser Ile Thr Cys	180
Asn Leu Ser Arg Pro		Ser Glu Lys Thr Ile		Phe Leu Lys Thr Met	



Phe Gly Val Ser	185	Gly Phe Cys Leu Leu	190	Phe Thr Phe Leu Glu Leu	195
Val Leu Leu Gly	200	Leu Gly Arg Trp Trp	205	Arg Thr Trp Lys His Lys	210
Ser Ser Ser Ser	215	Lys Tyr Phe Leu Thr	220	Ser Glu Ser Thr Arg Arg	225
His Lys Lys Ala	230	Thr Asp Ser Leu Pro	235	Val Val Glu Thr Lys Glu	240
Gln Phe Gln Glu	245	Ala Val Pro Gly Arg	250	Ser Leu Ala Gln Glu Lys	255
Gln Arg Pro Val	260	Gly Pro Arg Asp Ala	265		270
	275				

<210> 32  
 <211> 154  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2219267CD1

Met Val Thr Gly Leu	Ala Ser Leu Leu Leu	Leu Ala Gly Ala Gln
1	5	10
Tyr Leu Pro Gly Trp	Thr Val Leu Phe Leu	Ser Val Leu Gly Leu
	20	25
Leu Ala Ser Arg Ala	Val Ser Ala Leu Ser	Ser Leu Phe Ala Ala
	35	40
Glu Val Phe Pro Thr	Val Ile Arg Gly Ala	Gly Leu Gly Leu Val
	50	55
Leu Gly Ala Gly Phe	Leu Gly Gln Ala Ala	Gly Pro Leu Asp Thr
	65	70
Leu His Gly Arg Gln	Gly Phe Phe Leu Gln	Gln Val Val Phe Ala
	80	85
Ser Leu Ala Val Leu	Ala Leu Leu Cys Val	Leu Leu Leu Pro Glu
	95	100
Ser Arg Ser Arg Gly	Leu Pro Gln Ser Leu	Gln Asp Ala Asp Arg
	110	115
Leu Arg Arg Ser Pro	Leu Leu Arg Gly Arg	Pro Arg Gln Asp His
	125	130
Leu Pro Leu Leu Pro	Pro Ser Asn Ser Tyr	Trp Ala Gly His Thr
	140	145
Pro Glu Gln His		150

<210> 33  
 <211> 289  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2308629CD1

Met Val Ala Gly Ala	Val Ala Gly Ile Leu	Glu His Cys Val Met
1	5	10
Tyr Pro Ile Asp Cys	Val Lys Thr Arg Met	Gln Ser Leu Gln Pro
	20	25
Asp Pro Ala Ala Arg	Tyr Arg Asn Val Leu	Glu Ala Leu Trp Arg
	35	40
Ile Ile Arg Thr Glu	Gly Leu Trp Arg Pro	Met Arg Gly Leu Asn
	50	55
Val Thr Ala Thr Gly	Ala Gly Pro Ala His	Ala Leu Tyr Phe Ala
	65	70
Cys Tyr Glu Lys Leu	Lys Lys Thr Leu Ser	Asp Val Ile His Pro
	80	85
Gly Gly Asn Ser His	Ile Ala Asn Gly Ala	Ala Gly Cys Val Ala

	95		100		105
Thr	Leu	Leu	His	Asp	Ala
	110		115		120
Gln	Arg	Met	Gln	Met	Tyr
	125		130		135
Cys	Val	Arg	Ala	Val	Trp
	140		145		150
Arg	Ser	Tyr	Thr	Thr	Gln
	155		160		165
Ile	His	Phe	Met	Thr	Tyr
	170		175		180
Gln	Arg	Arg	Tyr	Asn	Pro
	185		190		195
Ala	Gly	Ala	Val	Ala	Ala
	200		205		210
Lys	Thr	Leu	Leu	Asn	Thr
	215		220		225
Ile	Thr	Gly	His	Ile	Thr
	230		235		240
Tyr	Gln	Val	Gly	Gly	Val
	245		250		255
Arg	Val	Ile	Tyr	Gln	Ile
	260		265		270
Tyr	Glu	Phe	Phe	Lys	Tyr
	275		280		285
Arg	Ala	Gly	Lys		

<210> 34  
 <211> 300  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2660038CD1

<400> 34

Met	Asp	Phe	Leu	Met	Ser	Gly	Leu	Ala	Ala	Cys	Gly	Ala	Cys	Val
1				5					10					15
Phe	Thr	Asn	Pro	Leu	Glu	Val	Val	Lys	Thr	Arg	Met	Gln	Leu	Gln
				20					25					30
Gly	Glu	Leu	Gln	Ala	Pro	Gly	Thr	Tyr	Gln	Arg	His	Tyr	Arg	Asn
				35					40					45
Val	Phe	His	Ala	Phe	Ile	Thr	Ile	Gly	Lys	Val	Asp	Gly	Leu	Ala
				50					55					60
Ala	Leu	Gln	Lys	Gly	Leu	Ala	Pro	Ala	Leu	Leu	Tyr	Gln	Phe	Leu
				65					70					75
Met	Asn	Gly	Ile	Arg	Leu	Gly	Thr	Tyr	Gly	Leu	Ala	Glu	Ala	Gly
				80					85					90
Gly	Tyr	Leu	His	Thr	Ala	Glu	Ala	Thr	His	Ser	Pro	Ala	Arg	Ser
				95					100					105
Ala	Ala	Ala	Gly	Ala	Met	Ala	Gly	Val	Met	Gly	Ala	Tyr	Leu	Gly
				110					115					120
Ser	Pro	Ile	Tyr	Met	Val	Lys	Thr	His	Leu	Gln	Ala	Gln	Ala	Ala
				125					130					135
Ser	Glu	Ile	Ala	Val	Gly	His	Gln	Tyr	Lys	His	Gln	Gly	Met	Phe
				140					145					150
Gln	Ala	Leu	Thr	Glu	Ile	Gly	Gln	Lys	His	Gly	Leu	Val	Gly	Leu
				155					160					165
Trp	Arg	Gly	Ala	Leu	Gly	Gly	Leu	Pro	Arg	Val	Ile	Val	Gly	Ser
				170					175					180
Ser	Thr	Gln	Leu	Cys	Thr	Phe	Ser	Ser	Thr	Lys	Asp	Leu	Leu	Ser
				185					190					195
Gln	Trp	Glu	Ile	Phe	Pro	Pro	Gln	Ser	Trp	Lys	Leu	Ala	Leu	Val
				200					205					210
Ala	Ala	Met	Met	Ser	Gly	Ile	Ala	Val	Val	Leu	Ala	Met	Ala	Pro
				215					220					225
Phe	Asp	Val	Ala	Cys	Thr	Arg	Leu	Tyr	Asn	Gln	Pro	Thr	Asp	Ala

Gln Gly Lys Gly	230	235	240
Leu Met Tyr Arg Gly	245	Ile Leu Asp Ala Leu Leu	255
Gln Thr Ala Arg	260	Gly Met Tyr Lys Gly	270
Gly Ala Ser Tyr	275	His Thr Ile Leu Ser	285
Phe Phe Trp Asp	290	Tyr Tyr Thr Asp Thr	300

&lt;210&gt; 35

&lt;211&gt; 382

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2670745CD1

&lt;400&gt; 35

Met Leu Arg Trp Thr	Val His Leu Glu Gly	Gly Pro Arg Arg Val	15
Asn His Ala Ala Val	Ala Val Gly His Arg	Val Tyr Ser Phe Gly	30
Gly Tyr Cys Ser Gly	Glu Asp Tyr Glu Thr	Leu Arg Gln Ile Asp	45
Val His Ile Phe Asn	Ala Val Ser Leu Arg	Trp Thr Lys Leu Pro	60
Pro Val Lys Ser Ala	Ile Arg Gly Gln Ala	Pro Val Val Pro Tyr	75
Met Arg Tyr Gly His	Ser Thr Val Leu Ile	Asp Asp Thr Val Leu	90
Leu Trp Gly Gly Arg	Asn Asp Thr Glu Gly	Ala Cys Asn Val Leu	105
Tyr Ala Phe Asp Val	Asn Thr His Lys Trp	Phe Thr Pro Arg Val	120
Ser Gly Thr Val Pro	Gly Ala Arg Asp Gly	His Ser Ala Cys Val	135
Leu Gly Lys Ile Met	Tyr Ile Phe Gly Gly	Tyr Glu Gln Gln Ala	150
Asp Cys Phe Ser Asn	Asp Ile His Lys Leu	Asp Thr Ser Thr Met	165
Thr Trp Thr Leu Ile	Cys Thr Lys Gly Ser	Pro Ala Arg Trp Arg	180
Asp Phe His Ser Ala	Thr Met Leu Gly Ser	His Met Tyr Val Phe	195
Gly Gly Arg Ala Asp	Arg Phe Gly Pro Phe	His Ser Asn Asn Glu	210
Ile Tyr Cys Asn Arg	Ile Arg Val Phe Asp	Thr Arg Thr Glu Ala	225
Trp Leu Asp Cys Pro	Pro Thr Pro Val Leu	Pro Glu Gly Arg Arg	240
Ser His Ser Ala Phe	Gly Tyr Asn Gly Glu	Leu Tyr Ile Phe Gly	255
Gly Tyr Asn Ala Arg	Leu Asn Arg His Phe	His Asp Leu Trp Lys	270
Phe Asn Pro Val Ser	Phe Thr Trp Lys Lys	Ile Glu Pro Lys Gly	285
Lys Gly Pro Cys Pro	Arg Arg Arg Gln Cys	Cys Cys Ile Val Gly	300
Asp Lys Ile Val Leu	Phe Gly Gly Thr Ser	Pro Ser Pro Glu Glu	315
Gly Leu Gly Asp Glu	Phe Asp Leu Ile Asp	His Ser Asp Leu His	330
Ile Leu Asp Phe Ser	Pro Ser Leu Lys Thr	Leu Cys Lys Leu Ala	345
Val Ile Gln Tyr Asn	Leu Asp Gln Ser Cys	Leu Pro His Asp Ile	360

Arg Trp Glu Leu Asn Ala Met Thr Thr Asn Ser Asn Ile Ser Arg  
 365 370 375  
 Pro Ile Val Ser Ser His Gly  
 380

<210> 36  
 <211> 287  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2676443CD1

<400> 36  
 Met Ala Ala Glu Ala Arg Val Ser Arg Trp Tyr Phe Gly Gly Leu  
 1 5 10 15  
 Ala Ser Cys Gly Ala Ala Cys Cys Thr His Pro Leu Asp Leu Leu  
 20 25 30  
 Lys Val His Leu Gln Thr Gln Gln Glu Val Lys Leu Arg Met Thr  
 35 40 45  
 Gly Met Ala Leu Arg Val Val Arg Thr Asp Gly Ile Leu Ala Leu  
 50 55 60  
 Tyr Ser Gly Leu Ser Ala Ser Leu Cys Arg Gln Met Thr Tyr Ser  
 65 70 75  
 Leu Thr Arg Phe Ala Ile Tyr Glu Thr Val Arg Asp Arg Val Ala  
 80 85 90  
 Lys Gly Ser Gln Gly Pro Leu Pro Phe His Glu Lys Val Leu Leu  
 95 100 105  
 Gly Ser Val Ser Gly Leu Ala Gly Gly Phe Val Gly Thr Pro Ala  
 110 115 120  
 Asp Leu Val Asn Val Arg Met Gln Asn Asp Val Lys Leu Pro Gln  
 125 130 135  
 Gly Gln Arg Arg Asn Tyr Ala His Ala Leu Asp Gly Leu Tyr Arg  
 140 145 150  
 Val Ala Arg Glu Glu Gly Leu Arg Arg Leu Phe Ser Gly Ala Thr  
 155 160 165  
 Met Ala Ser Ser Arg Gly Ala Leu Val Thr Val Gly Gln Leu Ser  
 170 175 180  
 Cys Tyr Asp Gln Ala Lys Gln Leu Val Leu Ser Thr Gly Tyr Leu  
 185 190 195  
 Ser Asp Asn Ile Phe Thr His Phe Val Ala Ser Phe Ile Ala Gly  
 200 205 210  
 Gly Cys Ala Thr Phe Leu Cys Gln Pro Leu Asp Val Leu Lys Thr  
 215 220 225  
 Arg Leu Met Asn Ser Lys Gly Glu Tyr Gln Gly Val Phe His Cys  
 230 235 240  
 Ala Val Glu Thr Ala Lys Leu Gly Pro Leu Ala Phe Tyr Lys Gly  
 245 250 255  
 Leu Val Pro Ala Gly Ile Arg Leu Ile Pro His Thr Val Leu Thr  
 260 265 270  
 Phe Val Phe Leu Glu Gln Leu Arg Lys Asn Phe Gly Ile Lys Val  
 275 280 285  
 Pro Ser

<210> 37  
 <211> 497  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3295764CD1

<400> 37  
 Met Asp Val Pro Gly Pro Val Ser Arg Arg Ala Ala Ala Ala Ala  
 1 5 10 15  
 Ala Thr Val Leu Leu Arg Thr Ala Arg Val Arg Arg Glu Cys Trp  
 20 25 30

Phe	Leu	Pro	Thr	Ala	Leu	Leu	Cys	Ala	Tyr	Gly	Phe	Phe	Ala	Ser
				35					40					45
Leu	Arg	Pro	Ser	Glu	Pro	Phe	Leu	Thr	Pro	Tyr	Leu	Leu	Gly	Pro
				50					55					60
Asp	Lys	Asn	Leu	Thr	Glu	Arg	Glu	Val	Phe	Asn	Glu	Ile	Tyr	Pro
				65					70					75
Val	Trp	Thr	Tyr	Ser	Tyr	Leu	Val	Leu	Leu	Phe	Pro	Val	Phe	Leu
				80					85					90
Ala	Thr	Asp	Tyr	Leu	Arg	Tyr	Lys	Pro	Val	Val	Leu	Leu	Gln	Gly
				95					100					105
Leu	Ser	Leu	Ile	Val	Thr	Trp	Phe	Met	Leu	Leu	Tyr	Ala	Gln	Gly
				110					115					120
Leu	Leu	Ala	Ile	Gln	Phe	Leu	Glu	Phe	Phe	Tyr	Gly	Ile	Ala	Thr
				125					130					135
Ala	Thr	Glu	Ile	Ala	Tyr	Tyr	Ser	Tyr	Ile	Tyr	Ser	Val	Val	Asp
				140					145					150
Leu	Gly	Met	Tyr	Gln	Lys	Val	Thr	Ser	Tyr	Cys	Arg	Ser	Ala	Thr
				155					160					165
Leu	Val	Gly	Phe	Thr	Val	Gly	Ser	Val	Leu	Gly	Gln	Ile	Leu	Val
				170					175					180
Ser	Val	Ala	Gly	Trp	Ser	Leu	Phe	Ser	Leu	Asn	Val	Ile	Ser	Leu
				185					190					195
Thr	Cys	Val	Ser	Val	Ala	Phe	Ala	Val	Ala	Trp	Phe	Leu	Pro	Met
				200					205					210
Pro	Gln	Lys	Ser	Leu	Phe	Phe	His	His	Ile	Pro	Ser	Thr	Cys	Gln
				215					220					225
Arg	Val	Asn	Gly	Ile	Lys	Val	Gln	Asn	Gly	Gly	Ile	Val	Thr	Asp
				230					235					240
Thr	Pro	Ala	Ser	Asn	His	Leu	Pro	Gly	Trp	Glu	Asp	Ile	Glu	Ser
				245					250					255
Lys	Ile	Pro	Leu	Asn	Met	Glu	Glu	Pro	Pro	Val	Glu	Glu	Pro	Glu
				260					265					270
Pro	Lys	Pro	Asp	Arg	Leu	Leu	Val	Leu	Lys	Val	Leu	Trp	Asn	Asp
				275					280					285
Phe	Leu	Met	Cys	Tyr	Ser	Ser	Arg	Pro	Leu	Leu	Cys	Trp	Ser	Val
				290					295					300
Trp	Trp	Ala	Leu	Ser	Thr	Cys	Gly	Tyr	Phe	Gln	Val	Val	Asn	Tyr
				305					310					315
Thr	Gln	Gly	Leu	Trp	Glu	Lys	Val	Met	Pro	Ser	Arg	Tyr	Ala	Ala
				320					325					330
Ile	Tyr	Asn	Gly	Gly	Val	Glu	Ala	Val	Ser	Thr	Leu	Leu	Gly	Ala
				335					340					345
Val	Ala	Val	Phe	Ala	Val	Gly	Tyr	Ile	Lys	Ile	Ser	Trp	Ser	Thr
				350					355					360
Trp	Gly	Glu	Met	Thr	Leu	Ser	Leu	Phe	Ser	Leu	Leu	Ile	Ala	Ala
				365					370					375
Ala	Val	Tyr	Ile	Met	Asp	Thr	Val	Gly	Asn	Ile	Trp	Val	Cys	Tyr
				380					385					390
Ala	Ser	Tyr	Val	Val	Phe	Arg	Ile	Ile	Tyr	Met	Leu	Leu	Ile	Thr
				395					400					405
Ile	Ala	Thr	Phe	Gln	Ile	Ala	Ala	Asn	Leu	Ser	Met	Glu	Arg	Tyr
				410					415					420
Ala	Leu	Val	Phe	Gly	Val	Asn	Thr	Phe	Ile	Ala	Leu	Ala	Leu	Gln
				425					430					435
Thr	Leu	Leu	Thr	Leu	Ile	Val	Val	Asp	Ala	Ser	Gly	Leu	Gly	Leu
				440					445					450
Glu	Ile	Thr	Thr	Gln	Phe	Leu	Ile	Tyr	Ala	Ser	Tyr	Phe	Ala	Leu
				455					460					465
Ile	Ala	Val	Val	Phe	Leu	Ala	Ser	Gly	Ala	Val	Ser	Val	Met	Lys
				470					475					480
Lys	Cys	Arg	Lys	Leu	Glu	Asp	Pro	Gln	Ser	Ser	Ser	Gln	Val	Thr
				485					490					495
Thr	Ser													

&lt;210&gt; 38

&lt;211&gt; 228

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3438320CD1

<400> 38

Met	Pro	Arg	Arg	Gly	Leu	Val	Ala	Gly	Pro	Asp	Leu	Glu	Tyr	Phe	
1				5					10					15	
Gln	Arg	Arg	Tyr	Phe	Thr	Pro	Ala	Glu	Val	Ala	Gln	His	Asn	Arg	
				20					25					30	
Pro	Glu	Asp	Leu	Trp	Val	Ser	Tyr	Leu	Gly	Arg	Val	Tyr	Asp	Leu	
				35					40					45	
Thr	Ser	Leu	Ala	Gln	Glu	Tyr	Lys	Gly	Asn	Leu	Leu	Leu	Lys	Pro	
				50					55					60	
Ile	Val	Glu	Val	Ala	Gly	Gln	Asp	Ile	Ser	His	Trp	Phe	Asp	Pro	
				65					70					75	
Lys	Thr	Arg	Asp	Ile	Arg	Lys	His	Ile	Asp	Pro	Leu	Thr	Gly	Cys	
				80					85					90	
Leu	Arg	Tyr	Cys	Thr	Pro	Arg	Gly	Arg	Phe	Val	His	Val	Pro	Pro	
				95					100					105	
Gln	Leu	Pro	Cys	Ser	Asp	Trp	Ala	Asn	Asp	Phe	Gly	Lys	Pro	Trp	
				110					115					120	
Trp	Gln	Gly	Ser	Tyr	Tyr	Glu	Val	Gly	Arg	Leu	Ser	Ala	Lys	Thr	
				125					130					135	
Arg	Ser	Ile	Arg	Ile	Ile	Asn	Thr	Leu	Thr	Ser	Gln	Glu	His	Thr	
				140					145					150	
Leu	Glu	Val	Gly	Val	Leu	Glu	Ser	Ile	Trp	Glu	Ile	Leu	His	Arg	
				155					160					165	
Tyr	Leu	Pro	Tyr	Asn	Ser	His	Ala	Ala	Ser	Tyr	Thr	Trp	Lys	Tyr	
				170					175					180	
Glu	Gly	Lys	Asn	Leu	Asn	Met	Asp	Phe	Thr	Leu	Glu	Glu	Asn	Gly	
				185					190					195	
Ile	Arg	Asp	Glu	Glu	Glu	Glu	Phe	Asp	Tyr	Leu	Ser	Met	Asp	Gly	
				200					205					210	
Thr	Leu	His	Thr	Pro	Ala	Ile	Leu	Leu	Tyr	Phe	Asn	Asp	Asp	Leu	
				215					220					225	
Thr	Glu	Leu													

<210> 39  
 <211> 273  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3986488CD1

<400> 39

Met	Ala	Ala	Thr	Ile	Met	Ile	Leu	Tyr	Val	Ser	Lys	Leu	Asn	Lys	
1				5					10					15	
Ile	Ile	His	Phe	Pro	Asp	Phe	Asp	Lys	Lys	Ile	Pro	Val	Lys	Leu	
				20					25					30	
Phe	Pro	Leu	Pro	Leu	Leu	Tyr	Val	Gly	Asn	His	Ile	Ser	Gly	Leu	
				35					40					45	
Ser	Ser	Thr	Ser	Lys	Leu	Ser	Leu	Pro	Met	Phe	Thr	Val	Leu	Arg	
				50					55					60	
Lys	Phe	Thr	Ile	Pro	Leu	Thr	Leu	Leu	Glu	Thr	Ile	Ile	Leu		
				65					70					75	
Gly	Lys	Gln	Tyr	Ser	Leu	Asn	Ile	Ile	Leu	Ser	Val	Phe	Ala	Ile	
				80					85					90	
Ile	Leu	Gly	Ala	Phe	Ile	Ala	Ala	Gly	Ser	Asp	Leu	Ala	Phe	Asn	
				95					100					105	
Leu	Glu	Gly	Tyr	Ile	Phe	Val	Phe	Leu	Asn	Asp	Ile	Phe	Thr	Ala	
				110					115					120	
Ala	Asn	Gly	Val	Tyr	Thr	Lys	Gln	Lys	Met	Asp	Pro	Lys	Glu	Leu	
				125					130					135	
Gly	Lys	Tyr	Gly	Val	Leu	Phe	Tyr	Asn	Ala	Cys	Phe	Met	Ile	Ile	
				140					145					150	
Pro	Thr	Leu	Ile	Ile	Ser	Val	Ser	Thr	Gly	Asp	Leu	Gln	Gln	Ala	

Thr	Glu	Phe	Asn	Gln	Trp	Lys	Asn	Val	Val	Phe	Ile	Leu	Gln	Phe
155									160					165
Leu	Leu	Ser	Cys	Phe	Leu	Gly	Phe	Leu	Leu	Met	Tyr	Ser	Thr	Val
170									175					180
Leu	Cys	Ser	Tyr	Tyr	Asn	Ser	Ala	Leu	Thr	Thr	Ala	Val	Val	Gly
185									190					195
Ala	Ile	Lys	Asn	Val	Ser	Val	Ala	Tyr	Ile	Gly	Ile	Leu	Ile	Gly
200									205					210
Gly	Asp	Tyr	Ile	Phe	Ser	Leu	Leu	Asn	Phe	Val	Gly	Leu	Asn	Ile
215									220					225
Cys	Met	Ala	Gly	Gly	Leu	Arg	Tyr	Ser	Phe	Leu	Thr	Leu	Ser	Ser
230									235					240
Gln	Leu	Lys	Pro	Lys	Pro	Val	Gly	Glu	Glu	Asn	Ile	Cys	Leu	Asp
245									250					255
Leu	Lys	Ser							265					270

<210> 40  
 <211> 206  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4378816CD1

Met	Gly	Ala	Glu	Trp	Glu	Leu	Gly	Ala	Glu	Ala	Gly	Gly	Ser	Leu
1					5				10					15
Leu	Leu	Cys	Ala	Ala	Leu	Leu	Ala	Ala	Gly	Cys	Ala	Leu	Gly	Leu
20					25				30					35
Arg	Leu	Gly	Arg	Gly	Gln	Gly	Ala	Ala	Asp	Arg	Gly	Ala	Leu	Ile
35					40				45					50
Trp	Leu	Cys	Tyr	Asp	Ala	Leu	Val	His	Phe	Ala	Leu	Glu	Gly	Pro
50					55				60					65
Phe	Val	Tyr	Leu	Ser	Leu	Val	Gly	Asn	Val	Ala	Asn	Ser	Asp	Gly
65					70				75					80
Leu	Ile	Ala	Ser	Leu	Trp	Lys	Glu	Tyr	Gly	Lys	Ala	Asp	Ala	Arg
80					85				90					95
Trp	Val	Tyr	Phe	Asp	Pro	Thr	Ile	Val	Ser	Val	Glu	Ile	Leu	Thr
95					100				105					110
Val	Ala	Leu	Asp	Gly	Ser	Leu	Ala	Leu	Phe	Leu	Ile	Tyr	Ala	Ile
110					115				120					125
Val	Lys	Glu	Lys	Tyr	Tyr	Arg	His	Phe	Leu	Gln	Ile	Thr	Leu	Cys
125					130				135					140
Val	Cys	Glu	Leu	Tyr	Gly	Cys	Trp	Met	Thr	Phe	Leu	Pro	Glu	Trp
140					145				150					155
Leu	Thr	Arg	Ser	Pro	Asn	Leu	Asn	Thr	Ser	Asn	Trp	Leu	Tyr	Cys
155					160				165					170
Trp	Leu	Tyr	Leu	Phe	Phe	Phe	Asn	Gly	Val	Trp	Val	Leu	Ile	Pro
170					175				180					185
Gly	Leu	Leu	Leu	Trp	Gln	Ser	Trp	Leu	Glu	Leu	Lys	Lys	Met	His
185					190				195					200
Gln	Lys	Glu	Thr	Ser	Ser	Val	Lys	Lys	Phe	Gln				
200					205									

<210> 41  
 <211> 235  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4797137CD1

Met	Gln	Gln	Arg	Gly	Ala	Ala	Gly	Ser	Arg	Gly	Cys	Ala	Leu	Phe
1					5				10					15
Pro	Leu	Leu	Gly	Val	Leu	Phe	Phe	Gln	Gly	Val	Tyr	Ile	Val	Phe

	20		25		30									
Ser	Leu	Glu	Ile	Arg	Ala	Asp	Ala	His	Val	Arg	Gly	Tyr	Val	Gly
	35		40		45									
Glu	Lys	Ile	Lys	Leu	Lys	Cys	Thr	Phe	Lys	Ser	Thr	Ser	Asp	Val
	50		55		60									
Thr	Asp	Lys	Leu	Thr	Ile	Asp	Trp	Thr	Tyr	Arg	Pro	Pro	Ser	Ser
	65		70		75									
Ser	His	Thr	Val	Ser	Ile	Phe	His	Tyr	Gln	Ser	Phe	Gln	Tyr	Pro
	80		85		90									
Thr	Thr	Ala	Gly	Thr	Phe	Arg	Asp	Arg	Ile	Ser	Trp	Val	Gly	Asn
	95		100		105									
Val	Tyr	Lys	Gly	Asp	Ala	Ser	Ile	Ser	Ile	Ser	Asn	Pro	Thr	Ile
	110		115		120									
Lys	Asp	Asn	Gly	Thr	Phe	Ser	Cys	Ala	Val	Lys	Asn	Pro	Pro	Asp
	125		130		135									
Val	His	His	Asn	Ile	Pro	Met	Thr	Glu	Leu	Thr	Val	Thr	Glu	Arg
	140		145		150									
Gly	Phe	Gly	Thr	Met	Leu	Ser	Ser	Val	Ala	Leu	Leu	Ser	Ile	Leu
	155		160		165									
Val	Phe	Val	Pro	Ser	Ala	Val	Val	Val	Ala	Leu	Leu	Leu	Val	Arg
	170		175		180									
Met	Gly	Arg	Lys	Ala	Ala	Gly	Leu	Lys	Lys	Arg	Ser	Arg	Ser	Gly
	185		190		195									
Tyr	Lys	Lys	Ser	Ser	Ile	Glu	Val	Ser	Asp	Asp	Thr	Asp	Gln	Glu
	200		205		210									
Glu	Glu	Glu	Ala	Cys	Met	Ala	Arg	Leu	Cys	Val	Arg	Cys	Ala	Glu
	215		220		225									
Cys	Leu	Asp	Ser	Asp	Tyr	Glu	Glu	Thr	Tyr					
	230		235											

&lt;210&gt; 42

&lt;211&gt; 147

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5470806CD1

&lt;400&gt; 42

Met	Ala	Ser	Leu	Arg	Leu	Phe	Leu	Leu	Cys	Leu	Ala	Val	Leu	Ile
1				5					10					15
Phe	Ala	Ser	Glu	Ala	Gly	Pro	Gly	Gly	Ala	Gly	Glu	Ser	Lys	Cys
			20						25					30
Pro	Leu	Met	Val	Lys	Val	Leu	Asp	Ala	Val	Arg	Gly	Ser	Pro	Ala
			35						40					45
Val	Asp	Val	Ala	Val	Lys	Val	Phe	Lys	Lys	Thr	Ala	Asp	Gly	Ser
			50						55					60
Trp	Glu	Pro	Phe	Ala	Ser	Gly	Lys	Thr	Ala	Glu	Ser	Gly	Glu	Leu
			65						70					75
His	Gly	Leu	Thr	Thr	Asp	Glu	Lys	Phe	Thr	Glu	Gly	Val	Tyr	Arg
			80						85					90
Val	Glu	Leu	Asp	Thr	Lys	Ser	Tyr	Trp	Lys	Ala	Leu	Gly	Ile	Ser
			95						100					105
Pro	Phe	His	Glu	Tyr	Ala	Glu	Val	Val	Phe	Thr	Ala	Asn	Asp	Ser
			110						115					120
Gly	His	Arg	His	Tyr	Thr	Ile	Ala	Ala	Leu	Leu	Ser	Pro	Tyr	Ser
			125						130					135
Tyr	Ser	Thr	Thr	Ala	Val	Val	Ser	Asn	Pro	Gln	Asn			
			140						145					

&lt;210&gt; 43

&lt;211&gt; 147

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5473242CD1



&lt;400&gt; 43

Met	Val	His	Leu	Thr	Asp	Ala	Glu	Lys	Ala	Thr	Val	Asn	Gly	Leu
1				5					10					15
Trp	Gly	Lys	Val	Asn	Pro	Val	Glu	Ile	Gly	Ala	Glu	Ser	Leu	Ala
				20					25					30
Ser	Leu	Leu	Ile	Val	Tyr	Pro	Trp	Thr	Gln	Arg	Tyr	Phe	Ser	Lys
				35					40					45
Phe	Gly	Asp	Leu	Ser	Ser	Val	Ser	Ala	Ile	Met	Gly	Asn	Pro	Gln
				50					55					60
Val	Lys	Ala	His	Gly	Glu	Lys	Val	Ile	Asn	Ala	Phe	Asp	Asp	Gly
				65					70					75
Leu	Lys	His	Leu	Asp	Asn	Leu	Lys	Gly	Thr	Phe	Ala	Ser	Leu	Ser
				80					85					90
Glu	Leu	His	Cys	Asp	Lys	Leu	His	Val	Asp	Pro	Glu	Asn	Phe	Arg
				95					100					105
Leu	Leu	Gly	Asn	Met	Ile	Val	Ile	Met	Met	Gly	His	His	Leu	Gly
				110					115					120
Lys	Glu	Phe	Thr	Pro	Ser	Ala	Gln	Ala	Ala	Phe	Gln	Lys	Val	Val
				125					130					135
Ala	Gly	Val	Ala	Ser	Ala	Leu	Ala	His	Lys	Tyr	His			
				140					145					

&lt;210&gt; 44

&lt;211&gt; 2701

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 264114CB1

&lt;400&gt; 44

gcggcgccgc	cagcttcttc	ggccggagg	gaggcgagac	cccaggcgag	gccgcggcgg	60
gagggccacg	cccccgacgc	cgcccgagg	ggggccagtg	tgagcggggc	caccggctgg	120
agcggtatccc	acacctccgg	accgagggac	gcggttactc	cacaggatcc	gctgaacata	180
ggatgttgcc	acaaaatcta	cctcgtgtat	ttttctcttt	cactcatgag	ctgcacaatt	240
gcagatttga	gcacaatgtc	tgcaagctgt	gttgaaaaac	tctgaagaac	ctaattaaca	300
caggatgacc	taggagtgat	tctaagctctg	tgtaacaaga	tattactcat	tagtgaatgt	360
gtcagtcttg	gtactgaatg	ctgcagataa	cagcaagtag	gttctccttt	atttctgaag	420
tattcacttg	accttccatc	agtaagacgg	acttttctaa	tctgttcctg	gagatattaa	480
tggaatacac	tcatgtccac	tcaagacgag	aggcagatca	atactgaata	tgctgtgtca	540
ttgttggaac	agttgaaact	gttttatgaa	cagcagttgt	ttactgacat	agtggttaatt	600
gttgagggca	ctgaattccc	ttgtcataag	atggttcttg	caacatgtag	ctcttatttc	660
agggccatgt	ttatgagtgg	actaagtga	agcaaacaaa	cccatgtaca	cctgaggaat	720
gtcgattgtg	ccaccttaca	gataataata	acttatgcat	acacgggtaa	cttggcaatg	780
aatgacagca	ctgtagaaca	gctttatgaa	acagcttgct	tcctacaggt	agaagatgtg	840
ttacaacgtt	gtcgagaata	tttaattaaa	aaaataaatg	cagagaattg	tgtacgattg	900
ttgagttttg	ctgatctctt	cagttgtgag	gaattaaaac	agagtgtctaa	aagaatggtg	960
gagcacaaat	tcatgtctgt	gtatcatcag	gacgcgttca	tgcagctgtc	acatgacata	1020
ctgatagata	ttctcagtag	tgacaattta	aatgtagaaa	aggaagaaac	cgttcgagaa	1080
gctgctatgc	tgtggctaga	gtataacaca	gaatcacgat	cccagtatct	gtcttctgtt	1140
cttagccaaa	tcagaattga	tgcaatttca	gaagtaacac	agagagcttg	gtttcaaggt	1200
ctggcaccaca	atgataagtc	agtgggtggt	caaggtctgt	ataagtccat	gcccaagttt	1260
ttcaaaccac	gacttgggat	gactaaagag	gaaatgatga	ttttcattga	agcatcttca	1320
gaaaatcctt	gtagtcttta	ctcttctgtc	tgttacagcc	cccaagcaga	aaaagtttac	1380
aagttatgta	gccaccagc	tgatttgcac	aaggttggga	ccgttgtaac	tcctgataat	1440
gatattctaca	tagcaggggg	tcaagttcct	ctgaaaaaca	caaaaacaaa	tcacagtaaa	1500
acaagcaaac	ttcagactgc	cttcagaact	gtgaattgct	tttattgggt	tgatgcacag	1560
caaaatacct	ggtttccaaa	gaccccaatg	ctttttgtcc	gcataaagcc	atctttggtt	1620
tgctgtgaag	gctatatcta	tgcaattgga	ggagatagcg	taggtggaga	acttaactcg	1680
aggaccgtag	aaagatacga	cactgagaaa	gatgagtgga	cgatggtaag	ccctttacct	1740
tgtgcttggc	aatggagtgc	agcagttgtg	gttcagtact	gcatttatgt	gatgacactg	1800
aacctcatgt	actgttattt	tccaaggtct	gactcatggg	tagaaatggc	catgagacag	1860
actagtaggt	cctttgcttc	agctgcagct	tttggtgata	aaattttcta	tattggaggg	1920
ttgcataattg	tcaccaattc	cggcataaga	ctcccctctg	gcactgtaga	tggtcttcca	1980
gtaactgtgg	aaatttatga	tgtgaataaa	aatgagtggg	aaatggcagc	caacatccct	2040
gctaagaggt	actctgaccc	ctgtgttaga	gctgttgtga	tctcaaattc	tctatgtgtg	2100
tttatgcgag	aaaccacact	aatgagcgca	gctaaatacg	tcacctacca	atatgacctg	2160
gaacttgacc	ggtggtctct	gcggcagcat	atatctgaac	gtgtactgtg	ggacttgggg	2220

```

agagattttc gatgcactgt ggggaaactc tatccatcct gccttgaaga gtctccatgg 2280
aaaccaccaa cttatctttt ttcaacggat gggacagaag agtttgaact ggatggagaa 2340
atggttgacac taccacctgt atagtgggga agttcagggg gtgcacgcct gagttatgtg 2400
ctttgtcatt ttctttgcta aacaaaagag gctatgaaag aactaaatat gagtacataa 2460
aattctatct ttgataaatt ttatttttat gccctactta atatttgcac cagtataata 2520
tatatcagtg agtcttacag aaagatatgc ttccataata tgaaatagat tattcaataa 2580
ttgagaaact ttatgtgtaa tcatgagagt ataagaatct ggattatcta acattgttag 2640
ccctgtgtat gtacagttca aaaagttcat ttataaaagt agtttcctgt tcctagttga 2700
t 2701

```

&lt;210&gt; 45

&lt;211&gt; 736

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1455669CB1

&lt;400&gt; 45

```

gagacttagc gacagacaga cgctgggacc cacgacgaca gaaggcgccg atggccgcgc 60
ctgctgagcc ctgcgcgggg caggggggtct ggaaccagac agagcctgaa cctgcgcgca 120
ccagcctgct gagcctgtgc ttcttgagaa cagcaggggt ctgggtaccc cccatgtacc 180
tctgggtcct tgggtcccatc tacctcctct tcatccacca ccatggccgg ggctacctcc 240
ggatgtcccc actcttcaaa gccaaagatgg tagctgccat ccctgggagc ctggaaccag 300
gcaatgttct ggggaggcag gggacaggct ggaacctggt gaagtcttaa agtagactcc 360
tcctatcggg gtgtagaagg gaatctgtta atcaaacaga gcaatattag aaaggctaca 420
gaggtcaact cagtggaaac cggttctctcc aaacagatct tgtaattccg aaaatccacg 480
catgcgcaaa catacgcata cactcccatg ttcttggaac gtttatagct accataacct 540
ggcattttcc aaacataacc atgtagactc ttggatacac aaggtaattt tagagccaca 600
ttaggatgaa ccttttaaaa agttatgcat ttatttttat gttccccac tggtgtatt 660
ataggacaat ttttatatgt gatatgtatt taccttagtg tgttaataa acactggcat 720
tccaagtgtg aaaaaa 736

```

&lt;210&gt; 46

&lt;211&gt; 1826

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2084989CB1

&lt;400&gt; 46

```

ccttagggcg cagggacagc cgagcggttac ctggtcccg gacagcgagc tctttaccca 60
ccccagttct ggttctgacg ccctagctca ttccgcaaat ttagggcttg ggtctggctt 120
gttccccctc ggctcgaacc acctcttctc tgagccgagc cagctaccgg ggctcctgga 180
attgccaccc ctccctgggc acccttgagg cctccgtgga gggacgtcac ggggcagagc 240
gggacgtgag cctgagtttg ctgcaggcgt gctctgtgtg gtggctgggt tctgccaatc 300
cccgtgccca ccgggtgggc gcggccggga agctcctgcc cctccctgct ggtcggcgctc 360
acgcgtgacg tcccgcgctg tggctgggag ggcccgccgg cgacagcgga ggcagagagg 420
aaggcggttc tgagagcttc agagagcgat ggaaagcaaa atgggtgaat tgcctttaga 480
catcaacatc caggaacctc gctgggacca aagtaacttc ctgggcagag cccggcactt 540
tttcaactgt actgacctc gaaatctgct gctgtccggg gcacagctgg aagcttctcg 600
gaacatcgtg cagaactaca gggccggcgt ggtgacccca gggatcaccc aggaccagct 660
gtggaggggc aagtatgtgt atgactccgc cttccatccg gacacagggg agaagtggt 720
cctgattggc cgcattgtcag cccagggtgcc catgaacatg accatcactg gctgcatgct 780
cacattctac aggaagaccc caaccgtggt gttctggcag tgggtgaatc agtccttcaa 840
tgccattggt aactactcca accgcagtgg tgacactccc atcactgtga ggcagctggg 900
gacagcttat gtgagtgcca cactgggagc gtggtggcac gccctgggac tcaaatccct 960
caccaagcac ctgccccctc tggctggcag atttgtgccc ttgacagcag tggcagctgc 1020
caactgcatc aacatcccc tgatgaggca gagagagctg caggtgggca tcccgggtggc 1080
tgatgaggca ggtcagaggc ttggctactc ggtgactgca gccaaagcagg gaatcttcca 1140
ggtggtgatt tcaagaatct gcatggcatg tcttgccatg gccatccac cactgatcat 1200
ggacactctg gagaagaaag acttcctgaa ggtaggcgac tgtacctctc ttgtcctgga 1260
atggggcgatg gctgggagaa gtgaccaggc cccaactctc tctccagcct cgcctgattc 1320
tctaagactt gccagccctt ctctgaccc ctgcaccgcc tctccacctc tcgttcatte 1380
agcaagaatg aactgggctg ggtgaaagga actctgcagg ggcaggagga gaggacaaa 1440

```

```

gaaggaaacc aacttcatca gtgttactcc agtggcttct gacacacaga aggggactgt 1500
catagtcatg cttgatctca tgctcattct tttacccctt agtgcctcca tactgagagg 1560
tacacacggg tgaacacgca cacacagaca tgaacaggac acgaaagcaa agcacaggaa 1620
caagctctgg ctcattcaca gaatcattta ttcacaaatg tattgagtgc catgcaccag 1680
gcatgtttta gggctgagga gatggcactg aacacaatgg ttatggcccc tgtcctcatg 1740
aagtttatag tctgatgcag aaaccaataa acaaggaggc acccacataa atacattctt 1800
agaaagtgtg aaaataaaaa aaaaaa 1826

```

&lt;210&gt; 47

&lt;211&gt; 1325

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2501034CB1

&lt;400&gt; 47

```

ccacgggtcc ggttctggac tgcagttgag tggaaatggg caacggcggg cggagcgggc 60
tgcagcaggg gaaggggaac gtggatgggg tggcagcgac tcctactget gcctcggcct 120
cctgccagta caggtgcatc gaatgcaacc aggaggccaa agagtgtgtac cgagactata 180
accacgggtg gctgaagata accatctgta aatcctgcca gaaacctgta gacaaatata 240
tcgagtatga tcctgttata atcttgatta atgctatatt gtgcaaagct caggcctaca 300
gacatattct tttcaatact caaataaata tccatggaaa actctgcata ttttgtttgc 360
tttgtgaagc atacctgagg tgggtggcagc ttcaagattc caaccagaat actgccccctg 420
atgacttgat cagatatgct aaggaatggg atttctatag aatgtttgcg attgctgctt 480
tagaacaacac tgcctatttt attggcattt ttaccttctt gtgggtagaa cggcccatga 540
cggcaaaaaa aaagcccaac ttcattttgc tgcctgaaagc attattatta tctagctacg 600
gaaaactctt gctgattcca gctgtcattt ggggaacatga ctacacatct gtgtgcctca 660
aactcattaa agtatttggt cttacatcaa attttcaggc aattagagtg accctaaaca 720
tcaaccgtaa gctctccttc ttggccgtgt tgagtggcct actgctggaa agcatcatgg 780
tctacttctt ccagagtatg gaatgggatg ttggaagtga ttatgccatc tttaaatctc 840
aggacttctg aagagtttta ttcttcttca ctatctgtgg catgaccagc tgtatctgaa 900
agagaaaaga catgaaatat aaaccaacct cctcatttct gttgagttaa atgaagcaaa 960
gattggaaac actttctgaa aaagaaagca atgataatag cgggtggatac ccacccccac 1020
aaatgcaccc aagagacaag ccatttacat acagatattc acagtcacac atagaaacac 1080
ccacatggac acaaggaatg ttgctgcaga gactgaatga catgcaacag gtgaaggttt 1140
atacgtttata cacaaggcca ggtaagcgct cataattcac acataataaa acatctaggt 1200
ttcattcctt tgacatgttt atatcttttt aattttaaag ttgttactgg cttaaaatat 1260
tttgtgttct tacaatagaa acgcttttaa taaagtcttt cagaataaac caaaaaaaaa 1320
aaaaa 1325

```

&lt;210&gt; 48

&lt;211&gt; 1832

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2745212CB1

&lt;400&gt; 48

```

tgggctgtcg ttggctggag cagcggctgc gcggtctcgc gtgctgtgag gtctgcgggc 60
gctggcaaat ccggcccagg atgtagagct ggcagtgctt gacggcgctt ctgacgcgga 120
gttgggtggg gtagagagta gggggcggtg gtcgggggtg gtgggagaag gaggaggcgg 180
cgaatcactt ataaatggcg ccgaagcagg accggaagcc taaattccag gagggtgagc 240
gagtgcctgt ctttcatggg cctcttcttt atgaagcaaa gtgtgtaaa gttgccataa 300
aggacaaaca agtgaataac ttcatacatt acagtgggtg gaataaaaaat tgggatgaat 360
gggttccgga gagcagagta ctcaaatacg tggacaccaa tttgcagaaa cagcgagaac 420
ttcaaaaagc caatcaggag cagtatgcag aggggaagat gagaggggct gccccaggaa 480
agaagacatc ttgtctgcaa cagaaaaatg ttgaagtga aacgaaaaag aacaaacaga 540
aaacacctgg aaatggagat ggtggcagta ccagtggagc ccctcagcct cctcggaaga 600
aaagggcccc ggtagatcct actgttgaaa atgaggaaac attcatgaac agagtgtgaag 660
ttaaagtaaa gattcctgaa gagctaaaac cgtggcttgt tgatgactgg gacttaatta 720
ccaggcaaaa acagctcttt tatcttcttg ccaagaagaa tgtggattcc attcttgagg 780
attatgcaaa ttacaagaaa tctcgtggaa acacagataa taaggagtat gcggttaaatg 840
aagttgtggc agggataaaa gaataactca acgtaatgtt gggtaaccag ctactctata 900
aatttgagag accacagtat gctgaaattc ttgcagatca tcccgatgca cccatgtccc 960

```

```

aggtgtatgg agcgccacat ctcctgagat tatttgtacg aattggagca atgttggcctt 1020
atacacctct ggatgagaag agccttgctt tattactcaa ttatcttcac gatttcctaa 1080
agtacctggc aaagaattct gcaactttgt tcagtggcag cgattatgaa gtggctcctc 1140
ctgagtacca tcggaaagct gtgtgagagg cactctcact cacttatgtt tggatctccg 1200
taaacacatt tttgttctta gtctatctct tgtacaaacg atgtgctttg aagatgttag 1260
tgtataacaa ttgatgtttg ttttctgttt gatttttaac agagaaaaaa taaaaggggg 1320
taatagctcc tttttctctc tttctttttt tttttcattt caaaattgct gccagtgttt 1380
tcaatgatgg acaacagagg gatatgctgt agagtgtttt attgcctagt tgacaaagct 1440
gcttttgaaat gctgggtggt ctattccttt gacactacgc acttttataa tacatgttaa 1500
tgctatatga caaaatgctc tgattcctag tgccaaaggt tcaattcagt gtatataact 1560
gaacacactc atccatttgt gcttttggtt ttttttatgg tgcttaaagt aaagagccca 1620
tcctttgcaa gtcattccat ttgttactta ggcattttat cttggctcaa attgttgaag 1680
aatgggtgct tgtttcatgg tttttgtatt tgtgtctaata gcacgtttta acatgataga 1740
cgcaatgcat tgtgtagcta gttttctgga aaagtcaatc ttttaggaat tgtttttcag 1800
atcttcaata aattttttct ttaattttca aa 1832

```

&lt;210&gt; 49

&lt;211&gt; 1211

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4833111CB1

&lt;400&gt; 49

```

gacagaccgg cctcaaacat ggcgggcgccc agcgcgcgag gacgtgatcc gcttctgctc 60
cggcttggtt tgtagccttg acgaggtctg agcgaccatg gaccggccgg gggttcgtggc 120
agcgctgggt gctgggtggg tagcaggtgt ttctgttgac ttgatattat ttccctctgga 180
taccattaaa accaggctgc agagtcccca aggatttagt aaggctgggt gttttcatgg 240
aatatatgct ggcggttcct ctgctgctat tggatccttt cctaattgctg ctgcattttt 300
tatcacctat gaatatgtga agtgggtttt gcatgctgat tcatcttcgt atttgacacc 360
tatgaaacat atgttggctg cctctgctgg agaagtgggt gcctgcctga ttcgagttcc 420
atctgaagtg gttaagcaga gggcacaggt atctgcttct acaagaacat ttcagatttt 480
ctctaacatc ttatatgaag aggggtatcca aggggttgat cgaggctata aaagcacagt 540
tttaagagag attccttttt ctttgggtcca gtttccctta tggggagtcct taaaagccct 600
ctgggtcctg aggcaggatc atgtgggtgga ttcttggcag tcagcagctc gtggagcttt 660
tgcaggtgga tttgcgctg cagtcaccac cctctagac gtggcaaaaga caagaattac 720
gctggcaaa gctgggtcca gcaactgctg tgggaatgtg ctctctgtcc tgcatgggg 780
ctggcggtca caggggctgg caggattatt tgcaggtgtc ttccctcgaa tggcagccat 840
cagtcgggga gggttcatct ttctgggggc ttatgaccga acgcacagct tgctgttgga 900
agttggcaga aagagtcctt gaagcagaga caagcctcac ctccacttct gtcaagagag 960
gggcctgcag tgcaaaccct ctccgctgga gcagctgtct gaactatagg cccagtgct 1020
gaagaccagt tgtgctaaga taccggcatg gagattgtgc catccgtggg ataggctggc 1080
tggtatgaag tcatggcct gtatgccaga gagctaagag aagaaaacgg ggtctgtggc 1140
ggtactctga acaatttctt cagaacctct taataaataa gtttggtaat gctgagaaaa 1200
aaaaaaaaa a 1211

```

&lt;210&gt; 50

&lt;211&gt; 1046

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 876677CB1

&lt;400&gt; 50

```

cccacgcgtc cgggaatgtc ttctactctc tcatactttc tcctctcccc tctcccaagc 60
acatctgagt tgctgcctgt tcttcacact tagctccaaa cccatgaaaa attgccaaagt 120
ataaaagctt ctcaagaatg agatggattc taggggtgtc tcacctgaga agcaagataa 180
agagaatttc gtgggtgtca acaataaacg gcttgggtgta tgtggctgga tctgtttttc 240
cctctctttc ctgttggtga tcattacctt ccccatctcc atatggatgt gcttgaagat 300
cattaaggag tatgaacgtg ctgttgattt ccgtctggga cgcattccaa ctgacaaaagc 360
caaggggcca gggttgatcc tggctcctgcc atgcatagat gtgtttgtca aagttgacct 420
ccgaacagtt acttgcaaca ttctccaca agagatcctc accagagact ccgtaactac 480
tcaggtagat ggagttgtct attacagaat ctatagtgct gtctcagcag tggctaattg 540
caacgatgtc catcaagcaa catttctgct ggtcctaaacc actctgagaa atgtcttagg 600

```

gacacagacc	ttgtcccaga	tcttagctgg	acgagaagag	atcgcccata	gcatccagac	660
tttacttgat	gatgccaccg	aactgtgggg	gatccgggtg	gcccgagtgg	aaatcaaaga	720
tggttcggatt	cccgtgcagt	tgccagagatc	catggcagcc	gaggctgagg	ccaccggga	780
agcgagagcc	aaggctcctt	cagctgaagg	agaaatgaat	gcttccaaat	ccctgaagtc	840
agcctccatg	gtgctggctg	agtctcccat	agctctccag	ctgcgctacc	tgccagacct	900
gagcacggta	gccaccgaga	agaattctac	gattgtgttt	cctctgcccc	tgaatatact	960
agagggcatt	ggtggcgta	gctatgataa	ccacaagaag	cttccaaata	aagcctgagg	1020
tcctcttgcg	gtagtcagct	attgca				1046

&lt;210&gt; 51

&lt;211&gt; 1660

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2326143CB1

&lt;400&gt; 51

gctcccctgc	ccaccccgcc	cccgtggcgg	agccccgggag	tcgagtggga	gtcggccggc	60
cggcgcgggc	agccgtgacc	cccgcggggg	acactgcagc	cggagcccgg	gagggggccgc	120
gccgccaccg	tctgaactag	gatgtcccga	catgaagggtg	tcagctgtga	tgcatgttta	180
aaaggaaatt	ttcgaggctg	cagatataag	tgtttaattt	gctacgatta	cgatcctttgt	240
gcatcttggt	atgaaagtgg	tgcaacaaca	acaaggcata	caactgacca	cccaatgcag	300
tgcatattaa	caagggtaga	ttttgattta	tactatgggtg	gggaagcttt	ctctgtagag	360
cagccacagt	cttttacttg	tccctattgt	ggaaaaatgg	gctatacgga	gacatctctt	420
caagaacatg	ttactttctga	acatgcagaa	acatcaacag	aagtgatttg	tccaatatgt	480
gcagcggttac	ctggaggcga	tcctaatacat	gtcacggatg	actttgcagc	tcattcttaca	540
cttgaacaca	gagcccctag	agattttagat	gaatcgagtg	gtgttcgaca	tgtacgtaga	600
atgtttcacc	ctggccgggg	attaggaggtc	cctcgtgctc	gtagatcaaa	catgcacttt	660
actagcagtt	ctactgggtg	actttctctt	tctcagagtt	catattctcc	aagcaatagg	720
gaagccatgg	atcctatagc	tgagctttta	tctcagttat	caggagtggag	acgttctgca	780
ggaggacagc	ttaattctct	tgccccttcc	gcttctcagt	tacaacaact	gcagatgcag	840
ctgcagctag	aacggcagca	tgcccaggca	gcacggcaac	aactggagac	cgcacgcaac	900
gcaacccggc	gtactaacac	aagcagtgctc	accactacaa	tcacacaatc	cacagcaaca	960
accaacatag	ctaatacaga	aagcagtcag	cagactctac	agaattccca	gtttctttta	1020
acaagggtga	atgatcctaa	aatgtctgaa	acggagcgcc	agtccatgga	aagcgcagct	1080
gcagaccgca	gcctgtttgt	ccaagagctc	cttctgtcca	ctttagtgcg	tgaagagagc	1140
tcatectcag	atgaggatga	tcgggggggag	atggcagatt	ttggtgctat	gggctgtgta	1200
gatattatgc	cttttagatgt	tgcttttagaa	aacctaaatt	taaaagagag	taataaagga	1260
aatgagcctc	caccacctcc	tctttgatga	catcccaatt	cgcagacaat	gtcctctgtg	1320
ctgtattttgc	caatgaaagt	ggacaacaac	tatcttgggt	ttgtttgggtg	attgttaattt	1380
caggtctgtc	actcttggtta	cattgtgtac	attcaaaagg	aagagagaaa	atataatatga	1440
taatcatttc	cacttaacta	atttttactt	ctagcaggta	aatgtaggta	gcagtgccag	1500
ggtgatctct	gcttctctga	ccttgacatg	caaaaggctc	tcctaatact	ccacattcaa	1560
actgaagagg	aaaattgaaa	tctctaataga	agctgctgtg	tgtatttatg	aataattaatg	1620
aataaaaaact	gcttggatgg	tttaccttaa	aaaaaaaaaa			1660

&lt;210&gt; 52

&lt;211&gt; 1110

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2786302CB1

&lt;400&gt; 52

cctttatttag	agaaagagtg	acctatctac	cccttacttg	gggaagactc	tctgcttcaa	60
cagtttcttc	tgtcttctgc	tcctggacat	ggcatgaaaa	cggaagtttg	agtattttcaa	120
ggataaagaa	taactctag	acaagagttg	ctttggggct	aaacctatct	cctgtattca	180
tttagatttg	gctttaaaag	cccatactat	tttattagct	cttctatgct	ttttaaaagt	240
ttttaaaaaa	attaatgtgg	gctttttta	tctttctcag	tgaggagggtg	agtttgaata	300
aacctttctt	ccacatgaga	agtattttac	aagttgcttg	tcaaatttta	aagaaaatga	360
tcaaatcttc	aagaaaatga	tcaaatcttc	tgtagacaaa	aatggacaa	atattcaca	420
ttgagtgtga	atgccatggt	gatgctaatt	ctgattattt	ttcttttctt	ttagccatac	480
cttcagagtg	ttattttctt	ttttgtcata	aggtgtctgg	aatgaagta	tggaatgaa	540
ataatgaata	aagaccagtg	tttcagaatc	tctccacgga	gtagagaaac	tcattcccaat	600

```

ccggaagagc ccgaagaaga agatgaagat gttcaagctg aaagagtcca agcagcaaat 660
gcactcactg ctccaaactt ggaggaggaa ccagtcataa ctgcaagctg tttacacaag 720
gaatattatg agacaaagaa aagtgtgttt tcaacaagaa agaagaaaaat agccatcaga 780
aatgttttct tttgtgttaa aaaagggtgaa gttttgggat tactaggaca caatggagct 840
ggtaaaagta cttccattaa aatgataact ggggtgcacaa agccaactgc aggagtgggtg 900
gtgttacaag gcagcagagc atcagtaagg caacagcatg acaacagcct caagtctctg 960
gggtactgcc ctcaggagaa ctcactgtgg cccaagctta ccatgaaaga gcacttggag 1020
ttgtatgcag ctgtgggaaag actggggcaa aaaagatgct gctctcagta tttcacgatt 1080
ggtgggaggt cttaagctcc aggaacaact

```

&lt;210&gt; 53

&lt;211&gt; 1120

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3735780CB1

&lt;400&gt; 53

```

gaaatccagt tatcaaaatt gactcaagaa gagagaacct aacagaacaa taacaatgga 60
agaaaattggg aacattatca caaagctatc atcctgccaa actccaggct cagatgtcac 120
aggttaaaaa aagtccttca tgaaaaagaa agatcttaag cagcatgatg gattcagaag 180
ctcatgaaaa gaggccacca atactaacat cttcaaaaca agatatatca cctcatatta 240
caaatgttgg tgagatgaag cattaactgt gtggctgctg tgcagccttc aacaatgtcg 300
caatcacatt tccattcag aaggtcctct ttcgacaaca gctgtatggc atcaaaaacc 360
gggatgcaat acttcagttg agaagggatg gatttcgaaa tttgtatcgt ggaatccttc 420
ccccattgat gcagaagaca actacgcttg cacttatgtt tggctctgtat gaggatttat 480
ctgccttctt ccacaagcat gtcagtgtct cagagtttgc aaccagtggc gtggcggcag 540
tgccttgagg gacaacagaa gcaattttca ctccactgga aagagtccag acattgcttc 600
aagaccacaa acatcatgac aaatttacca acacttacca ggctttcaag gcactgaaat 660
gtcatggaat tggagagtat tatcgaggct tgggtgccat tcttttcggg aatggactca 720
gcaatgtctt gtttttcggc cttcgaggct ccattaagga gcatctgcct accgcaacga 780
ctcacagtgc tcactctggc aatgatttta tctgtggagg tctattgggt gccatgttgg 840
gattcttggt ttttccaatt aatgttgtaa aaactcgcat acagtctcag attggtgggg 900
aatttcagtc tttccccaag gttttccaaa aaatctggct ggaacgggac agaaaactga 960
taaatctttt cagaggtgcc catctgaatt accatcggtc cctcatctct tggggcataa 1020
tcaatgcaac ttatgagttc ttgttaaagg ttatatgaaa aaaccatcag ttaagtgcc 1080
tttatcaact gaatagacct tctaagaaga aaaaaaaaaa

```

&lt;210&gt; 54

&lt;211&gt; 886

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 039026CB1

&lt;400&gt; 54

```

ggccgagggt cctgtccaga ccctgacctt ccctcccaag gctcaaccgt cccccaacaa 60
ccgccagcct tgtactgatg tcggatgcga gagcctgtgc ttaagtaaga atcaggcctt 120
attggagaca ttcaagcaaa gggttgacaa ctacttttcc agaacagaaa ggaaactcat 180
gcatcagaaa aggtgactaa taaaggtaac agaagaatat ggctgcacaa ataccagaat 240
ctgatcagat aaaacagttt aaggaatttc tggggaccta caataaactt acagagacct 300
gctttttgga ctgtgtttaa gacttcacaa caagagaagt aaaacctgaa gagaccacct 360
gttcagaaca ttgcttacag aaatatttaa aaatgacaca aagaatatcc atgagatttc 420
aggaatatca tattcagcag aatgaagccc tggcagccaa agcaggactc cttggccaac 480
cacgatagag aagtcctgat ggatgaactt ttgatgaaag attgccaca gctgctttat 540
tgaaaatgag gactcatctg atagaatccc ctgaaagcag tagccacctt gttcaacct 600
ctgtcatgac tgtttggaac atggaaaccg ctggagaaac aaaattgcta ttaccagga 660
ataatcacia tagaaggctt tattgttcag tgaaataata agatgcaaca tttgttgagg 720
ccttatgatt cagcagcttg gtcacttgat tagaaaaata aaccattgtt tcttcaattg 780
tgactgttaa ttttaaagca acttatgtgt tcgatcatgt atgagataga aaaattttta 840
ttactcaag taaaataaat ggaaatatca ctgaaaaaaa aaaaaa

```

&lt;210&gt; 55

&lt;211&gt; 2336

<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 260607CB1

<400> 55  
taatacgcctc actatagggga atttggccct cgagcagtaa ttcggcacga ggaccatctc 60  
tttaggatata atttttaaat tctttgaaac acataaccaa aatggtttga ttcactgact 120  
gactttgaag ctgcactctgc cagttacacc ccaaattggct ttaatcccct ctcgggtctg 180  
gttgccctttt gcagtttggg ttgtggactc agtcctctgt aggggtctgg ttaggagaga 240  
gccattttta aggacaggga gttttatagc ccttttctac ttccctcccc tcctccagt 300  
ccttatcaat cttttttcct ttttctgac cccctccttc tggaggcagt tgggagctat 360  
ccttgtttat gcctcactat tggcagaaaa gaccccatctt aaaaccaga gaacactgga 420  
gggggatgct ctagtgtgtt ctgtgtccat tttctctgt gccaaagaca gacagacaga 480  
ggctgagaga ggctgttcct gaatcaaagc aatagccagc ttctgacaca tacctggctg 540  
tctgaggagg aaggcctcct ggaaactggg agctaagggc gaggcccttc ccttcagagg 600  
ctcctggggg attaggggtgt ggtgtttgcc aagccaaggg gtaggaggcc gagaaattgg 660  
tctgtcggct cctggttgca ctttggggaa ctcttgctc cacaaggaga caggtagctc 720  
cctgttgtgg gactgctctg tccccctgcc ctactgcaga gatagcactg ccgagttccc 780  
ttcaggcctg gcagacgggc agtgaggagg ggcctcagtt agctctcaag ggtgccttcc 840  
cctcctccca acccagacat accctctgcc aaactgggaa ccagcagtgcc tagtaactac 900  
ctcacagagc ccagagggc ctgcttgagc cttcttgctc cacaggagaa gctgtgctc 960  
ctaggcaacc ccttctcccc acctctcatc aggggtgggg gttctccttt ctttccccctg 1020  
aagtgtttat ggggagatcc tagtggtctt gccattcaaa ccactcgact gtttgccctgt 1080  
ttcttgaaaa ccagtagaag ggaaacagca cagcctgtca cagtaattgc aggaagattg 1140  
aagaaaaatc ctcatcaatg ccaggggaca taaaagccat ttcccttcca aatactcgac 1200  
aathtagatg cagaacattt ctctgtattc agacttagag taacaccagc tgaaaactgc 1260  
agtttctttc ctttggatac ataaggcttc tctatcgggg tacgggacag ggaggaggcc 1320  
tcatgtctga agggggattt aggggagaga gccgcagccc tgaccctcgg tcctgtgcac 1380  
cgctttgggg cacagtctga tggcgccctt gctggcgccct tagtatggtt gactccggat 1440  
ggacaaaaga aaaaaaattt tttttcttga atgaaatagc aggaagctcc tcgggagcat 1500  
gtgttttgat taaccgtagt gatggatgct acgagtataa atggattaac tacctcaatc 1560  
cttacagtaa gattggaact aagggcaggg actcatgcat aagggtatga atccagcca 1620  
ggacaagtga gttgaggcct gtgccacaaa aggtttgtcc ttgggggaaca ggcaggcctg 1680  
ccaggatccc ccccatatcg attgggctgg gagggctggc cgtgagggtcc ccactttctg 1740  
ctttccttgc ccatgtgtca cccctttggc ctccagcttg tccctctctc actttctata 1800  
gctttgttgg accagatggt gaggaaagga atggcctctt ccttctctaga gggggtggc 1860  
tggagtgaaga cctggggcctt ggccctggaac ccaccacaca gccccaaagt caggaagcct 1920  
ggggaaacca gagctgagac ctcttcaaca ggtttctttt gagatcctac acctccattg 1980  
ggcccttttt cagtcttcaa tggggggcca gttggctcta gaaggagaag aggtgaagca 2040  
ggatcctttt ccctggggga gtctgagggc gcggtccttg gactcattca ggccgtcttt 2100  
gtagttgggg gagttccact gggcgatccc agccccctcc caccaccctt ctaatggacc 2160  
tcctcataga agccccattt cacttttggt ttatctacct cttagcaaaa caatagataa 2220  
attaggtaga ggcagctcca cttgcttagg ttaggggggg aaaaagattt ctttttccaa 2280  
aggaaaaaaa tattaccttg agaatacttt ccaaaaaata aaatttaaaa aaaaaa 2336

<210> 56  
<211> 2200  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1429651CB1

<400> 56  
gagaacttta cgcttgatc tcatctaact gacacagaaa ccctgtaagg atccagaggt 60  
ctcgttcagg accatggaga gcggcaccag cagccctcag cctccacagt tagatcccct 120  
ggatgcgttt ccccgagaag gcttgaggcc tggggacatc gcggtgctag ttctgtactt 180  
cctctttgtc ctggctgttg gactatggtc cacagtgaag accaaaagag acacagtga 240  
aggctacttc ctggctggag gggacatggt gtggtggcca gtgggtgcat ccttgtttgc 300  
cagcaatggt ggaagtggac atttcattgg cctggcaggg tcagggtgctg ctacgggcat 360  
ttctgtatca gcttatgaac ttaattggctt gttttctgtg ctgatgttgg cctggatctt 420  
cctaccatc tacattgctg ttcaggctcac cagatgcca gaatacctac ggaagcgtt 480  
cggtggcatc agaattccca tcatcctggc tgtactctac ctatttatct acatcttcac 540  
caagatctcg gtagacatgt atgcagggtgc catcttcatc cagcagctct tcacactgga 600

tctgtacctg	gccatagttg	ggctactggc	catcactgct	gtatacacgg	ttgctgggtg	660
cctggctgct	gtgatctaca	cggatgcccc	gcagacgctg	atcatgctta	taggagcgct	720
caccttgatg	ggctacagtt	tcgccgcggt	tgggtgggatg	gaaggactga	aggagaagta	780
cttcttggcc	ctggctagca	accggagtg	gaacagcagc	tgcgggctgc	cccgggaaga	840
tgcttccat	attttccgag	atccgctgac	atctgatctc	ccgtggcccg	gggtcctatt	900
tggaatgtcc	atcccatccc	tctgggtactg	gtgcacggat	caggtgattg	tccagcgga	960
tctggctgcc	aagaacctgt	cccatgccaa	aggaggtgct	ctgatggctg	catacctgaa	1020
ggtgtgccc	ctcttcataa	tgggtgtccc	tgggatggtc	agccgcatcc	tcttcccaga	1080
tcaagtggcc	tgtgcagatc	cagagatctg	ccagaagatc	tgcagcaacc	cctcaggctg	1140
ttcggacatc	gcgtatcccc	aactcgtgct	ggaactcctg	cccacagggc	tccgtgggct	1200
gatgatggct	gtgatgggtg	cggctctcat	gtcctccctc	acctccatct	ttaacagtgc	1260
cagcaccatc	ttcaccatgg	acctctggaa	tcacctccgg	cctcgggcat	ctgagaagga	1320
gctcatgatt	gtgggcagg	tgtttgtgct	gctgtctggtc	ctgggtctcca	tcctctggat	1380
ccctgtggtc	caggccagcc	agggcggcca	gctcttcatc	tatatccagt	ccatcagctc	1440
ctacctgcag	ccgcctgtgg	cgggtgtctt	catcatggga	tgtttctgga	agaggacca	1500
tgaagagggt	gccttctggg	gcctgatctc	gggcctgtct	ctgggcttgg	ttaggtggt	1560
cctggacttt	atttacgtgc	agcctcgatg	cgaccagcca	gatgagcgcc	cggctcctggt	1620
gaagagcatt	cactacctct	acttctccat	gatcctgtcc	acggtcaccc	tcactactgt	1680
ctccaccgtg	agctggttca	cagagccacc	ctccaaggag	atggctcagcc	acctgacctg	1740
gtttactcgt	cacgaccccg	tgttccagaa	ggaacaagca	ccaccagcag	ctcccttgtc	1800
tcttaccctc	tctcagaacg	ggatgccaga	ggccagcagc	agcagcagcg	tccagttcga	1860
gatggttcaa	gaaaacacgt	ctaaaaccca	cagctgtgac	atgaccccaa	agcagttcaa	1920
agtgtgtaag	gccatcctgt	ggctctgtgg	aatacaggag	aagggcaagg	aagagctccc	1980
ggccagagca	gaagccatca	tagtttccct	ggaagaaaac	cccttgggtga	agaccctcct	2040
ggacgtcaac	ctcattttct	gcgtgagctg	cgccatcttt	atctggggct	attttgcctta	2100
gtgtgggggtg	aaccaggggg	tccaaactct	gtttctcttc	agtgtccat	ttttttaatg	2160
aaagaaaaaa	taataaagct	tttgtttacc	aaaaaaaaaa			2200

&lt;210&gt; 57

&lt;211&gt; 2823

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2069971CB1

&lt;400&gt; 57

gaaagacata	cacacttcat	gtaatgctac	ctgcaagtct	ccctagaaaa	gcagtttttg	60
taggtgaaaa	caatgaagcc	aggtaatat	gcaaggaggc	tgtaatTTTA	gcagacctac	120
caacaacact	gatgtaggaa	gctcattatt	ttaatTTTctg	gagcctTTTA	atTTTtctt	180
tagaaagtgt	ataaataatt	gcagtgtctg	tttgtctcca	aaactgggca	gtgagttcaa	240
caacaacgac	aacaacagcc	gcagctcatc	ctggccgtca	tggagtttct	tgaagaaacg	300
tatcttgtga	atgataaagc	tgccaagatg	tatgctttca	cactagaaaag	aaggagctgc	360
aaatgaacac	ttcatagcaa	tgtggaactc	caacagaaac	cgggtgaataa	agatcagtgt	420
cccagagaga	gaccagagga	gctggagtca	ggaggcatgt	accactgcca	cagtggctcc	480
aagccacag	aaaagggggc	gaatgagtac	gcctatgcca	agtggaaact	ctgttctgct	540
tcagcaatat	gcttcatttt	catgattgca	gaggtcgtgg	gtgggcacat	tgtgggaggt	600
cttgtgtttg	tcacagatgc	tgcccacctc	ttaatTgacc	tgaccagttt	cctgtctcagt	660
ctcttctccc	tgtggttgtc	atcgaagcct	ccctctaagc	ggctgacatt	tggatggcac	720
cgagcagaga	tccttgggtc	cctgctctcc	atcctgtgca	tctgggtggt	gactggcgctg	780
ctagtgtacc	tggcatgtga	gcgcctgtctg	tatcctgatt	accagatcca	ggcgactgtg	840
atgatcatcg	tttccagctg	cgcagtggcg	gccaacattg	tactaactgt	ggttttgcac	900
cagagatgcc	ttggccacaa	tcacaaggaa	gtacaagcca	atgccagcgt	cagagctgtc	960
tttgtgcatg	cccttgga	tctattttcag	agtatcagtg	tgctaattag	tgcaattatt	1020
atctacttta	agccagagta	taaaatagcc	gacccaatct	gcacattcat	cttttccatc	1080
ctggtcttgg	ccagcaccat	cactatctta	aaggacttct	ccatcttact	catggaaggt	1140
gtgccaaaga	gcctgaatta	cagtgggtgtg	aaagagctta	tttttagcagt	cgacgggggtg	1200
ctgtctgtgc	acagcctgca	catctgtgtc	ctaacaatga	atcaagtaat	tctctcagct	1260
catgttgcta	cagcagccag	ccgggacagc	caagtgggtc	ggagagaaat	tgctaaagcc	1320
cttagcaaaa	gctttacgat	gcactcactc	accattcaga	tggaaatctcc	agttgaccag	1380
gaccccgact	gccttttctg	tgaagacccc	tgtgactagc	tcagtcacac	cgtcagtttc	1440
ccaaatttga	caggccacct	tcaaacatgc	tgtctatgag	tttctgcac	atagaaaaata	1500
aggaaccaaa	ggaagaaatt	catgtcatgg	tgcaatgcac	atTTTtatcta	tttatttagt	1560
tccattcacc	atgaaggaag	aggcactgag	atccatcaat	caattggatt	atatactgat	1620
cagtagctgt	gttcaattgc	aggaatgtgt	atatagatta	ttcttgagtg	gagccgaagt	1680
aacagctgtt	tgtaaactatc	ggcaatacca	aattcatctc	ccttccaata	atgcatcttg	1740
agaacacata	ggtaaatTTg	aactcaggaa	agtcttacta	gaaatcagtg	gaaggggacaa	1800



```
atagtcacaa aattttacca aaacattaga aacaaaaaat aaggagagcc aagtcaggaa 1860
taaaagtgc tctgtatgct aacgccacat tagaacttgg ttctctcacc aagctgtaat 1920
gtgatttttt tttctactct gaattggaaa tatgtatgaa tatacagaga agtgcttaca 1980
actaattttt atttacttgt cacatttttg caataaatcc ctcttatttc taaattctaa 2040
cttgtttatt tcaaaacttt atataatcac tgttcaaaag gaaatatttt cacctaccag 2100
agtgtttaa cactggcacc agccaaagaa tgtgggtgta gagaccagaga agtcttcaag 2160
aacagccgac aaaaacattc gagttgaccc caccaagttg ttgccacaga taatttagat 2220
atttacctgc aggaaggaat aaagcagatg caaccaattc attcagtcga cgagcatgat 2280
gtgagcactg ctttgtgcta gacattgggc ttagcattga aactataaag aggaatcaga 2340
cgcagcaagt gcttctgtgt tctggtagca actcaacact atctgtggag agtaaaactga 2400
agatgtgcag gccaacattc tggaaatcct atgtcactgg gtttgggttg gaaacctgga 2460
cttctgcatt tttaaaagtt acccagagat gcttctaaag atgagccata gtctagaaga 2520
ttgtcaacca caggagtcca ttgagtggga catcagata catcattgg cagctacaat 2580
agtatcatga attgcaatga tgtagtgggg tagtaaaagga aagcagtgga tattgcgga 2640
tgggcatggc cagtgatgtt tcacgtcatt gaggtgacag ctctgctgga ctttgaatta 2700
catatggagg ctctccagga agacgaagaa gagaaggaca ttctaggcaa aaagaagact 2760
aggcacaagg cacacttatg tttgtctgtt agcttttagt tgaaaaagca agatacaggg 2820
tcg
```

<210> 58  
<211> 1491  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2329339CB1

```
<400> 58
cgcctccctc cagctgcgag tgcggcctcg gctggcgggc gcaccaggcc acagttgtaa 60
gggatcttgt ggctgtcagg atggcagagg agcaggaggt caccagagtc tgcaagttgc 120
ctgcacagcc ctcacacca cactgcgtga acaacaccta ccgcagcgca cagcactccc 180
aggctctgct ccgagggctg ctggctctcc gggacagcgg aatcctcttc gatgttgtgc 240
tgggtggtgga gggcagacac atcgaggccc atcgatcct gctggctgctg tcctgcgatt 300
acttcagagg aatgtttgct gggggattga aggagatgga acaggaagag gtccctgatcc 360
acggtgtgtc ctacaatgct atgtgccaaa tcctacattt catatacacc tccgagctgg 420
agctcagcct gagcaatgta caagagacac tgggtggctgc ctgccagctt cagatcccag 480
aaattatcca tttctgctgt gatttcctca tgtcctgggt ggacgaagag aacattctcg 540
atgtctaccg gctggcagag ctgtttgact tgagccgctt gactgagcaa ctggacacct 600
atatcctcaa aaactttgtg gccttctctc ggactgacaa gtaccgccag ctccattgg 660
agaaggtcta ctccctctc agcagcaatc gcctggaggt ctctgcgag accgaggtat 720
atgagggggc ccttctctac cattatagcc tggagcaggt gcaggctgac cagatctcgc 780
tgcacgagcc cccaaagctc cttgagacag tgcggtttcc gctgatggaa gctgaggtcc 840
tgcagcggct gcatgacaag ctggacccca gccctttgag ggacacagtg gccagcggcc 900
tcattgtacca ccggaacgag agcctacagc ccagcctgca gagcccgcaa acggagctgc 960
ggtcggactt ccagtgcgtt gtgggcttcg ggggcattca ctccacgccg tccactgtcc 1020
tcagcgacca gccaaagtat ctaaacctct tactgggaga gtggaagcac ttcactgctc 1080
ccctggcccc ccgcatgtcc aaccagggga tgcggtgct caacaacttc gtatactga 1140
ttggagggga caacaatgtc caaggatttc gagcagagtc ccgatgctgg aggtatgacc 1200
cacggcacia ccgctggttc cagatccagt ccctgcagca ggagcagccg gacctgtccg 1260
tgtgtgttgt aggcaggtac atctacgctg tggcggggcg tgactaccac aatgacctga 1320
atgctgtgga gcgctacgac cctgccacca actcctgggc atacgtggcc ccactcaaga 1380
gggaggtgta tgcccacgca ggcgcgacgc tggaggggaa gatgtatatc acctgcggac 1440
gcaagcttat tccctttagt gagggttaat tttagcttgc actggccgctc g 1491
```

<210> 59  
<211> 986  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2540219CB1

```
<400> 59
ggacgccaac ctgcgtttga agtccagggc agtggctcct gcggacagcc agcataccag 60
gggccagtgct actgcattac aaccattgtg aggaatgagg gcctgtgccc tgtggctggc 120
gggcggcatg gcaggagcaa tttcttgggg gacagcgact cctatggatg tcgtgaaaag 180
```

tcgactccaa	gctgatgggg	tttattttaa	caaataataa	gggtgcctgg	actgtatctc	240
ccagagttac	cagaaggaaag	gtctttaaagt	gtttttcaga	ggcatcactg	tgaacgcggg	300
gcggggcttc	cccatgagtg	cggccatggt	ccttgggtac	gagctgtcgc	tgcaggctat	360
ccgcggggac	cacgcagtga	cgagcccata	agcgccagga	ggtgaacaca	ggatgactac	420
agtgttcccc	tgggcctcat	ctctgcatgt	gaagccctga	gagctgcaga	tgtttgccct	480
ttggacctcc	aagtggacat	caattagcaa	gcgtgggcta	ggatgggtga	gacactgacg	540
tggcccttct	gatgcctggg	atgcctcatg	agtcactgat	tcaagccctc	caagggtctg	600
atccccaatg	cccactctgc	taggctggca	tcaaagagct	ttccaagaaa	tgtttgggtc	660
agctgagaag	tcttgaccat	gagcaccagg	gagccagaaa	ccaccagag	aaacgttgct	720
tcactcctct	gtctgaggat	ggggaggggc	cagtgaagtc	tgggctcagc	cactccctcc	780
agtctcaagt	aacacgtccc	cgtgcctcca	gtctcctctc	agcaccgacc	aggtttttcc	840
ccgctcctgc	acccgtggat	cctgaggaca	gcggtagcgc	cttcctcacc	gcacgctgag	900
tccagtgcgt	gtcctcact	gtgcacttat	tagtgtctgt	tgagtgatta	aatcacatcc	960
tcaggctctgc	agcaataaaa	tgaag				986

<210> 60  
 <211> 4023  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2722462CB1

<400> 60	
gtccggggcg	gcgcgatg
gctctggggg	tgagagttcc
aggcagcaat	ctacctccag
atgccaaagg	gctggcgggc
tggccacggc	cctgctgctg
tccggcttgg	catctatgtc
ttgaactctg	catgaagtta
ccatggtcaa	gacctcggg
ggcagatgtc	ccatgtgcgg
ggtattgcgg	tggcgctccg
tggacatcct	cctgctgctg
tggtctcccc	taacccttca
tggttgcctc	tctgaccaca
accctgggtc	ctgctctctc
acctgcttct	ggctgtgggt
ctttgctact	gcacaagcga
ggaggcctgc	cggcatctcc
ggatgagtg	cagggagcgc
tgctcagcct	aaaggacttt
agaaaaacag	agagcactgg
gtattaatat	ccttgtgaag
tcaacggggg	ctggatcctc
agcacgtgcc	ctggagtta
agggttgcgg	cctgggccc
ccgtgacagt	gttcgccttc
atctcatcgt	ggctcctgcg
accgcaacgt	gctggacacc
ccctgctcat	cttttactac
tcttcccaa	ctgctgcaac
ccgtggggca	caggaccgtg
tcctcaacag	ctttgtgacc
tggaaaggcg	cacctctcag
ttgtgacct	ggtgggtgat
gaatgaacta	cagccgcaag
aggaaatctc	caaagaagag
cctcctcgga	tgtcaccagg
attccatgg	gtttctggga
accaggagga	gatccaggag
aactcagcag	cagtgcagcc
agaccgttac	ctagcccagc
tactctactg	cgtgttacgg
tttatataca	ggaagaaaaa
tgctcttctt	aactccagaa
ccatccacga	gcccagtttg
ccgcacacaa	ctgggagatg
acaacgacaa	gttcttcacc
cacacaatca	cctctcttac
ccctgtgcga	ggcccccgcc
ggccatcggt	gtgttggtac
cttcatccgg	cacaagcgga
ggccatcggt	gtgttggtac
cattttccgt	gtggactgtc
ccagtccctg	ccgcccttca
tgccatcctc	ggtttctact
ggagaacagc	atcgctcagc
gatgccctcc	tactcccga
cgagctgtat	ttcatcatga
tgagaaacgc	aagttcaagt
ccgcctgctc	atcagccaga
catgcgcttc	tacaagcccc
gaatcagaac	aacacacccc
tgctttgaag	tggaaaggcca
ggcgctcctc	atcttcaaa
catgtacttg	gtgggtggcag
aggtgggaac	ttcttctcca
tgggtggag	ctgttctctga
atggaacttg	tttgacttct
cctcaacatg	gagcccttct
gtttaagttg	aaggagcgct
gatggccagc	ctgggcccga
ggagtctctc	tgccgggacg
ctaccgctgg	cgcaaccaca
tctcaataat	tttgacaaca
caacaactgg	tacatcatca
ctacttcatg	acctttttaca
cctcgaggcc	ttcgtcttcc
tggtggcatc	acccttgaga
ctaccgggag	gcacgggggg
gatggagaga	taccagcaac
cgacctgagc	ctgaagatgt
ggagcaagag	cagcagcgac
aggcagccgc	cagcgctccc
cgtctcttct	acaatagtat
tgtgtacaca	tatgcacata
ggtttataac	caccttgccc
gggtgcggcc	accaggtctg

```

agctcttccct actgtggaag gctccagaag gcccttcaca aggagacccc tcacctggat 2640
ccagtcgact gcggggcttg cccctcatgt gggctggcct ccatcggccca cgtccaaagc 2700
tgtcactgct actgcttcag gctcacatcc ccccgacctg atggcggtgcc cgccccctct 2760
cctgcgggcc catgccacag gttctctgtg tttgcttttag ggacagaacc acttaggaaa 2820
gaaagaacte ccggtctcca ggggtggtatt tcagtgtctg tgataatgtc acgcaacacc 2880
tcttcgggga ccagtgccca ggatctaatg gaagcggaat tggggcaact gggcccctgt 2940
ggccagagct cagttagcca gtgcccggcg gccacagatt acactgacca atctcctccc 3000
ttggctctgc aagcctccca cccagccttc tctggcttaa cccttgttgg cgaaaactct 3060
tccacagtgg cctccttggg gacccagaac ccggaggaag gggcatgagg caggaagtgg 3120
ggccgatgtc tgcaaccagg accacttcgt ggaatgggct cttgacccaa tccctttttt 3180
ttgcgattta cccgttcaag caaaacaacg ttttgggtta ctaaggattg tgctaaagcc 3240
gataccaggt ccttcacacg tgtgcactag gaacaggagc gaacagcaca gagagacgct 3300
ccctgtggga cgcagcagcc ccgtggcccc ggcccagttc ccagccaccc tccctggctc 3360
tgctcacacc agagatttcc atagcaggag cgggttgggtg agaagtaggt tcagatgaac 3420
ctcagttaac gtccccacce ctccctccac catggtaccc tgtaggagcc ctgtatgaca 3480
tctgagcgtg gtggaggtag gaggggttgc agctgcagtg accctggcac agagtagcagg 3540
tcagtgcaga ggtcgctttg gttccgcttc cctggggcac agaacggaac acagcatagg 3600
ttctgcagca ggagccgcag tggcaggatg gaggggtgcga agggcaagga gtgcactgct 3660
gggcatttct ggccagcccc ctccctctgg ggccctctgg tgctgcttc ctgtgacttc agaaggcagg 3720
tggacagagc tccctctctg ccttgctctc ttccagcca cagaacgggc agggtagcac 3780
ccgacccagg gggagcagta cctggtcccc caccctctcc tcccaaccac ctccaaggcc 3840
aagctgggtc ccatagccag cacggcatgg tttccctctt cccctcttcc caggtcaggg 3900
gagttggaca agtagcaggt gtttgttttt aaagcacagc cctttgggaa agcaacacat 3960
tattgagact cactgtgatt cccccgggag tcagactggc tttgtctctt tctctctgga 4020
ggg 4023

```

&lt;210&gt; 61

&lt;211&gt; 2345

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2739264CB1

&lt;400&gt; 61

```

gaaaggaggc ggaactcggt gatctgactg gcggtttccc ggccggactg agaaggggag 60
cgcgctgcgc gtcgcaggag taacctactt ggtctcctgc tttcgcgaca tggccttcaa 120
ttttggggct ccctcgggca cctccgggtac cgtcgcagcc acccgggccc ccgcggtggg 180
gtttggaggga tttgggacaa catctacaac tgcaggttct gcattcagct tttctgcccc 240
aactaacaca ggcactactg gactcttttg tggtactcag aacaaagggt ttggatttgg 300
tactggtttt ggcacaacaa cgggaactag tactggttta ggtactgggt tgggactggg 360
actgggattt ggaggattta atacacagca gcagcagcaa actacattag gtggtctctt 420
cagtcagcct acacaagctc ctacccagtc caaccagctg ataatactg cgagtgcctc 480
ttctgctcca acgctgttgg gagatgagag agatgctatt ttggcaaat ggaatcaact 540
gcaggccttt tggggaacag gaaaagggtta tttcaacaat aatattccgc cagtggaaat 600
cacacaagaa aatccctttt gccgatttaa ggcagtaggt tatagttgca tgcccagtaa 660
taaagatgaa gatgggctag tggtttttagt tttcaacaaa aaagaaacag agattcgaag 720
ccaacaacaa cagttggtag aatcattgca taaagttttg ggaggaaacc agacccttac 780
tgtaaatgta gagggcacta aaacattgcc agatgatcag acagaagttg ttatttatgt 840
tgttgagcgt tcgccaatg gtacttcaag aagagttcca gctacaacgc tatatgcccc 900
ttttgaacaa gccaatataa aaacacaatt gcagcaactt ggtgtaacce tttctatgac 960
tagaacagaa ctttctcctg cacagatcaa acagctttta cagaatcctc ctgctgggtg 1020
tgatcctatt atctgggaac aggccaaaggt agataacccct gattctgaaa agttaattcc 1080
tgtaccaatg gtgggtttta aggaacttct ccgaagactg aaggttcaag atcagatgac 1140
taagcagcat caaaccagat tagatatcat atctgaagat attagtgagc tacaaaagaa 1200
tcaaactaca tctgtagcca aaattgcaca atacaagagg aaactcatgg atctttccca 1260
tagaacttta caggctctaa tcaaacagga aattcaaagg aagagtgggt atgccattca 1320
ggctgatgaa gagcagttgc gagttcagct ggatacgatt caggggtgaa taaatgcacc 1380
tactcagttc aagggccgac taaatgaatt gatgtctcaa atcaggatgc agaattcatt 1440
tggagcagtc agatctgaag aaagggtatta catagatgca gatctgttac gagaatcaa 1500
gcagcatttg aaacaacaac aggaaggcct tagccatttg attagcatca ttaaagacga 1560
tctagaagat ataaagctgg tcgaacatgg attgaatgaa accatccaca tcagaggtgg 1620
tgtcttttag tgacagttca caaacttgtg taaaggtttg tgaaatgcat cttcttactg 1680
catcagacct tctttaagaa tgaaccgac cacatggagg gaaaaagaaa acaattcttt 1740
cttggatttg ttttttgaga agtttactga caaattactg ttcatacaat ctgaaatagt 1800
cacctcacag ctcttcaaaag aaaacctttg aaagatttat atctaaaagc tgtatttact 1860
ttaaaagaag tgcataatta ccaaaattgt atgtactatt gtacattttt, acaacagcat 1920

```

```

tttcttaaac ataatctgtg tttaatgatt attgtccatt gagcctgtac tctgctttcc 1980
ataccaagta aatatgaaat aatctacttt gcacataaca gaacaaacta taattacttg 2040
gctgttgagg atttgtactt gagtataaat gtacaccagt ttttgtattt gtgaactcat 2100
ctgtgggagg agtaaagaaa atccaaaagc atttaaatgtt ttgtttttgt tctataaaga 2160
tatgaaaatg tattttttata ttattttact tattttggaat ttacagagca cacctaagca 2220
attaggatat aacaaaacta ctttaaccatt tttgcaacca ttttgttttt taagcctttt 2280
tattttctaaa aagatgaaaa cttataaata aattcttaat ttgtaattac ttttaaaaaa 2340
aaaaa 2345

```

<210> 62

<211> 2085

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 2758310CB1

<400> 62

```

cggagatgtg gcgaccgttt ctggcatcat tctgagactc ggcagttgct tctcactgct 60
gcggccgggc ctgtctgtgg gagctgcac ctcctcatct gcaggcgtg gaaaaccaga 120
cacgatcgga catgcatgtg gttctgcggc caaagcacgc cctttggttg tgaacttcat 180
gatacctgtg tgcagctctg tcatttccac tctgctctgc tgcacagaag gcagaagccc 240
tggcgtctcc ctgctgtgtt cttcaggaga aacgtcaggg gccttcctcc aaggttctcc 300
agccccacac ccctgtggag gaaggtgctc tccaccgcgg tagtgggggc gccctgctc 360
ctcggagccc gctatgtcat ggcagaggca cgggagaaga ggaggatgcg gctcgtggtg 420
gatggcatgg ggcgctttgg caggtctctg aaggtcggcc tgcagatctc cctggactac 480
tgggtggtgca ccaatgttgt ccttcgaggg tggaaagagcc caggctactt ggaggtgatg 540
tctgctgtgc accagcgggc ggctgatgcc ctgggtggcag gggccatcag caacgggggc 600
ctctacgtga agctgggcca ggggctgtgc tccttcaacc acctgcttcc ccccgagtat 660
accgggacct tgcgctgtct agaggacagg gccctcaagc ggggcttcca ggaggtggat 720
gagttgttcc ttgaggactt ccaggccctc ccccacgagc tcttccagga gtttgactac 780
cagccaattg ctgccgccag cctggcacag gtgcacagag ccaagctgca cgatggcacc 840
agcgtggctg tgaaggtgca gtacatcgac ctgcgggacc gctttgatgg ggacatccac 900
accctggagc tcctgctgcg gctcgttgag gtcatgcacc ccagctttgg cttcagctgg 960
gtcctccagg acctgaaggg gacctggcc caggagctgg acttcgagaa tgagggccgc 1020
aacgcagagc gctgtgcgcg ggagctggcg cacttcccct acgtcgtggt gccccgcgtg 1080
cactgggaca agtccagcaa gcgctgtctc actgcccact tctgcgccgg ctgcaaggtc 1140
aacgatgttg aggccatcag gagccagggg ctggcagtgc atgacatagc agaaaagctc 1200
atcaaggcct ttgctgagca gatattttac accggcttca tccactcgga cccacatcct 1260
ggcaacgttc tgggtgcggaa agggccggac gggaaagcgg agctggtgct gctggaccac 1320
gggctctacc agttcctgga ggagaaggac cgcgagccc tctgccagct gtggcggggc 1380
atcatcctgc gggacgacgc cgccatgagg cgcacgcag ccgcaactgg ggtgcaagac 1440
tacctcctgt tcgccgagat gctcatgcag cgcctcgctg gcctggggca gctgtggggc 1500
tcgcacctac tgagccgcga agaggcggcc tacatggtgg acatggcccg cgagcgcttc 1560
gaggccgtca tggcggtgct cagggagctg ccgcgcccca tgcgtctggt gctgcgcaac 1620
atcaacaccg tgcgctgtat caacgtggcc cctggcgccc cctgggaccg ctacttctt 1680
atggctaaaa gggtgttccg gggctggagc cgcctggcgg gcgccacgta tcggggtgtc 1740
tacggcacca gcctcctgcg ccacgccaa gtcgtctggt agatgctcaa gtttgaagt 1800
gcgctcaggc tggagacctt ggccatgcgg ctgaccgcc tcctggctcg tgctctggtc 1860
cacctgagcc tcgtgcccc agcggaggag ctctaccagt acctggagac ctagggtgca 1920
gccgcccagg gccggcgggg ccctttttac cttgggctga cggaggtggc ggggctagag 1980
gtgtagacac cccgagcccc gtgggcactc gcactggggg gctgtgacag cagctggggc 2040
aggaggcctg gtaatgacca cacactcctc tcaagcaaaa aaaaa 2085

```

<210> 63

<211> 3014

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 2762348CB1

<400> 63

```

ggaggcaaa ggcaggctga ggatcagggt ggcccgggtg gcagcgggga ggcgctgcat 60
gctggaggct gtgctgagtg cccggtgcag gtgagccggt cctgcggagt tgtgccgagt 120
gcctgctgca gaccgaggct gggccaagat ggcgtctgtg tttcgaagcg aggagatgtg 180

```

```

tttgtcacaa ctgtttctcc aggtggaagc tgcattattgc tgtgtggctg agctcggaga 240
gctcggattg gttcagttca aagattttaa tatgaatgtg aacagctttc aaaggaaatt 300
tgtgaatgaa gtcagaaggt gtgaatcact ggagagaatc ctccgttttc tggaagacga 360
gatgcaaaat gagattgtag ttcagttgct cgagaaaagc cactgaccc cgctcccacg 420
ggaaatgatt accctggaga ctgttctaga aaaactggaa ggagagttac aggaagccaa 480
ccagaaccag caggccttga aacaaagctt cctagaactg acagaactga aatacctcct 540
gaagaaaacc caagacttct ttgagacgga aaccaattta gctgatgatt tctttactga 600
ggacacttct ggctccttgg agttgaaagc agtgcctgca tatatgaccg gaaagtgtgg 660
gttcatagca ggtgtgata caacagggaa gaggatggct tcctttgagc ggttactgtg 720
gcgagtctgc cgaggaaacg tgtacttgaa gttcagttag atggacgcc ctctggagga 780
tcctgtgacg aaagaagaaa ttcagaagca catattcctc atattttacc aaggagagca 840
gctcaggcag aaaatcaaga agatctgtga tgggtttcga gccactgtct acccttgcct 900
agagcctcgc gtggagcgca gagagatgtt ggagagcgtc aatgtgaggc tggaagattt 960
aatcaccgtc ataacacaaa cagagtctca ccgcagcgc ctgctgcagg aagcctgc 1020
caactggcac tcctggctca tcaaggtgca gaagatgaaa gctgtctacc acatcctgaa 1080
catgtgcaac atcgacgtca cccagcagtg tgtcatcgcc gagatctggt tcccggtggc 1140
agatgcacac cgtatcaaga gggcactgga gcaaggcatg gaactaagtg gctcctccat 1200
ggccccatc atgaccacag tgcaatctaa aacagccctt cccacattta acaggaccaa 1260
taaatccaca gctggcttcc agaataattgt tgatgcctat ggtgtcggca gctaccggga 1320
gataaaacca gccccctaca ccatcatcac tttcccttct ctgttcgctg tgatgtttgg 1380
agactgtggt catggaaccg tgatgctcct ggctgcactt tggatgattc tgaatgagag 1440
acgcttgctc tccagaaga cagacaatga gatttggaac accttcttcc acgggcgcta 1500
tctgactcta ctatgggca tcttctccat ctacacgggt ttgatctaca atgactgctt 1560
ctccaagtcc ttgaacatct ttggctcttc tggagtgctc ccaacctatgt tcagaacagg 1620
cacatggaat actcatgtaa tggaggaaag tctatatctg cagctggacc cagccatacc 1680
aggagtgtat tttggaaatc cataccgtt tgggattgat ccgatttggg acttggcttc 1740
aaacaaactc acatttctga actcgtataa aatgaagatg tccgtgatcc tgggaattgt 1800
ccagatgggt tttgggtgca tccctcagct tttcaatcac atatacttca gaagaactct 1860
caacatcatt ctgcaattta tccctgagat gatttttatc ctgtgtctgt ttggatacct 1920
ggttttcatg atcattttca aatggtgctg ctttgacgtc cacgtatctc agcacgcccc 1980
cagcatcctc atccacttca tcaacatggt tctgtttaac tacagtact cttccaacgc 2040
acccctctac aaacatcagc aagaagtcca aagtttcttt gtggttatgg ctttgatttc 2100
tgtgccgtgg atgcttctga ttaagccgtt tattcttaga gccagtcac ggaaatccca 2160
gctgcaggca tccaggatcc aagaagatgc cactgagaac attgaagggt atagctccag 2220
cccttctagc cgttctggcc agaggacttc tgcagatacc cacggggctc tggacgacca 2280
tggagaagag ttcaactttg gagacgtctt gtccacca gcatccaca ccacgagta 2340
ctgcctgggc tgcatttcaa acacagcctc ttaacctgcy ctctggggcc tcagcctggc 2400
tcatgcacaa ctgtctgaag tgctctggac tatggtgatg aacagcgcc ttacagacgc 2460
aggctgggga ggaatcgtc ggggttttat tattttggc gtatttgcgt tcctgacagt 2520
agccatcctt ctgatcatgg agggcctctc tgetttctg cagccctgc gactgcactg 2580
ggttgagttc cagaacaagt tctatgtcgg ggatggttac aagttttctc catttctct 2640
taaacacatc ctggatggca cagccgagga gtaggctgag ggtgcacct cccacgggtg 2700
tcaccatgcc aatgaaggaa gttcagtcct gtctttgata tcagccctg caaggcgctc 2760
aatgggaagg ttgttcttgg ctcacctgaa gcatgaaact gtgtattatt tggagctcag 2820
cctgtggatt tgatacgact taaccacgtc agaggaagga ctttggcaag tgatattgtc 2880
ttcatgtggg gtattaattc tcaaataata aagtaattga caaatgagg gagaatgcta 2940
aacagatgtc ttcttgcaat attttaataa ttgtatttga gaaaataaac atctgagtca 3000
ttcaaaaaaa aaaa 3014

```

&lt;210&gt; 64

&lt;211&gt; 1726

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3715961CB1

&lt;400&gt; 64

```

tgaaaatgcc tgctgcgtac caaggtatgt actagggcat ctggggtaag taaaaacaaa 60
cacatagagc ctgcctggag aagctcatgg tctgatggaa agataagcaa gaagagttaa 120
tttctaatac atatgataaa aaggtcagag agcagtttct gaaaaacatg tttttgagtt 180
gagtcttgaa agacaaggag atgttagtaa agcagagaag ggagaattca tctagaagaa 240
atcagacagt gtgtgggaag ggcagagtct gaaaagagca tgccccattt ggagaagcat 300
caagaagccc acgtgttaga agcaccggcc ccatgagaca aagacacagc tagagagatt 360
gactaggcca tgtcggaaat tctcttatt ttatacatat ataagcatat agatacatat 420
agccaaagtt acctttttaa tgatcttttt taccagtggt attctggagg tcgaatggct 480
acatatgaac atctccgaga ggttgtgttt ggcaaaagtg aagatgagca ttatcccttc 540

```

```
tggaaatcag tcattggagg gatgatggct ggtgttattg gccagttttt agccaatcca 600
actgacctag tgaagggttca gatgcaaagt gaaggaaaaa ggaaactgga agggaaacca 660
ttgctgatttc gtggtgtaca tcatgcatat gcaaaaatct tagctgaagg aggaatacga 720
gggctttggg caggctgggt acccaatata caaagagcag cactggtgaa tatgggagat 780
ttaaccactt atgatacagt gaaacactac ttggtattga atacaccact tgaggacaat 840
atcatgactc acggtttatc aagtttatgt tctggactgg tagcttctat tctgggaaca 900
ccagccgatg tcatcaaaag cagaataatg aatcaaccac gagataaaca aggaagggga 960
cttttgtata aatcatcgac tgactgcttg attcaggctg ttcaagggtga aggattcatg 1020
agtctatata aaggcttttt accatccttg ctgagaatga ccccttggtc aatgggtgtc 1080
tggtctactt atgaaaaaat cagagagatg agtggagtca gtccatttta aaccctctaa 1140
gatgcaaccc ttaaagatac agtgttcagt attattgaaa tatgggcctc tgcaacacat 1200
accccttatt atttctacct ctttaggaag acacctattc cacagagact gatttatagg 1260
gggcagcact ttattttttt ctggaaaccc aagttctctt tgactcctct ttttgtccaa 1320
aagtgtctg gtcggatctc acaaggccat ccaatgagac cccgcacagc attttctaaa 1380
gaagaatcga agcctgacca ctttcacctt gggcaagaag gtttggcctt tgagttgcta 1440
ttctatgctg aagagcctgc tttagaggag agtaccagga gggagccagc atttcagatc 1500
tgaagtagac gataggaatg tggagaaca catacatagt gcttaagaaa tacatttaac 1560
ctgttatgtc agtatttatc aatgaagttt gataattcac tttctgtca ttgttaaagc 1620
gtacatactg taaattaaag ggaggtgaat ggaattaat gaataaacat tttgagtttc 1680
cctagtgttg aaggaaagtg tactttttct tgtcagaaag ataaaa 1726
```

<210> 65  
<211> 899  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5108194CB1

```
<400> 65
gcgcggcgga cgcgtccggg ccggtgaggg ggcgcggggg gcgcggggg ggcccaagcg 60
tcagcgcggc cgcctgtcgg gctgaactga ggaccgagtc tcctgccatt ccgagcagcg 120
ctggtatggg taatgggtgtg aaggaaggcc cggtgcgatt gcatgaggat gctgaggctg 180
tcctgtcctc gtccgtctca tcaaagcgtg accacaggca agtgcctcagc tccctgctgt 240
ctggggccct ggctgggtgcc cttgccaaaa cagcggtagc tccctggac cgaacaaaa 300
tcattctcca agtgtcttca aaaagatttt ctgccaaagg ggcttccgg gtccctctact 360
acacctacct caacgaggga tttctcagct tgtggcgcg ggactcggcc accatgggtg 420
gcgtgggtgcc ctacgcggcc atccagttca gcgcacacga ggagtacaag cgcattcttg 480
gcagctacta tggcttccgt ggagaagccc tgcccccttg gcctcgctc ttcgcggcg 540
cactggctgg aacgacagcc gcttcaactg cctacccctt ggacctggtc agagcgcgga 600
tggccgtaac ccgaaggaa atgtacagca acatctttca tgccttcac cgcattctga 660
gagaagaggg gctgaagact ctctaccatg gatttatgcc caccgtgctg ggggtgctg 720
cctacgctgg cctgagcttc ttacattatg agacgctcaa gagcttcac agagagtaca 780
cgggccgcaa gcttattccc tttagtggag gttaatTTTA gcttggcact ggccgctgct 840
ttacaacgct gtgactggga aaaccctggc gttacccaac ttaatcgctt tgcagcaca 899
```

<210> 66  
<211> 643  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5503122CB1

```
<400> 66
ctttaagctg tagctgtggg ttctgcagca attttgtttt tgccttgaaa gaggtgctct 60
ggattatcac acctccatgt atgacaattt gtacctgcat ggaattgaag actcggagcg 120
tggttcagcg gattcctaca caagcaggcc gtctgactcc gatgtctctt tggaagagga 180
ccgggaagca attcgacagg agagagaaca gcaagcagct atccagcttg agagagcaaa 240
gtccaaacct gtagcatttg ccgtgaagac aaatgtgagc tactgcggcg ccctggacga 300
ggatgtgcct gttccaagca cagctatctc ctttgatgct aaagactttc tacatattaa 360
agagaaatat aacaatgatt ggtggatagg aaggctgggtg aaagagggct gtgaaattgg 420
cttcattcca agtcactca gattggagaa catacggatc cagcaagaac aaaaaagagg 480
acgtttttcac ggaggggaaat caagtggaaa ttcttcttca agtcttgag aaatggatc 540
tgggacattc cgagcaactc ccacatcaac aggtgagggg tgtagttaaa ctctttttca 600
tacactgtat tcctttttaaa aatatttgaa cacacatgca agc 643
```

<210> 67  
<211> 2574  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5517972CB1

<400> 67  
gcgcctccca cgccggccgc cgccgacgtg atcgctcggg cgccgggggc gtgggtgggg 60  
gaaggggttg tgccgcgcga cggctctgcgt gctgtgccca ctcaaaaggt tccgggcgcg 120  
caggacggaa gacgcagtg cgcactcc cactgagatt gagagacgcg gcaaggaggc 180  
agcctgtgga ggaactgggt aggatttagg aacgcaccgt gcacatgctt ggtgggtcttg 240  
ttaagtggaa actgctgctt tagagtttgt ttggaaggctc cgggtgactc atcccaacat 300  
ttacatcctt aattgttaaa gcgctgcctc cgagcgcacg catcctgaga tcctgagcct 360  
ttgggttaaga ccgagctcta ttaagctgaa aagataaaaa ctctccagat gtcttcagat 420  
aatgtcgaag tttttatccc agtgtcacaa ggaaacacca atggcttccc cgcgacagct 480  
tccaatggac tgaaggcatt tactgaagga gctgtgttaa gttttcataa catctgctat 540  
cgagtaaaac tgaagagtgg ctttctacct tgctgaaaaa cagttgagaa agaaatatta 600  
tcgaatatca atgggatcat gaaacctggg ctcaacgcca tcctgggacc cacagggtgga 660  
ggcaaatctt cgttattaga tgtcttagct gcaaggaaaag atccaagtgg attatctgga 720  
gatgttctga taaatggagc accgcgacct gccaatctta aatgtaattc aggttacgtg 780  
gtacaagatg atgttgtgat gggcactctg acggtgagag aaaacttaca gttctcagca 840  
gctcttcggc ttgcaacaac tatgacgaat catgaaaaaa acgaacggat taacagggtc 900  
attcaagagt taggtctgga taaagtggca gactccaagg ttggaactca gtttatccgt 960  
gggtgtgtctg gaggagaaaag aaaaaggact agtataggaa tggagcttat cactgatcct 1020  
tccatcttgt tcttgatga gcctacaact ggcttagact caagcacagc aaatgctgtc 1080  
cttttgctcc tgaaggatgt gtctaagcag ggacgaacaa tcattcttctc cattcatcag 1140  
cctcgatatt ccattctcaa gttgtttgat agcctcacct tattggcctc aggaagactt 1200  
atgttccacg ggcctgctca ggaggccttg ggataacttg aatcagctgg ttatcactgt 1260  
gaggcctata ataaccctgc agacttcttc ttggacatca ttaatggaga ttccactgct 1320  
gtggcattaa acagagaaga agactttaaa gccacagaga tcatagagcc ttccaagcag 1380  
gataagccac tcatagaaaa attagcggag atttatgtca actcctcctt ctacaaagag 1440  
acaaaagctg aattacatca actttccggg ggtgagaaga agaagaagat cacagtcttc 1500  
aaggagatca gctacaccac ctcttcttgt catcaactca gatgggtttc caagcgttca 1560  
ttcaaaaact tgetgggtaa tccccaggcc tctatagctc agatcattgt cacagtcgta 1620  
ctgggactgg ttataggtgc catttacttt gggctaaaaa atgattctac tggaatccag 1680  
aacagagctg gggttctctt ctctcagcag accaaccagt gtttcagcag tgtttcagcc 1740  
gtggaactct ttgtggtaga gaagaagctc ttcatatcat aatacatcag cggatactac 1800  
agagtgtcat cttatttctt tggaaaactg ttatctgatt tattacccat gaggatgta 1860  
ccaagtatta tattacctg tatagtgtac ttcatgttag gattgaagcc aaaggcagat 1920  
gccttcttcg ttatgatgtt tacccttatg atgggtggctt attcagccag ttccatggca 1980  
ctggccatag cagcaggtca gagtgtggtt tctgtagcaa cacttctcat gaccatctgt 2040  
tttgtgttta tgatgatttt ttcaggtctg ttgggtcaatc tcacaaccat tgcattcttg 2100  
ctgtcatggc ttcagtactt cagcattcca cgatatggat ttacggcttt gcagcataat 2160  
gaatttttgg gacaaaactt ctgcccagga ctcaatgcaa caggaaacaa tccttgtaac 2220  
tatgcaacat gtactggcga agaataattt gttaaagcag gcacgatctc ctcaccctgg 2280  
ggcttggga agaatacagc ggcttgggt tgtatgattg ttattttcct cacaattgcc 2340  
tacctgaaat tgttatttct taaaaaatat ccttaaatc ccccttaatt cagtatgatt 2400  
tatcctcaca taaaaaagaa gcactttgat tgaagtattc aatcaagttt ttttgttgtt 2460  
ttctgttccc ttgccatcac actgttgcac agcagcaatt gttttaaaga gatacatctt 2520  
tagaaatcac aacaaactga attaaacatg aaagaaccca agaaaaaaa aaaa 2574

<210> 68  
<211> 1571  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5593114CB1

<400> 68  
ggcgcgcgga cagctagggt tcacggccac tggggcagag gagccgcgag aagatgtggg 60  
tttttggtta cgggtccctg atctggaagg tggatttccc ctatcaggac aagctggctg 120  
gatacatcac caactacagc aggcgcttct ggcaaggcag cacggaccac cgccgggtcc 180  
ccggcaagcc tgggaagagt gtgactcttg ttgaagatcc tgcgggatgt gtatgggtgt 240

```

ttgcttacag attgccagta ggaaaggaag aagaagtaaa agcatacctt gacttcagag 300
aaaaaggagg ctacagaacc acaacagtca ttttttatcc aaaagatccc acaacaaaac 360
cattcagtggt attgctatat attggaacat gtgataatcc tgattatctt ggtcctgcac 420
ctctggaaga cattgctgaa caaatTTTTa atgcagctgg tccaagtgga agaaatacag 480
aatatctttt tgaacttgca aattctatta ggaaccttgt gccagaagaa gcagatgagc 540
atcttttcgc tttggaaaaa ttagtaaaagg aacgtttaga agggaaaacag aacctcaatt 600
gcatataaatt tagtcttcag agaattaact tcagtgcaca atgacaatat gatttggaaa 660
tacgtttact taaagatctt atttttaatg tagtgaggat attattttaa cttttatttt 720
aactggaaat gtcccgaaac acatatttaa aatattggga tacagtgaaa gaaaaattca 780
aattttaata acataaagat ttcctaactt tatgttattg aacacttact cactagaagt 840
gagttcttta gaaaaataca gtgaaggact cagttcagtc ttgtttttat cagagtgata 900
atcatcctgt ttcacatccc aatactattt tgaaattcta aacaatttaa ccaaaattcc 960
aatataatata aggttatgcc ttcaatata tctatataca ttctgttaac atggtttaaa 1020
atacacaagc ttaaaataac atgcttagaa atacacaata atatgaacag tatttcagcc 1080
ttaattgtga atttccttgt tattcaagta ttaaatgaaa tcttttgagt ttttagccaa 1140
aaattggcat ttttaaaata cgaaaatttc cttggaatta taatgtactg tacctctctt 1200
tttttaaaata aaggcatttt actatatgga aaataactca ctaaagcata aattacatta 1260
tacaatcat gatcactaat gatgtagtct gtcattcact ttgtattaat cttataccaa 1320
aactgaaaaa gatgggctga tactacaaat taatggcaca tataatgaaa atttagtttt 1380
taaacacagc tttggaattc tttgtctgtc actatctcaa tttgtgtgtg tgtgtgtgta 1440
tatacataaa tatacatata aaattttttt tttctttgca gcctgcgtct ggccatccca 1500
caggctggaa agtgtaacct ctggcagaag ccaagaacag gcacctcttg gaattataat 1560
tttgttttgt t 1571

```

&lt;210&gt; 69

&lt;211&gt; 1549

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 044775CB1

&lt;400&gt; 69

```

cgcgcttagg caggcggtgg cgcggctgga gtgccgcggg gagggctgtg ccggttgctt 60
tctgcagcgg catctcggcc agctctcctc gcgctccccg gggcgctgtg cgtctccagt 120
ccgggaccga agccgcctgc cgtagcgggc ggccagatcc gcgtccccgc tcagcgcccg 180
gaggacatgc gggagagaga atgagccaga gggacacgct ggtgcatctg tttgccggag 240
gatgtgggtg tacagtggga gctattctga catgtccact ggaagttgta aaaacacgac 300
tgagtcacgc ttctgtgacg ctttatattt ctgaagttca gctgaacacc atggctggag 360
ccagtgtcaa ccgagtagtg tctcccgac ctcttcattg cctaaagggtg atcttggaaa 420
aagaaggggc tcgttccttg tttagaggac taggccccaa tttagtgggg gtagccctt 480
ccagagcaat atactttgct gcttattcaa actgcaagga aaagttgaat gatgtatttg 540
atcctgatgc taccgaagta catatgattt cagctgcaat ggcaggcatt tactgtacat 600
ttctcccgag aaaagagtga gatcgtgtca tctcatgctc cccatccgca ggtcacttcc 660
tgtagaataa tggactaact taaacctcgt tttactgcaa tcacagcaac caacccatt 720
tggcttataa agactcgggt acagcttgat gcaaggaaac gcggggaaaag gcgaatgggt 780
gcttttgaat gtgttcgtaa agtgtatcag acagatggac taaaaggatt ttataggggc 840
atgtctgctt catatgctgg tatatcagag actgttatcc attttgttat ttatgaaagt 900
ataaaacaaa aactactgga atataagact gcttctacaa tggaaaaatga tgaagagtct 960
gtgaaagaag catcagattt tgtgggaatg atgctagctg ctgccacctc aaaaacttgt 1020
gccacaacta tagcatatcc acatgaagtt gtaagaacaa gactacgtga agagggaaca 1080
aaatacagat ctttttttca gactctatct ttgcttgctc aagaagaagg ttatgggtct 1140
ctttatcgtg gtctgacaac tcacttagtg agacagattc caaacacagc cattatgatg 1200
gccacctatg aattggtggt ttacctactc aatggatagc agcacgagga ctgctgtact 1260
gcaaaaaaag aagacaaaaa gattacagtg gaccatggga tacagaagcc agcatggcag 1320
acagaagaaa aatagtttgg gaacatgtaa ctattctaag tggaaagttt gttgtaggaa 1380
ttatagtaat cacaccacat tacttggcct ttcggtaatg tgaaaaaaa aaaaaaacct 1440
cagagcctcc aaggaaatgc ctttagaagc actcctctct caaaattgct attttctcta 1500
ccatgtcccc cagacacagt tgggttttgt tgatttatgg cagtcttct 1549

```

&lt;210&gt; 70

&lt;211&gt; 2237

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature



&lt;223&gt; Incyte ID No: 116588CB1

&lt;400&gt; 70

```
gtcactgggtg ccacggggcc tcagcgcact tctgtcttaa gtcctctgggc ctcccttattt 60
tccccctttgc gatcgattcc agccacacct gtggatgttg ctagttactc cgcgtccgga 120
acgtgggaggt cgagggactg agctggggcga gttttgtggc actcctttgc tcttcagcag 180
ctatcttttgc tatgataatc ctgctgcccc tcagactcaa gttaaaccag atattgcaagt 240
gaataaccacg aaattcatgc tgctgtatgc ctgggtattct tggcccaatg tagttttgtg 300
tttcttttggg ggcttttttga tagaccgagt attttggaata cgatgggggca caatcatttt 360
tagctgctttt gtttgcattg gacagggttgt ttttgccttg ggtggaatat ttaatgcttt 420
ttggctgatg gaatttggaa gatttgtatt tgggattgggt ggcgagtcct tagcagttgc 480
ccagaataca tatgctgtga gctgggttaa aggcaaagaa ttaaaccttg tgtttggact 540
tcaacttagc atggctagaa ttggaagtac agtaaaccat aacctcatgg gatggctgta 600
ttctaagatt gaagctttgt taggttctgc tggtcacaca accctcggga tcacacttat 660
gattgggggtg gtaacgtgta ttctttcact aatctgtgcc ttggctcttg cctacttgga 720
tcagagagcga gagagaatcc ttcataaaga acaaggaaaa acaggtgaag ttattaaatt 780
aactgatgta aaggacttct ccttaccctt tggtggtata tttatcatct gtgtctgcta 840
ttatgttgcct gtgttccctt ttattggact tgggaaagtt ttctttacag agaaatttgg 900
atcttcttcc caggcagcaa gtgcaattaa cagtgttgta tatgtcatat cagctcccat 960
gtccccgggtg tttgggctcc tgggtggataa aacagggaag aacatcatct gggttctttg 1020
cgcagtagca gccactcttg tgtcccatat gatgctggcc tttacgatgt ggaacccttg 1080
gattgctatg tgtcttcttg gactctccta ctatttgctt gcctgtgcat tgtggccaat 1140
ggtggcattt gtagtctctg aacatcagct gggaaactgca tatggcttca tgcagttccat 1200
tcagaatctt ggggttgcca tcatttccat cattgctggg atgatactgg attctcgggg 1260
gtatttgttt ttggaagtgt tcttcattgc ctgtgtttct ttgtcacttt tatctgtggg 1320
cttactctat ttggtgaatc gtgcccaggg tgggaacctt aattattctg caagacaaag 1380
ggaagaaata aaattttccc atactgaatg agaagttaaa atgaatgtgt catgagaatg 1440
ggcttaacac atcgttgggt tgaaaacttc catttttaa aatttagagt ttagtcatta 1500
gaaaaaataa tggactggaa agttatatat atatccaaat atacctattt caaagtgtat 1560
ttgtgaggcc tgttttagcc tgtgtctttt gtattgtgtg ttgctaaaga attctacttt 1620
tagtaggcta atcaacaatg aaagggttag ggaacatcc ttgagataat aggtgaaatt 1680
caggaaagac agtgaaaaat ggaacatcc ttgagataat cttcattagg 1740
tatatatctt agggatacag ccttttcttt atcttatagc aggaaaaaaa aacttttgag 1800
ggaatatagaa gggctgcgtt acacaaaata aacaatggca ttgtcatagg ccttcccttt 1860
actagtaggg cataatgcta gggaatatgt ctactctgta gttatgtgga ttgccagca atgaccttt 1920
cacgaacaat agcttgcgct ctactctgta gttatgtgga ttgccagca atgaccttt 1980
tcaatttctt atttctgtgt tactgaggac cctaactact tagggatgta attttatagt 2040
ataaactttc tgtacagttt ttcttatagt ctaataagta aaaagtgtcc ttcaaattat 2100
gataattggc tatgtacatg gataaattaa aacctgcac acggagtaaa aaaaaaaa 2160
aaaaaaaaaa aaaaaatgag cggccgcaag cttattccct ttagtgaggg ttaattttag 2220
cttggcactg gccgtcg 2237
```

&lt;210&gt; 71

&lt;211&gt; 1114

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 875369CB1

&lt;400&gt; 71

```
tgggaacgga gcagccccgg gggccccctt gaggcggcga ggccgcgaag ggcgcggggc 60
tggagggtag gagagcgcg gaaagcgccc cagacgccac tcgcggcgga cgcgggccag 120
ttcccagggg tttggagccg gcccgcgcg ccatggctca cgtcggctcc cgcaagcgct 180
cgaggagtcg cagccgggtcc cggggacggg ggtcgaaaaa gagaaagaag aagagcagga 240
aagacacctc gaggaactgc tcggcctcca catccaagg tcgcaaggcc agcacggccc 300
ctggggcgga ggcctcacct tctccctgca tcacagagag aagcaagcag aaggcccgga 360
ggagaacaag atccagctcc tcctcctctt ctccagttc ttctagctcc tcttcttctt 420
cctcgctctc ctccctcttc tccagtgatg gccggaagaa gcgggggaag tacaaggaca 480
agaggaggaa gaagaagaag aagaggaaga agctgaagaa gaagggcaag gagaaggcgg 540
aagcacagca ggtggaggct ctgcggggcc cagtgggcac ccagtggcac cgatcagctg 600
gggaggaaga ggatggccca gtcctgacgg atgagcagaa gtcccgaatc caggccatga 660
agcccattgac caaggaggag tgggatgccc ggcagagcat catccgcaag gtggtggacc 720
ctgagacggg cgcaccagag cttattaagg gagatggcga ggtcctagag gaaatcgtaa 780
ccaaagaacg acacagagag atcaacaagc aagccacccg aggggactgc ctggccttcc 840
agatgcgagc tgggttgctt ccctgagggc ccccgctggc caaggcctgt ggacgacgct 900
ggcggccag cctgggcag tttcagggtg ccagtgggaa gcctgatggg tgctggtggc 960
```

```

ctttcccccg tggattgggc tctggcccag cccagtcctc tctcaggggc agggggtgga 1020
ggttgggggc accggcctgc ttggcacccc catctgaaag agcagcactt ctcagctatt 1080
aaaggccccc tggatagaca aaaaaaaaaa aaaa 1114

```

```

<210> 72
<211> 998
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1325518CB1

```

```

<400> 72
gactaaaaaa gccatgtatt ctttcgtttc tctctaaaag aagaaaaata taatttaaaa 60
atacattgcg tattttctaa aacaataaat ttatagtgtt aatattcata gggatcaatca 120
aatgaagct tctccttttg gcctgcattg tatgtgttgc ttttgcaagg aagagacggg 180
tccccttcat tgggtgaggat gacaatgacg atggtcaccc acttcatcca tctctgaata 240
ttccttatgg catacgggaat ttaccacctc ctctttatta tcgcccagtg aatacagtc 300
ccagttaccc tgggaatact tacactgaca cagggttacc ttcgtatccc tggattctaa 360
cttctcctgg attcccttat gtctatcaca tccgtggttt tcccttagct actcagttga 420
atgttcctcc tctccctcct aggggtttcc cgtttgtccc tcttcaagg ttttttcag 480
cagctgcagc acccgctgcc ccacctattg cagctgagcc tgctgcagct gcacctctta 540
cagccacacc tgtagcagct gagcctgctg caggggcccc tgttgcagct gagcctgctg 600
cagaggcacc tgttgagct gagcctgctg cagaggcacc tgttgcagct gagcctgctg 660
cagaggcacc tgttgagct gagcctgctg cagaggcacc ttcaccagct gagcctgctg 720
cagccaagcc tgctgcccc aacccctacc cttctcctc tcttgaacag gcaaatcagt 780
gaaattctct agaagagtac catgggttca tttctatact gatgcagaaa taagtgaat 840
ctacaaaagt tttctttctt ttccaaagac tatttcattc tgttgatttc agagtattca 900
tctcactaca ttgatttgtt tgtggtagtt tttccttgga cttaatttat attgaaaaaa 960
cattgataat taaataaata aaatagataa tttagaca 998

```

```

<210> 73
<211> 2348
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2060987CB1

```

```

<400> 73
gggccagccg gctcgcccg ggcccatggc agcagcggct actgcagccg agggggtccc 60
cagtcggggg cctcccggg aagtcaccca tctgaatgtg ggaggcaaga gattcagtac 120
ctctcgccag actctcacct ggatcccaga ctctctcttc tccagtcctc tgagcggacg 180
catctcgacg ctgaaagatg agaccggagc aatcttcctc gacagggacc ctacagtcct 240
cgccccatc ctcaacttcc tgccgaccaa agagttggat cccaggggtg tccacgggtc 300
cagcctcctc catgaagccc agttctatgg gctcactcct ctgggttcgtc gcctgcagct 360
tcgagaggag ttggatcgat cttcttgtgg aaacgtcctc tccaatggtt acctgccgcc 420
accagtgttc ccagtgaagc ggccgaaccg gcacagccta gtggggcctc agcagctagg 480
aggacggcca gcccctgtcc gacggagcaa cagatgccc cccaaccttg gcaatgcagg 540
gctgctgggc cgaatgctgg atgagaaaac cctccctca cctcaggac aacctgagga 600
gccggggatg gtgcgcctgg tgtgtggaca ccataattgg atcgctgtgg cctataccca 660
gtttctagtc tgctacaggt tgaaggaaag ctctggctgg cagctggtgt tttccagccc 720
ccgcctggac tggcccatcg aacgactggc gctcacagcc cgggtgcatg gtggggcttt 780
gggtgaacat gacaagatgg tggcagcagc caccggcagc gagatcctgc tatgggctct 840
gcaggcggaa ggccgtggct ccgagatagg ggtctttcat ctgggggtgc ctgtggaggc 900
cttgttcttc gtcgggaacc agctcattgc tacaagccac acagggcgca tcggggtgtg 960
gaatgccgtc accaagcact ggcaggtcca ggaggtgcag cccatcacca gttatgacgc 1020
ggcaggctcc ttctcctcc tgggctgcaa caacggctcc atttactacg tggatgtgca 1080
gaagttcccc ttgcgcatga aagacaacga cctccttgtc agcagctct atcgggaccc 1140
agcggagatg ggggtcaccc cctcagtggt ctacctacc cccaagacca gtgacagtgg 1200
gaactggatc gagatcgct atggcaccag ctcagggggc gtgcgggtca tcgtgcagca 1260
cccgagact gtgggctcgg ggcctcagct cttccagacc ttcactgtgc accgcagccc 1320
tgtcaccaag atcatgctgt cggagaagca cctcatctca gtctgtgccg acaacaacca 1380
cgtgcggaca tggctgtgta ctgccttccg cgcatgatt tccaccagc cggctccac 1440
cccactcgct tctttaaga tcttggtctt ggagtcggca gatgggcatg gcggctgcag 1500
tgctggcaat gacattggcc cctacgggtg gcgggacgac cagcaagtgt tcatccagaa 1560

```

```

ggtggtgccc agtgccagcc agctcttctg gctctctca tctactgggc agcgggtgtg 1620
ctccgtgcgc tccgtggacg gctcacccac gacggccttc acagtgcctgg agtgccaggg 1680
ctcccgccgg ctcggctctc ggccccggcg ctacctgctc actggccagg ccaacggcag 1740
cttgcccatg tgggacctaa ccaccgccat ggacggcctc ggccaggccc ctgcaggttg 1800
cctgacggag caagagctga tggaaacagct ggaacactgt gagctggccc cgccgggtcc 1860
ttcagctccc tcatggggct gtctccccag cccctcacc cgcactctcc tcaccagcct 1920
ccactcagcc tccagcaaca cctccttgct tggccaccgt gggagcccaa gcccccgca 1980
ggctgaggcc cggcgcctgt gtggggggcag ctttgtggaa cgctgccagg aactgggtgcg 2040
gagtgggcca gacctccgac ggccacccac accagccccg tggccctcca gcggtctcgg 2100
cactcccttc acacctccca agatgaaact caatgaaact tccttttgaa caacgcagct 2160
gccatgatgc cttgggatgc cctggctctg ggggactcag gtgcctccct gattcctgtg 2220
ggaaccccggt gttcaggggc agggcctcct tggataaaat ggttattgtt actagggtccc 2280
caccttccct cttttctgga agccaaagtc accctcccca ataaagtcct cactgccaaa 2340
aaaaaaaaa 2348

```

<210> 74  
 <211> 1139  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2172064CB1

```

<400> 74
ctcagagctgg gatgtgtggc aggttctctg ggcggctgct ggcggaggag agccggcgct 60
ccacccccgt ggggcgcctc ttgcttcccc tgctcctggg attccgcctt gtgctgctgg 120
ctgccagtgg gcctggagtc tatggtgatg agcagagtga attcgtgtgt cacacccagc 180
agccgggctg caaggctgcc tgcttcgatg ccttccaccc cctctccccg ctgcgttcc 240
gggtcttcca ggtcatcttg gtgctgtac ccagcgccct ctatatgggt ttactctgt 300
atcacgtgat ctggcactgg gaattatcag gaaaggggaa ggaggaggag accctgatcc 360
agggacggga gggcaacaca gatgtcccag gggctggaag cctcaggctg ctctgggctt 420
atgtggctca gctgggggct cggcttgctc tggagggggc agccctgggg ttgcagtacc 480
acctgtatgg gttccagatg cccagctcct ttgcatgtcg ccgagaacct tgccttggtg 540
gtataacctg caatctgtcc cgccccctct agaagaccat ttccctaaag accatgtttg 600
gagtcagcgg ttctgtctc ttgtttactt ttttgagct tgtgcttctg ggtttgggga 660
gatggtggag gacctggaag cacaaatctt cactcttctaa atacttccct acttcagaga 720
gcaccagaag acacaagaaa gcaaccgata gcctccaggt ggtggaaacc aaagagcaat 780
ttcaagaagc agttccagga agaagcttag cccaggaaaa acaaagacca gttggacca 840
gagatgcctg agttggagat gaactttggc caactttcct catcaccata cttaaaatcc 900
tgtccaagga ggagcttatt caccattttc tatacatgtg acatatgtag cagcataacc 960
gacaactggg actgcgctgc cttgactcca cctctacata caatgactca gctaaccaga 1020
ctaataaaaag ccatgtttgc accattgtct agggaggcat tgctttgggg aattattccc 1080
agtgtcctcc ttacttatcg caagtaataa aaatcccctg ggaaatcttc aaaaaaaaaa 1139

```

<210> 75  
 <211> 863  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2219267CB1

```

<400> 75
tcagagctag cttccgccgc agcctggcac ctcaggtgcc gaccttctac ctgccctact 60
tcctggaggc cggcctggag gcggcagcct tgggtcttcc gctcctgacg gcagattgct 120
gtggacgccg ccccgctgct ctgctgggca ccatggctac aggcctggca tcctctgtgc 180
tcctcgctgg ggcccagtat ctgccaggct ggactgtgct gttcctctct gtctggggc 240
tcctggcctc ccgggctgtg tccgcactca gcagcctct cgcggccgag gtcttccca 300
cggtgatcag gggggccggg ctgggcctgg tgctgggggc cgggttccct ggccaggcag 360
ccggccccc cggacacctg cagggccggc agggcttctt cctgcaacaa gtcgtcttcg 420
cctcccttgc tgtccttgcc ctgctgtgtg tctctgtgct gcctgagagc cgaagccggg 480
ggctgcccc gtcactgcag gacgccgacc gcctgcgcgg ctcccactc ctgcggggc 540
gcccccgcca ggaccacct cctctgctgc cgccctccaa ctctactgg gccggccaca 600
cccccgagca gcactagtcc tgctgtgtgg ccctgggagc caggatggga ccaaagtcaa 660
ggcctggggc atggctgagt accccagacg tctggctcag ggcagacaca ttctctcag 720
aagcccgtgt ctcagtgcag gtggagccgt ggggacagcg tgaagggtgc tccagccagg 780

```

ccccaggcac tgggaggccc tgggtctccc cccagccaca cccagtaggt gtggaggata 840  
aaggcttctg tggaaaaaaa aaa 863

<210> 76  
<211> 1322  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2308629CB1

<400> 76  
ccggggggccc ggcgcgcggg gagggccgggg cctgcaggcc cccgggtacga caagatccgg 60  
actccggggccc ggactacgag gcgctgcccgg ctggagccac tgtcaccacg cacatggtgg 120  
caggcgcccggt ggaggggagc ctggagcact gcgtgatgta ccccatcgac tgcgtcaaga 180  
cccggatgca gagtctacag cctgacccag ctgcccgcta tcgcaatgtg ttggaggccc 240  
tctggaggat tataagaacg gagggcctat ggaggcccat gagggggctg aacgtcacag 300  
caacaggcgc agggcctgcc cacgcccttt attttgcctg ctacgaaaag ttaaaaaaga 360  
cattgagtga tgtaatccac cctgggggca atagccatat tgccaatggt gcggccgggt 420  
gtgtggcaac attacttcat gatgcagcca tgaaccctgc ggaagtggtc aagcagagga 480  
tgcagatgta caactcacca taccaccggg tgacagactg tgtacgggca gtgtggcaaa 540  
atgaaggggc cggggccttt taccgcagct acaccaccca gctgaccatg aacgttcctt 600  
tccaagccat tcacttcatg acctatgaat tccctgcagga gcactttaac cccagagac 660  
ggtacaaccc aagctccac gtccctctctg gagcttgccg aggagctgta gctgccgag 720  
ccacaacccc actggacgtt tgcaaaacac tgcctcaacac ccaggagtcc ttggctttga 780  
actcacacat tacaggacat atcacaggca tggctagtgc cttcaggacg gtatatcaag 840  
taggtgggggt gaccgcctat ttccgagggg tgcaggccag agtaatttac cagatccccct 900  
ccacagccat cgcattggtc gtgtatgagt tcttcaaata cctaactact aaaaggcaag 960  
aagagtggag ggctggcaag tgaagttaga ctgaacgaag ccagggggttc agatgacact 1020  
gctgcatect ggtcacattc tctgtctcct ggaatgctcc cactcaagt ggagttagaa 1080  
ggaaggtaga ggggctctcc cccaggattt tgggtgtttg actaacacca gttcctgcca 1140  
acctctgttg ccaccacctt tccctccagg ccctaagcac gtgcagcaaa gcacaccaca 1200  
gcacctttga taacctctct ccatcctggg cctgatgacc tgctctagac tgttatagag 1260  
ggataagcag ctcatctccc tggttcctaa taaaagcct ttaaattaaa aaaaaaaaaa 1320  
aa 1322

<210> 77  
<211> 1869  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2660038CB1

<400> 77  
caggagcgt cattcctggc cagagtccctg acttctttct cggatcgaga tctcgttgct 60  
ggctcgagga aatcacccgc tcttctcccg gatccttttc ttctctcact tgctggcttt 120  
cttcttctctg cctccctggc ttccatctcc caaccccggg attccctcct ctactcccgt 180  
gtctcccgctg cccgccatcc tgagccatcc cacttgcaac cttctgtctt ttgcccctcc 240  
ttgacctcag agggctcctgc cttaagcttc tcaccagaat ctccctagatt tctatctctt 300  
ccctgcttgc cagctcttga ctcccaaatt ccagctgacg ttgaccact tgacatttga 360  
ccctgacacc cttgactgca aatctaaatt cttatcttct gcaacctgta ctgctgaaac 420  
aggccttccc cctgtcttcc aaccccgtt tctgacaccc atttttactt tcttactctt 480  
gggtcagttc ctgctacagc tatcccacca tggacttctt gatgagtggc ctggcagcct 540  
gcggggcctg tgtattcacc aatcccctgg aggtggtgaa gaccaggatg cagttgcaag 600  
gagaactgca ggcccctggc acataccagc ggcactaccg aaatgtcttc catgccttca 660  
tcaccatcgg caaggtggat ggccttgctg cccctgcagaa aggcctggcc cccgccctct 720  
tgtaccagtt cctgatgaat ggcacccgac tgggcaaccta tgggctggct gaggtgggg 780  
gtacctgca cacagccgaa gcgaccaca gtccctgccg cagcgagca gctggggcca 840  
tggctgggggt cactgggagc tacttgggga gcccactcta catggtgaag acacacctgc 900  
aggcacaggc agcctcagaa attgctgtag ggcaccagta taagcatcag gccatgtttc 960  
aggcgctaac cgagattggc cagaaacatg gtctggtggg gttatggcgt ggggctcttg 1020  
gcggcctgcc ccgagttatc gtcggttctt ccaccagct gtgcaccttc tcatccacca 1080  
aggacctcct gagccagtgg gagatcttct cctcccagag ctggaagttg gcgctggtgg 1140  
ctgccatgat gagtggcatt gcagttgtct tggccatggc accctttgat gtggcctgca 1200  
caaggctcta caaccagccc acagatgcac agggcaaggg cctcatgtac cgggggatac 1260

```
tggacgctct gctgcagaca gctcggaccg agggcatttt tggcatgtac aaggggtatag 1320
gtgcctccta cttccgcctc ggccccaca ccatcctctc cctcttcttc tgggaccagc 1380
tgcgctccct ctactacaca gacactaaat aacagccgct tccccagctt ccaccaaattg 1440
agcactcctt ggccacttgt gcctccacca ctatgtcctg gtgactactg attaggtgac 1500
ctttcatcca tccatggggg acagccaacc ccactcccca tctgttctca gggttgaatc 1560
actacaagag atgagtttcc cttctttcct tgggtgttgc tttaaacctt ccctacccat 1620
tccctgggta actcacacc cttctctcagg gctgaacgag tcatcccaaa gtgtatttcc 1680
tcccactcac cactgccacc cttgagtcct tctgtctccc atgcacagt ttaaactcct 1740
ccctccaaaa ccaaagggaa tcgagagacc caattcccag gcgtctggga ccagggtgtc 1800
ctgttagatt caaaggcaca gagattatat tgattataaa gcaagtttat tctgaaaaaa 1860
aaaaaaaaa 1869
```

<210> 78

<211> 1881

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 2670745CB1

<400> 78

```
gaagaaccga gcttggctgt gtttatctcg ttggggacta aggcgtcggg tggcgcgcaa 60
cgggttctag gctgcaggca gctcgaggac ccgcggcccc gccccggctc ggcctggcag 120
atagcagagg cagcaggccg tgccgggggg gcatgttgct gtaaccagtg gccaggggga 180
tgttacgggtg gacagtgcac ctggagggcg ggccccgcag ggtgaaccat gctgcagtgg 240
ctgtcgggca tcgggtatac tccttcgggg gttactgtct tgggtgaagac tatgagacac 300
tgcgtcagat agatgtgcac attttcaatg cagtgtcctt gcgttggaac aagctgcccc 360
cgggtgaagtc tgccatccgt gggcaagctc ctgtggtacc ctacatgcgc tatggacact 420
caaccgtcct catcgacgac acagtcctcc tttggggcgg gcggaatgac accgaagggg 480
cctgcaatgt gctctatgcc tttgacgtca atacgcacaa gtgggttcaca ccccgagtgt 540
cagggacagt tcctggggcc cgggatggac attcagcctg tgtcctaggc aagatcatgt 600
acatttttgg gggctacgag cagcaggccg actgttttct caatgacatt cacaagctag 660
ataccagcac catgacatgg actcttatct gtacaaaggg cagccctgca cgctggaggg 720
acttccactc agccacaatg ctgggaagtc acatgtatgt ctttgggggc cgtgccgacc 780
gctttggggc attccattcc aacaatgaga tttactgcaa ccgcattcga gtctttgaca 840
ccagaactga ggcttggctg gactgtcccc cgactccagt gctgcctgag gggcgccgga 900
gccactcggc ctttggctac aatggggagc tgtacatctt tgggtggttat aatgcaaggc 960
tgaaccggca cttccatgac ctctggaagt ttaatcctgt gtcctttacc tggaaaaaga 1020
ttgaaccgaa ggggaagggg ccatgtcccc gccggcgcca gtgctgctgt attgttgggtg 1080
acaagattgt cctctttggg ggtaccagtc catctcctga ggaaggcctg ggagatgaat 1140
ttgaccttat agatcattct gacttacaca ttttggactt tagccctagt ctgaagactc 1200
tgtgcaaaact ggccgtgatt cagtataacc tagaccagtc ctgtttgcct catgatata 1260
ggtgggagct gaatgccatg accaccaaca gcaatatcag tcgccccatc gtctcctccc 1320
atgggtagga ggaagtctct gccacctccc cctctgagcc tgctgtcatc ttactgtccc 1380
ctgcccatct gtcacccacc tgctcctttg accctggac ttggtatacc tccatgtgga 1440
gttgttgggc gagaggtgtt ctctgtgctg tgaattcagt ggggagctgt agcggggtgg 1500
gggctagggt cctccccctt tgggcccagg gcccttccc cttggtgtct tgtccccatc 1560
cacctccttt cagctgtctc tgggcctcag ctctgcccag ggccagccag gttctgctgg 1620
gaagggaagg gaatggggag aagggagaag ctgagcctgt ctgagcctca ggagcttccc 1680
cctccccctt tgcttatccc ctccccctct cttgagcctt gagccttgac tgggagctga 1740
aaggagtgtc agctgttggc atgagacctc cttctccccg tcttggggag gtggggacca 1800
gcagataaat ccacacctc cttgagctgt cgctgtactc tgaagttcag ccagctcaga 1860
ttttataaaa attaatataa a 1881
```

<210> 79

<211> 2004

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 2676443CB1

<400> 79

```
aatcgcaggg ggcgtggcct agcgggtccc cccccggggc gcgcgcgcgc attggctgtg 60
cggggtgcgg gcgcgcgggc ggcgctttga accgggcgcg gggcgcgggg cgcggggcgc 120
tgcgcccggt acacgccggg gtagggccgg ggtcgggttg tggtcgggcc gggattgggc 180
```

tctcctgggc	catggcagcc	gagggcgcg	tgctcgctg	gtacttcggg	gggctggcct	240
cctgcggggc	cgcttgcctg	acgcacccgc	tggacctgct	caaggtgcat	ctgcagacgc	300
agcaggaggt	gaagctgcgc	atgacgggca	tggcgctgcg	ggtggtgctg	accgacggca	360
tcttggcact	ctacagcgcc	ctgagcgcc	cgctgtgcag	acagatgacc	tactccctga	420
ctcggttcgc	catctacgag	actgtgcggg	accgcgtggc	caagggcagc	caggggcctc	480
tccccctcca	cgagaaggtg	ttgctgggct	ccgtcagcgg	tttagctgga	ggcttcgtgg	540
ggacggccgc	agacttgggtc	aacgtcagga	tgcagaacga	cgtgaagctg	ccccagggtc	600
agcggcgcaa	ctacgcccc	gcgttggatg	gcctgtaccg	cgtagctcgt	gaagagggtc	660
tcaggagact	gttctcggtg	gcaacctggg	catccagccg	aggggcctta	gtcactgtgg	720
gccagctgtc	ctgctacgac	caggccaagc	agctgggtcct	tagcacgggg	tacctctctg	780
acaacatctt	cactcacttt	gtcgccagct	ttattgcagg	tggatgtgcc	acgttcctgt	840
ggcagccctt	ggatgtgctg	aagactcgcc	tgatgaactc	caagggggag	tatcaggggc	900
ttttccactg	gcctgtggag	acagcgaagc	tcgggcctct	ggccttttac	aagggcctcg	960
tcccagctgg	catccgcctc	atccccaca	ccgtgtctac	ttttgtgttt	ctggaacagc	1020
tacgcaaaaa	ctttggcctc	aaagtgccat	cctgaccagc	cgtgggaatg	gctgggctgc	1080
caggccagac	acgctaggtt	cttccaaaga	gtcccaagcc	cagcacctgc	tcctgggggc	1140
acgagctccc	tggcgtggc	cacccatcct	ccgcagcagg	ccctgctgtg	ccccccacct	1200
gctggctgag	ctcctcctgg	cctcgctccc	tctcagctgt	agctgcacca	cccccgctct	1260
ggctaccagg	ctctcccgcc	tgggcactgc	gtggccttgc	ccctctcccc	ctggcagctc	1320
ctcaggggaa	caggggctac	cagaggctga	tttctcccc	ctcctggggc	aggggagggg	1380
tattatccct	gcctcctgcc	cccgatgccc	aaagcagcat	cttccagcac	tttccatcga	1440
ggacttgggt	ggcagagtgt	gggtgcagcc	tggctgttgc	tcacccaagt	gctagctctg	1500
cacttcgtgt	ctgctgagag	caaccagacc	ttccatgtcc	tcgggcagct	gcaactcccc	1560
gcgagacccc	cgactgggtg	gggatgaaca	agcaacgcag	accacaagcg	agtgcctggg	1620
agggagtggc	ccaggggtgt	tctggagcca	ttgtgggtga	gggtcgaggg	ccacctgagc	1680
cccgcgcacc	gctgcctgcc	ctgcagtggc	tttaacagtt	agttttgcca	aagcctctcc	1740
actcaccagc	aggcggtctc	tgtcttcagg	gatttgtcct	gcgtccctcg	ggcacctggg	1800
cccccccgc	tggctccctg	ggggaatggc	ccaggcgggc	cgcgggttct	ccttagggcc	1860
ttctccccga	caaggagtcc	gacggggcgg	atgctgcata	ctctgcctcc	ctgggtcgtg	1920
ggcttcaccc	cacctgggaa	gggcagtgtg	ctctgtgggg	gctgcaatca	ataaatgccg	1980
ggagctgcca	aaaaaaaaaa	aaaa				2004

&lt;210&gt; 80

&lt;211&gt; 3555

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3295764CB1

&lt;400&gt; 80

gaccgtgggc	gagtccagaa	cgctcctggcc	ttacaggagg	aaggcgctcac	tcgcggttac	60
aagtgcctga	ccctcactcc	agttggcgga	ggaggagaa	gaagggggccg	ggccgggttc	120
cctccccctg	cgccccggat	ggatgtgccc	ggcccggtgt	ctcggcgggc	ggcgcgggcg	180
cgggccactg	tgctcctgcg	gaccgctcgg	gtccgctcgg	aatgctgggt	cttgccgacc	240
gcgctgctct	gcgcctacgg	cttcttcgcc	agcttcaggc	cgtccgagcc	cttccctgac	300
ccgtacctgc	tggggccgga	caagaacctg	accgagaggg	aggtcttcaa	tgaattttat	360
ccagtatgga	cttactctta	cctgggtgcta	ctgtttcctg	tgttccttgc	cacagactac	420
ctccgttata	aacctgttgt	tctactgcag	gggctcagcc	ttattgttac	atggtttatg	480
ctgctctatg	cccagggact	gctggccatt	caatttctag	aattttttta	tggcatcgcc	540
acagccactg	aaattgccta	ttactcttat	atctacagtg	tgggtggacct	gggcatgtac	600
cagaaagtca	caagttactg	tcgaagtgcc	actttgggtg	gctttacagt	gggctctgtc	660
ctagggcaaa	tccttgtctc	agtggcaggc	tggtcgctgt	tcagcctgaa	tgctatctct	720
cttacctgtg	tttcagtggc	gttctgtgtg	gcctgggttt	tacctatgcc	acagaagagc	780
ctcttctttc	accacattcc	ttctacctgc	cagagagtga	atggcatcaa	ggtacaaaat	840
ggtggcattg	ttactgacac	cccagcttct	aaccaccttc	ctggctggga	ggacatttag	900
tcacaaatcc	ctctaaatat	ggaggagcct	cccgtggagg	aaccggaacc	caagccagac	960
cgtctccttg	tattgaaagt	actatggaat	gatttctctg	tggtctactc	ctctcgccct	1020
cttctctgct	ggtctgtgtg	gtgggcccct	tctacctgtg	gctattttca	agttgtgaac	1080
tacacacagg	gcctgtggga	gaaagtgatg	ccttctcgct	atgctgctat	ctataatggt	1140
ggcggtggagg	ccgtttcaac	cttactgggt	gctgttgtct	tgtttgagct	tggttatata	1200
aaaatatcct	ggtcaacttg	gggagaaatg	acattatctc	tcctttctct	cctgattgct	1260
gctgcagtgt	atatcatgga	cactgtgggt	aacatttggg	tgtgctatgc	atcctatggt	1320
gtcttcagaa	tcactacat	gttactcatc	acgatagcaa	cttttcaaat	tgtgtcaaac	1380
ctcagctatg	aacgctatgc	cctagtattt	ggtgtaaata	ccttcattgc	cctggcagct	1440
cagacgctgc	tcactcta	tgtggttagat	ggcagtgggc	ttggattaga	aattaccact	1500
cagtttttga	tctatgccag	ttattttgca	ctcatcgctg	tggttttctc	ggccagtggt	1560

```

gcagtcagtg ttatgaagaa atgtagaag ctggaagatc cacaatcaag ttctcaagta 1620
accacttcat aatatactgc tgaagggtt ctctcttatag caagaactct gcacagcaac 1680
tgcttgatg tatttgattt ttaaagcgt agacatatat ttatgaatgt gcatttcttg 1740
acttcacagc agccacttga ctaatacctt gtgttccggg aataacatga tactattcag 1800
aggagccaga agtaaaagttt atttcattga ttatttatga gagctaattt aaggatgact 1860
ttttttctga ttcaaaagtg aacttgattt taaaaaccag tcaagagcaa tcaaagcagc 1920
acatgggtgt gtatacttca ttagcaagtg agtttggtgt tttataggtc acatatgtct 1980
gtatctactt agccagatgc ttggcctggg gggaccaggg ctccacagag gccacaaaat 2040
gttgcaagtc atgatggatg gaaatatgtt ctaacagcat ctgctcttat tcaatttaat 2100
tcttatttct gtgttactca tgtacattgg tcttcttaca tagttattct atcactggca 2160
atatttggtt tgggttagtg ttctgtattt taaggtgtac gtatcatttc taattttaag 2220
ttattttaaa aaaattcatc atatgaatgt tcttggttcc cattgtgacg attatttatt 2280
tctgtaaaat ttgttttagaa gtacgttttt gcattattca tatgcttccc agagaagctc 2340
atttagttag aaaataaggc aagttttgaa gcctgctaaa tgaagagact taagaaagct 2400
taaggtagtc ttgcttgtct ttaaactctt aatatgaagg actattaatt ccaagattaa 2460
aagttcatat ataggctaaa gatgtaacta ggccatttgt atttgtattc ccttttattt 2520
ccaaaataaa atgaaaaatc tttttttaat aatttcatcc ctatttatag tttttatatt 2580
aatttgtttt tcttatccaa gtaaagatgt caataggaat tgcattagtc caaggccttt 2640
ttcataaact gagcctcttt tcaattattt caatgggaca ggaactagga tagatgtgat 2700
tctgcattt ttttacctta aatctgcctt tgtttctaaa ggtagatcat cttgaatatt 2760
tgcttaaaat tgctagtgat ttcattacca agttacttga aaaaatgttc tatatgcatt 2820
taattctgaa atcagctctac caaggggctg ctagtatatg tcagacatga aaactatttt 2880
aaagctgact ttgttgccct atcttgaaaa gaatctagat aggtgctttt aactgggta 2940
ttaacttttt tagaatgaca cagctgaaca gtgttaataa tagtgtgtca agattgcaa 3000
gtcgacatac tcatttggtt taagcaggaa tcttagaagc aaatggatgg ggataaagat 3060
aggctatttt ctattcacca tcttttacta ttaagggaaa ggaaaagaac actagctaag 3120
gaagggaaa ggaagtgata tcataaaagt agcaaccttc attttacatt ctgtctgttt 3180
tctttttttt gctttgtttt gtttggtgta atttgggaat tgtgtactcc gaaacaagta 3240
gaaaagtgtc gtttgaggga ttttatataa ttttttttta atggaatgtg gtacaaattg 3300
ttcatgttac caaagcaata tttccctgga atttaattca aagtttgtgg catacaacct 3360
gagccttttc ttatataaga caagaatatg ttcacatctt ggtatgtggc catatttata 3420
gaatgctgaa ctcaatgtgc aagttgtact gtatgcagtt ttgtaataaa gtgaaaaata 3480
tttgtgtgac tttttattca attctgtata gattataaaa ttatttttat taaataaata 3540
ttttacagta ttttt 3555

```

&lt;210&gt; 81

&lt;211&gt; 1293

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3438320CB1

&lt;400&gt; 81

```

atatgccca cgcgggactc atactacgtt tcccgtgaac acgtgcagtc caaaccccg 60
ccctgatatt tatctcagtg gacggtggcc ggaagagac aatggtttcc atgtcagcgg 120
ataaacgctc tcccctcggc tcccgagcgc gacggaggtc gtagtagtag tgagtacgtg 180
ctgaggagca aaggagtaac caagagatcc agtgaccgac agagcaagag ccatgcccg 240
ccggggcctg gtggctgggc cagacttgga gtattttcag cgctcgctatt tcacgccggc 300
ggaggtggcc caacataaca ggcccgaaga cctctgggta tcttacctgg gacgcgtgta 360
cgacctaacg tcattggcac aggaatacaa ggggaacctg ctgctgaaac ccactctgga 420
agttgcagcg caggatatca gccactgggt tgatccaaag accagagaca tccgcaagca 480
catagatccg ctgaccggct gcctgaggtg ctgcaccccg cggggccgct ttgtgcacgt 540
tccgcctcag ctgccctgtt cggactgggc caacgatttt ggaagccct ggtggcaggg 600
gtcgtattat gaggtggggc ggctgtctgc caagaccgag agcatccgca tcattaacac 660
gctcacgtcg caggagcaca cactggaggt gggggttctg gagtccatat gggaaatcct 720
acaccgctat ctcccctata actcacatgc tgcagctac acgtggaat atgaagggaa 780
gaacctgaac atggatttta ccctggaaga gaatggatc cggtatgagg aggaagaatt 840
tgactatctc agtatggacg gtacacttca cacacctgca atacttctgt acttcaatga 900
tgatctcacg gagttgtagg caaggagatg tacactcgtg tagactcaag acgtatttctg 960
agtttggtct tttctgtgcc ttgagggaaa gtggtggggc cgaggggtgc ctggaccag 1020
atctccactc ctctccagga gctagcctgt gcccttctga agtgtaaaag cctattccc 1080
tgccttcatt acagtttgc ctgagaaaat tagtgaatta atctttggga atgatacaag 1140
aagatcaagt accttggttt agggagatgt agaagaggat agtcagagtt caggcagaac 1200
tgtttgatag ttaagagaga gtatttctac aggggtgagg gatggaagga cttttttggc 1260
aatgatggaa atgagatgtc tgcaggagat ggg 1293

```

<210> 82  
<211> 1489  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 3986488CB1

<400> 82  
cggggcgtggc gggcggcgct cttctacggg acctgctcct tcctcatcgt gcttgtcaac 60  
aaggcgctgc tgaccaccta cggtttcccg tcaccaattt tccttggaat tggacagatg 120  
gcagccacca taatgatact atatgtgtcc aagctaaaca aaatcattca cttccctgat 180  
tttgataaga aaattcctgt aaagctgttt cctctgcctc tcctctacgt tggaaaccac 240  
ataagtggat tatcaagcac aagtaaatta agcctaccga tgttcaccgt gctcaggaaa 300  
ttcaccattc cacttacctt acttctggaa accatcacac ttgggaagca gtattcactc 360  
aacatcatcc tcagtgtctt tgccattatt ctcggggctt tcatagcagc tgggtctgac 420  
cttgctttta acttagaagg ctatatTTTT gtattcctga atgatatctt cacagcagca 480  
aatggagttt ataccaaaca gaaaatggac ccaaaggagc tagggaaata cggagtactt 540  
ttctacaatg cctgcttcat gattatccca actcttatta ttagtgtctc cactggagac 600  
ctgcaacagg ctactgaatt caaccaatgg aagaatgttg tgtttatcct acagtttctt 660  
ctttcctgtt ttttgggggt tctgtgatg tactccacgg ttctgtgcag ctattacaat 720  
tcagccctga cgacagcagt ggttggagcc atcaagaatg tatccgttgc ctacattggg 780  
atattaatcg gtggagacta cattttctct ttgttaaact ttgtagggtt aaatatttgc 840  
atggcagggg gcttgagata ttcttttta acactgagca gccagttaaa acctaaacct 900  
gtgggtgaag aaaacatctg tttggatttg aagagctaaa gagtctgcag caggattgga 960  
gactgacttg tgactgcggg ctgggggggc attcccagta ggaatgtgaa gccagaggtt 1020  
tcgggttcgt gacatccacc ccctgggcaa gtgagagcat ctgcaaaatg caaagagAAC 1080  
tacctcatat gcaggatgag ccaatggcag tctcaagaaa tgtactcggg cgacacctta 1140  
cctgtggaAA gcaaattctt tcaaaataag ccactgggac tcggtaggtg gagcccagc 1200  
tgctcttcta gggacctatg gggccttctg ggcattctct tgctgtgtgc tggggaggag 1260  
gttgatgtaa tgggtgactct tttctgatca gcaccttggc cgtgattccc aagggtccag 1320  
ccaaagcaaa gggccagtgt tttcagttta aacagacatg tctttagtct aataaaatta 1380  
gttaactgcc agtaaagtta tttgttagct ttgatgaaag ctatgttggt atctttccct 1440  
aatcatcaaa gtaaataaaa aatcatttct atgtaaaaaa aaaaaaaaaa 1489

<210> 83  
<211> 927  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 4378816CB1

<400> 83  
ctccctgctt tcctctgccg catggctcctg ggccgttggc gtcggaagcc tgaagcatgg 60  
gcgctgagtg ggagctgggg gccgaggctg gcggttcgct gctgctgtgc gccgcgctgc 120  
tggcggcggg ctgcgccctg ggcctgcgcc tgggcgcggg gcagggggcg gcggaccgag 180  
gggcgctcat ctggctctgc tacgacgcgc tgggtgactt cgcgctggaa ggcccttttg 240  
tctacttgct tttagtagga aacgttgcaa attccgatgg cttgattgct tctttatgga 300  
aagaatatgg caaagctgat gcaagatggg tttattttga tccaaccatt gtgtctgtgg 360  
aaattctgac cgtcgccctg gatgggtctc tggcattgtt cctcatttat gccatagtca 420  
aagaaaaata ttaccggcat ttctctcaga tcaccctgtg cgtgtgcgag ctgtatggct 480  
gctggatgac cttctcccca gagtggctca ccagaagccc caacctcaac accagcaact 540  
ggctgtactg ttggctttac ctgtttttt ttaacgggtg gtgggttctg atcccaggac 600  
tgctactgtg gcagtcatgg ctagaactca agaaaatgca tcagaaagaa accagtccag 660  
tgaagaagtt tcagtgaact ttcaaaacca taaacaccat tatctaactt catgaaccag 720  
aatgaatcaa atctttttgt ttggccaaaa tgtaatacat tccagtctac actttgtttt 780  
tgtattgttg ctctgaaca acctgtttca aattggtttt aaggcgacca gttttcgttg 840  
tattgttggt caattaaatg gtgatatagg gaaaagagaa caaatttgaa tttgtaataa 900  
taaaatgttt aattatacaa aaaaaaa 927

<210> 84  
<211> 970  
<212> DNA  
<213> Homo sapiens



<220>  
<221> misc\_feature  
<223> Incyte ID No: 4797137CB1

<400> 84  
ggatgcagca gagaggagca gctggaagcc gtggctgcgc tctcttccct ctgctgggcg 60  
tcctgttctt ccagggtgtt .tatatcgtct tttccttgga gattcgtgca gatgcccattg 120  
tccgagggtta tggttgagaa aagatcaagt tgaatgcac tttcaagtca acttcagatg 180  
tcaactgacaa gcttactata gactggacat atcgccctcc cagcagcagc cacacagtat 240  
caatattttca ttatcagtct ttccagtacc caaccacagc aggcacattt cgggacgga 300  
tttcctgggt tggaaatgta tacaaagggg atgcacttat aagtataagc aacctacca 360  
taaaggacaa tgggacattc agctgtgctg tgaagaatcc cccagatgtg caccataata 420  
ttcccattgac agagctaaca gtcacagaaa ggggttttgg caccatgctt tcctctgtgg 480  
cccttctttc cctcctgtgc tttgtgccct cagccgtggg ggttgctctg ctgctgggtga 540  
gaatggggag gaaggctgct gggctgaaga agaggagcag gtctggctat aagaagctcat 600  
ctattgaggt ttccgatgac actgatcagg aggaggaaga ggcgtgtatg gcgaggcttt 660  
gtgtccgttg cgctgagtgc ctggattcag actatgaaga gacatattga tgaaagtctg 720  
tatgacacaa gaagagtcac ctaaagacag attccactgg cagctaaagc 780  
ctgtcagaga aagtggagct ggcctggacc atagcgatgg acaatcctgg agatcatcag 840  
taaagacttt aggaaccact tatttattga ataaatgttc ttgttgatt tataaactgt 900  
tcagggaagtc tcataagaga ctcatgactt cccctttcaa tgaattatgc tgtaattgaa 960  
tgaagaaatc 970

<210> 85  
<211> 594  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5470806CB1

<400> 85  
gacaggatgg cttcccttcg cctgttccct ctctgcctcg ctgtactgat atttgcgctt 60  
gaagctggcc ctgggggtgc tggagaatcc aagtgtcctc tgatgggtcaa agtcctggat 120  
gctgtccgag gcagccctgc tgttgatgtg gccgtgaaag tgttcaaaaa gactgcagac 180  
ggaagctggg agccgtttgc ctctgggaag accgccgagt ctggagagct gcacgggctc 240  
accacagatg agaagttcac ggaaggggtg tacagggtag aactggacac caaatcgtac 300  
tggaaaggctc ttggcatttc cccattccat gaatacgcag aggtggtttt cacagccaat 360  
gactctgggc atcgccacta caccatcgca gcctgtctca gcccgtactc ctacagcacc 420  
actgctgtcg tcagtaacct ccagaactga gggacccagc ccagtaggac caggatcttg 480  
ccaaagcagt agcttcccat ttgtactgaa acagtgttct tgctctataa accgtgttag 540  
caactcggga agatgccgtg aaacgatctt attaaaccac ctgtgatgcc aaaa 594

<210> 86  
<211> 618  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5473242CB1

<400> 86  
gtgttgactc gcaacctcag gaacagacac catggtgcac ctaactgatg ctgagaaggc 60  
tactgttaat ggcctgtggg gaaaggtgaa cctgtgtgaa attggcgctg agtcccttgc 120  
cagtctgctg attgtctacc cttggaccca gaggtacttt tctaaatttg gggacctgtc 180  
ctctgtctct gctatcatgg gtaaccccca ggtgaaggcc catggcgaaa aggtgataaa 240  
cgcttccgat gatggcctga aacacttgga caacctcaag ggcacctttg ccagcctcag 300  
tgaactccac tgtgacaagc tgcattgtga tcctgagaaac ttcaggctcc tgggcaatat 360  
gatttgtgatt atgatgggcc accacctggg caaggaattc accccgagtg cacaggctgc 420  
cttccagaag gtgggtggctg gagtggccag tgccttggtc cacaagtacc actaaacctc 480  
ttttcctgct cttgtctttg tgcaatggtc aattgttccc aagagagctt ctgtcagttg 540  
ttgtcaaaat gacaaagacc tttgaaaatc tgtcctacta attaaagcat ttggttcaag 600  
tggtctgttg agataccc 618